



Effect of Conservation Tillage and Precision Nitrogen Management on Wheat: A Review

Aditya Shukla ^{a*}, Mukesh Kumar ^a and Akanksha Shukla ^b

^a Department of Agronomy, Sardar Vallabhbhai Patel University of Agriculture, Meerut, India.

^b Department of Plant Pathology, Indira Gandhi Krishi Vishwavidyalaya, Raipur, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2022/v34i2131245

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/89523>

Review Article

Received 02 May 2022
Accepted 08 July 2022
Published 09 July 2022

ABSTRACT

The most significant source of protein in the world is wheat, which accounts for roughly 21% of daily dietary protein requirements on average. With more than 60% of the population living in rural regions, the wheat crop system in Indian agriculture represents the backbone and the future of the country's economy. By speeding the oxidation and disintegration of organic matter, which results in the deterioration of soil characteristics, intensive ploughing causes a decrease in soil organic matter. To conserve resources, slow down soil degradation, adjust cropping systems to climate extremes, and increase agricultural sustainability over the long term, conservation agriculture (CA), which involves minimising soil disturbances, is widely supported. One of the greatest restrictions for CA is the immobilisation of N, which is mostly related to the presence of a continuous residue cover on the soil surface. The employment of some technologies, such as the SPAD metre, NDVI sensor, site-specific nutrient management through soil test crop response, nutrient expert LCC, etc., during the growing season for precision N management aids in meeting the crop's nitrogen needs with the least amount of environmental impact. This study aims to examine the research studies where precision nitrogen management technologies were applied in conservation agriculture and discuss the effects of these tools and techniques on wheat growth, productivity, and nutrient usage efficiency (NUE).

Keywords: Conservation tillage; Leaf color chart; nutrient use efficiency; precision nitrogen management; SPAD meter.

1. INTRODUCTION

The most significant staple food for around 36% of the world's population is wheat (*Triticum aestivum* L.), sometimes known as the king of cereals. It is the most significant and strategically important cereal crop. Wheat provides around 18% of the daily calorie needs in underdeveloped nations. The most significant source of protein in the world is wheat, which accounts for roughly 21% of daily dietary protein requirements on average [1]. With more than 60% of the population living in rural regions, the wheat crop system in Indian agriculture represents the backbone and the future of the country's economy.

In India, the crop helps support a vast number of people's livelihoods. It is mostly seeded using heavy tillage farming techniques. Traditional/conventional farming practises have a number of drawbacks, including the deterioration and compaction of the soil, the depletion of water supplies, and the loss of biodiversity. Water is becoming increasingly limited in the western Indo-Gangetic Plains (IGP) as a result of increased competition from the urban and industrial sectors [2]. Overexploitation and poor groundwater management have had detrimental effects on the ecology and the water table in several areas of the region [3]. Other widespread issues in the area include excessive residue burning and degradation of land quality brought on by various types of soil degradation [4]. These variables prompt the idea of conservation agriculture (CA) to be taken into account for long-term productivity, financial success, and soil quality [5]. Three principles govern CA: (i) reducing mechanical soil disturbance; (ii) using cover crops and/or crop residues (primarily residue retention); and (iii) diversifying crop types in associations, sequences, and rotations to increase system resilience. According to reports, CA increases input usage effectiveness, soil organic carbon (SOC) content, and has the ability to lower greenhouse gas emissions [6].

In 2021, the last year for which we have official production statistics, the world produced an estimated 776.5 million tonnes of wheat with China, India, and Russia accounting for 38% of global production. When India gained independence in 1947, wheat production and

productivity were extremely low. In 1950–51, just 6.46 million tonnes of wheat were produced, and the productivity was only 663 kg per hectare, which was insufficient to feed the Indian population. Under PL-480, the nation used to import a lot of wheat to meet the demands of its citizens from various nations, including the USA. The tall plant habit that causes lodging when grown in fertile soils, the poor tillering and low sink capacity of the varieties used, the higher susceptibility to diseases, the higher sensitivity to thermo & photo variations, etc., resulting in poor adaptability, and (e) the longer crop duration that results in a long exposure of plants to the climatic variations and insect pests were the reasons for the low production and productivity of wheat at that time. In 1961, the Indian government established a commission to examine the viability of raising crop output given the country's current ecological circumstances. The wheat situation in our nation has entirely changed as a result of several actions the Indian government has taken. Our nation used to import wheat to meet our needs throughout the post-Independence era, but when the "Green Revolution" of the late 1960s saw a dramatic increase in wheat production and productivity, our nation was able to produce enough wheat to meet its own needs. Currently, the nation produces more wheat than is needed, and godowns are overflowing with it.

Today National wheat production is 109.52 Mt with an acreage of 31.6 million hectare and productivity of 3.46 t ha⁻¹. The top 3 producing states are Uttar Pradesh, Madhya Pradesh, Punjab. U.P. is a top producing state in India with 35.5 Mt of production and contributing 32% of total national production.

1.1 Conservation Agriculture

Three crop management principles, namely crop rotation, surface residue retention, and minimal or no-tillage (less soil disturbance), make up the majority of conservation agriculture (CA). The impact of CA on agricultural production components and its use from diverse farming vantage points are hotly contested topics worldwide. The current crop production methods, which involve repetitive tillage and the removal of straw, cause soil compaction and surface crusting, which decrease water infiltration and increase soil erosion, ultimately leading to a

general decline in the physical health of the soil. For sustainable crop production, effective use of natural resources, and improved response to additional inputs, it is necessary to maintain soil physical health at its optimum level. Numerous advantages have been linked to conservation agriculture and its components, such as increased soil water storage [7,8,9], improved soil quality [10,11], reduced erosion [12], and in some cases, higher yield and net farm income [13,14,9].

Initially developed as a response to the US Dust Bowl, conservation agriculture [15]. Since then, CA has gained popularity quickly, especially in North America, South America, and Australia [16]. It is mostly used on large, automated farms and calls for heavy doses of pesticides to combat weeds that are typically managed by tillage. In South Asia and Sub-Saharan Africa, there are currently coordinated efforts to promote CA in smallholder systems [17,18]. Based on the available evidence, it is uncertain whether CA, which was developed for high-input systems in more temperate locations, can function and produce profits in smallholder systems in the tropics and subtropics. This issue requires additional research.

1.2 Constraints for the Adoption of Conservation Agriculture

To change the attitudes of farmers, it is vital for farmers, technicians, extensionists, and researchers to shift their thinking away from soil-degrading tillage practises and toward sustainable production techniques like no-tillage (Derpsch, 2001). Hobbs and Govaerts [19] pointed out that overcoming the bias or mindset about tillage is likely the most crucial element in the implementation of CA. It is argued that one of the biggest obstacles to adopting CA widely is persuading farmers that effective cultivation is still achievable with reduced-tillage or without tillage. In many situations, it might be challenging to persuade the farmers of the potential advantages of CA outside of its potential to lower production costs, primarily by reducing tillage. CA is currently regarded as a path toward sustainable agriculture. The spread of conservation agriculture, therefore, will call for scientific research linked with development efforts.

The following are a few important constraints that impede the broad-scale adoption of CA.

- A lack of suitable seeders, particularly for small and medium-sized farmers: Although significant efforts have been made to develop and promote machinery for seeding wheat in no-till systems, successful adoption will require accelerated effort to develop, standardise, and promote quality machinery aimed at a variety of crops and cropping sequences. These would involve managing crop wastes during harvest and creating permanent bed and furrow planting techniques.
- The extensive utilisation of agricultural wastes for animal feed and fuel: Farmers, particularly in rainfed environments, experience a shortage of crop residues due to lower biomass production of various crops. For agricultural residue, there is rivalry between CA practise and livestock feeding. This poses a significant obstacle to the promotion of CA in rainfed environments.
- Burning agricultural residues: Farmers prefer to seed the crop in a timely manner by burning the residue rather than using technology for sowing under CA systems. The rice-wheat system in north India now frequently includes something like this. For the area, this causes environmental issues.
- Lack of awareness among agricultural leaders, extension agents, and farmers of the potential of CA: This suggests that the full spectrum of conservation agriculture practises, such as planting and harvesting, managing water and nutrients, controlling diseases and pests, etc., must be developed, assessed, and matched in the context of new systems.
- Skilled and scientific labour: Managing conservation agricultural systems would necessitate improved scientific capacity to approach issues from a systems viewpoint and be able to collaborate closely with farmers and other stakeholders. Mechanisms for sharing knowledge and information must be strengthened.

1.3 Precision Nutrient Management

According to specific field circumstances throughout a given cropping season, precise nutrient management is a science-based method for supplying nutrients to crops as and when they are required. Precision nutrient management is the science of managing spatial and temporal

variability in the inherent nutrient supply from soil to increase productivity, efficiency, and profitability of agricultural production systems. It involves the use of cutting-edge, innovative, and site-specific technologies. Understanding the geographic variability of soils is necessary (Jin and Jiang, 2002).

The 4 R's of precise nutrient management are

- Right Source
- Right Rate
- Right Time
- Right Place

Nitrogen is the nutrient that is most frequently and carelessly used in crop cultivation. Precision nitrogen management techniques can be developed using the spectral characteristics of plant leaves as an index since the dynamics of nitrogen supply to plants control the concentration of chlorophyll in plants. In order to make precise decisions about nutrient management while taking into account spatial and temporal variability in nutrient supply from the inherent sources, other techniques are being used as the plant demand for nutrients other than nitrogen cannot be easily accessed from the spectral properties of the leaves. Recent developments suggest that geospatial technologies like the global positioning system (GPS), geographical information system (GIS), remote sensing, real-time and/or variable rate applications (VRA) can assure need-based nutrient management in crop fields [20,21,22]. By addressing the issue of over- and underfertilization, the need-based variable-rate fertiliser delivery method offers a significant chance of increasing the efficiency of fertiliser use [23].

1.4 Precision Nitrogen Management Tools and Techniques

1.4.1 Optical sensors

There are many optical sensors that fall under the multispectral and hyperspectral categories. In contrast to hyperspectral sensors like ASD FieldSpec (350-2500 nm), which have a fine spectral resolution (1-2 nm) and continuous wavebands (2150) across the electromagnetic spectrum, multispectral sensors like Crop Circle (450-880 nm) and CropScan (440⁻¹750 nm) have a wide spectral resolution (10 to 20 nm) and a limited number of wavebands (3 to 16) used to describe nitrogen, biomass variation, Utilizing

spectral indices computed using univariate and multivariate regression techniques, the spectral reflectance data can be interpreted.

1.4.2 Chlorophyll meters

As trustworthy substitutes for conventional tissue analysis as plant N nutritional diagnosis techniques, chlorophyll metres are available. The handheld Minolta SPAD-502 chlorophyll metre is the most popular model. A rapid, non-destructive hand-held instrument called the SPAD 502 chlorophyll metre (Soil-Plant Analysis Development) was created by Minolta in Osaka, Japan. By utilising two LEDs (light-emitting diodes) that emit red (=650 nm) and infrared (= 940 nm) light, the unplucked leafy tissue is clamped in the metre, instantly providing an estimate of leaf N status as chlorophyll concentration [24]. The leaf is constructed to allow the red and infrared radiations to flow through. A silicon photodiode detector transforms the light's transmission through the leaf and partial absorption into an electrical signal.

1.4.2 Leaf Colour chart (LCC)

A high-quality plastic strip with several shades of green, from light yellowish green to dark green, is called a leaf colour chart. LCC technology was reportedly used in Japan, according to Furuya [25]. The International Rice Research Institute (IRRI) collaborated with the agricultural research systems of various Asian nations to develop an upgraded version of the six-panel LCC (IRRI-LCC, six-panel).

1.4.3 Nutrient management models

In general, computer-based decision support systems for precise nutrient management in crop production use the Nutrient Expert (NE) and QUEFTS models. The models are developed to ensure need-based nutrient treatments while taking into account regional and temporal variability in nutrient supply. The nutrient expert (NE) creates farmers' unique fertiliser recommendations based on information about the growing environment, residue content, organic and inorganic fertilisers used, achievable yield, and historical yield data going back three to five years. It considers resource availability when estimating their yield target. Site-specific nutrient management (SSNM) standards were used to construct the algorithm for determining fertiliser requirements in Nutrient Expert. It is a highly interactive computer-based instrument that

provides quick information on a field's fertiliser needs.

1.4.4 Omission Plot Technique (OPT)

The omission plot approach is used to calculate the amount of fertiliser needed to get a target yield. With the exception of the nutrient of interest, which is the missing nutrient, all the major nutrients are used in this procedure. The method gives an approximation of the soil's natural supply of nutrients. The yield will be constrained by the local supply of P, for instance, if all the nutrients are applied to the P-omission plot except for P. The amount of fertiliser needed is then determined by comparing the yield gap between the maximum yield possible and the yield obtained using the omission plot technique.

2. EFFECT ON GROWTH

When it comes to precision N-management techniques and tillage crop establishment procedures, furrow irrigated raised bed (FIRB) and N 80-LCC measured wheat plant height at harvest significantly higher than conventional tillage and farmer fertilisation approaches. Similar to how conventional tillage and farmer fertilisation approaches recorded the lowest dry-matter accumulation at 90 DAS of wheat crop, FIRB and N-80 LCC recorded significantly the highest. When wheat was fertilised with N-80 LCC and elevated beds that were irrigated by furrows, the number of tillers increased

dramatically at 90 DAS. Wheat plants gradually grew taller till they were fully mature [26]. Similar results were also reported by Biradar [27], and Sulochna et al. [28], where, precision nutrient management treatments recorded maximum plant height of wheat than farmers fertilizer practice as balanced application of nutrients as per crop need under N 80- LCC enhances the nutrient use efficiency.

Budhar [29] stated that a consistent supply of N supplied at the seedling stage served to generate a favourable influence on growth features, which may be the cause of the higher plant height and greater number of tillers at LCC 4 and 5. With higher LCC values of N application ranging from 3 to 5, there has been a progressive and noticeable rise in tiller output at 80 DAS. When it came to producing tillers per hill among the LCC values used for N application, LCC 5 stood out as being superior, especially at the last stages of growth at 60 and 80 DAS when the tiller count at LCC 5 was the greatest and much greater than LCC 3 and 4. According to Mohanty et al. [30], the maximum NAR was produced by STCR and absolute control at 0-30 and 60-90 DAS, respectively, for precision N-management strategies in wheat. However, at 30–60 DAS, various precision N-management techniques had little to no impact on NAR. It was abundantly obvious that the direct and indirect effects of enhanced nutrition gave the crop an initial boost, increasing RGR and NAR under conservation agriculture.

Table 1. Effect of tillage crop establishment methods and precision nitrogen management on plant height, dry-matter accumulation and tillers m⁻² of wheat crop

Treatments	Plant height at harvest (cm.)	Dry matter accumulation at 90 DAS (g m ⁻²)	Tiller m ⁻² at 90 DAS
Tillage crop establishment methods (TCM)			
T ₁ – Zero tillage	97.1	846.9	352.7
T ₂ – Roto Tillage	93.3	840.9	333.3
T ₃ – Raised bed	104.7	871.5	368.3
T ₄ – Conventional tillage	89.1	822.9	339.9
C.D. (P=0.05)	1.7	14.5	9.8
Precision nitrogen management			
F ₁ - N 80:20	95.6	821.1	339.2
F ₂ – Nutrient expert 33:33:33	98.1	864.5	360.1
F ₃ - N 80-LCC	100.1	916.9	373.0
F ₄ - Recommended dose of N	93.9	819.9	343.1
F ₅ - FFP	92.5	805.5	327.5
C.D. (P=0.05)	1.2	8.5	6.9

Source: Gawdiya, 2020

3. EFFECT ON YIELD

The amount of crop dry matter accumulated, the number of tillers, the number of grains spike⁻¹, and the grain weight, among other factors, all affect grain output. The most crucial factor for assessing the results of applied treatments is grain yield. Crop productivity is the pace at which a crop accumulates biomass and is primarily dependent on photosynthesis, which is the process by which green plants transform light energy into chemical energy.

Mahajan [31] reported that (FIRB) significantly outperformed all other treatments in terms of maximum grain output as seen in Table 2 (52.6 q ha⁻¹). Nevertheless, treatment T1 (wheat sown with no tillage) outperformed the other treatments. During the experiment, T5 outperformed reduced (T2) and roto tillage (T3). When compared to T4 (wheat sown on FIRB) and T5 (wheat sown by conventional tillage), the decline in grain production caused by reduced tillage and roto tillage procedures was 24.6 and 9.1 percent, respectively. However, compared to the other tillage practise plots, the yield of wheat planted in raised beds with furrow irrigation increased by 18.4%.

According to Biradar et al. [27], fertiliser application based on SSNM principles produced noticeably higher grain yields than farmer practise (FP) and the required amount of fertiliser (RDF). Averaging 3.7 t ha⁻¹, or 23 percent greater than the RDF and 39 percent more than the FP, was the average wheat yield in the SSNM practise. According to Khurana et al. [32], a N management schedule that took real-time variation in crop N requirement at critical wheat growth stages into consideration. Average grain production increased from 4.2 to 4.8 t ha⁻¹ as compared to current farmers' fertiliser practises (FFP), while plant N, P, and K accumulations increased by 12 to 20% when using SSNM.

Singh et al. [33] reported from Meerut that the mean grain yield of wheat was 5.30 t ha⁻¹, 4.47 t ha⁻¹ and 3.8 t ha⁻¹ under OPT, SR and FFP, respectively. On an average, the OPT out-yielded the SR and FFP by 0.82 t ha⁻¹ and 1.5 t ha⁻¹ or by 18% and 39%, respectively. According

to Hussain et al. [34], the crucial SPAD value of 37.5 is suitable for directing Pakistan's need-based N top-dressing of rice. By utilising SPAD value 35 as the threshold SPAD value in Bangladesh, Kyaw et al. [35] produced noticeably greater yields with 3 to 12 percent less fertiliser-N use than the general advice. In comparison to a blanket dose of 150 kg N ha⁻¹, Maiti et al. [36] discovered that a SPAD value of 37 is adequate for achieving high yields with 27.5 to 45.5 kg N ha⁻¹ less fertiliser. When managing N in direct wet-seeded rice, Huan et al. [37] used 32 and 29 SPAD threshold values. The grain yield from the SPAD 32 and 29 based N applications was 20 and 40 kg N ha⁻¹ less, respectively, than the blanket recommendation for fertiliser N, which is 120 kg N ha⁻¹. For various varietal groups, it can also be necessary to employ different thresholds for SPAD values [38,39]. According to Takebe et al. [40], there is a strong correlation between grain protein concentration at harvest and leaf colour at the full heading stage. A grain protein content of more than 120 g kg⁻¹ was achieved with the application of 30 kilogramme N ha⁻¹ (if SPAD value was 50–52) or 60 kg N ha⁻¹ (if SPAD value was 45–50) during the full heading stage.

According to research by Uppal et al. [41] on the influence of NPK levels on the development and production of durum wheat, grain yield rose with each increment of N by 40 kg/ha up to 200 kg ha⁻¹. Tayebbeh et al. [42] conducted research on the impact of organic and inorganic fertilisers on wheat grain production and protein banding pattern, and they found that the plants produced the highest grain yield when fertilised with 160 kg N and 30 t compost per hectare. Spikes/plant, seeds/spike, and 1000 kernels weight among yield components all increased significantly as nitrogen levels rose.

Alam et al. [43] found that LCC-based N management consistently increased grain yield production across two rice seasons compared to farmers' fertiliser practises. Depending on the village and season, the LCC-based N management increased grain output by 0.1 to 0.7 t ha⁻¹. In addition to the prescribed P, K, S, and Zn fertilisation, the use of LCC for regulating N enhanced mean yield by 0.3–0.4 t ha⁻¹ [44].

Table 2. Effect of tillage crop establishment methods and nitrogen management on yield and test weight of wheat

Treatments	Grain yield (quintal ha ⁻¹)	Straw yield (quintal ha ⁻¹)	Test weight (g)	Harvest Index (%)
Tillage Crop establishment methods (TCM)				
T ₁ ZT	48.7	57.5	41.2	45.8
T ₂ Reduced tillage	41.9	50.9	39.6	45.2
T ₃ Roto tillage	37.4	44.8	37.4	45.0
T ₄ Raised bed	52.6	63.8	42.3	45.2
T ₅ Conventional tillage	43.6	56.6	40.3	43.5
SEm±	0.51	0.53	0.4	0.45
C D (P=0.05)	1.68	1.75	1.31	1.45
Precision N management				
F ₁ Control	33.8	45	36.5	42.9
F ₂ RDF (150:60:40)	45.4	55.3	40.3	45.1
F ₃ SPAD	47.1	58.2	41.4	44.7
F ₄ Targeted yield	51.7	59.4	41.6	46.5
F ₅ LCC	46.2	55.7	40.9	45.4
SEm±	0.78	1.01	0.65	0.74
C D (P=0.05)	2.22	2.93	1.87	2.13

Source: Mahajan (2018)

4. EFFECT ON NUTRIENT USE EFFICIENCY (NUE)

Ghosh et al. [45], reported that, wheat's Agronomic N use efficiency (AEN), N recovery efficiency (REN), internal N use efficiency (IEN) and partial factor productivity of applied N (PFPN) were significantly impacted by N management practises (Table 3). The maximum AEN and REN were obtained with medium rate N application at medium SPAD level (N25S40), which was higher than those obtained with high (N35) and low (N15) rate N topdressing at high (S42) and low SPAD (S38) levels, as well as FTNM and FFP. In comparison to low rate of N topdressing at low SPAD level (N15S38), high rate of N topdressing at high SPAD level (N35S42), and FTNM, N25S40 increased AEN and REN by 29.2 and 50.9 percent, 50 and 8.8 percent, and 58.5 and 15.1 percent, respectively. Regardless of SPAD levels, increasing the rate of N topdressing in wheat caused the IEN and PFPN to gradually drop. As a result, moderate (N25) and low (N15) rates of N topdressing at medium (S40) and low (S38) SPAD levels respectively resulted in significantly higher IEN and PFPN than high rate of N topdressing at high SPAD level (N35S42). FTNM also considerably reduced IEN and PFPN as compared to low to moderate (N15-25) rates of N topdressing at low to medium SPAD levels (S38-40). Moderate rate of N topdressing at medium SPAD level (N25S40) caused higher IEN (+19.1%) and PFPN (+41.5%) than FTNM.

The harvest index for potassium, phosphorus, and nitrogen was considerably higher in SSNM and significantly lower in absolute control. Absolute control yielded significantly higher harvest indices of nitrogen and phosphorus among the various N precision management techniques, however the harvest index of potassium was not significantly impacted. Regardless of the wheat treatments used, the nutrient harvest index was in the order of P>N>K, indicating that more P was ingested in the grain than K was in the straw [46].

According to Mahajan [31], ZTW (T1) treatments had the highest nitrogen, phosphorus, and potassium harvest index, while FIRB (T4) treatments had the lowest. Absolute control (F1) produced a significantly greater nitrogen and phosphorus harvest index across the various precision N management techniques, while the harvest index of potassium was not significantly impacted. The order of P, N, and K in the nutrient harvest index was highest to lowest. The ability of the soil to give nutrients and the plant's innate ability to use nutrients are both demonstrated by the recovery of any added nutrients. It also depends on how it is applied and the environment in which it grows. Different tillage techniques and precise N management techniques dramatically affected apparent nutrient recovery in wheat. Due to a high amount of potassium in the soil and lower potassium fertiliser doses, the crop recovered more than 100% of the potassium that was applied [47-49].

Table 3. Effect of nitrogen management practices on nutrient use efficiency

Treatment	N rate (kg ha ⁻¹)	REN (kg kg ⁻¹)	AEN (kg kg ⁻¹)	PFPN (kg kg ⁻¹)	IEN (kgkg ⁻¹)
S38N15	70	0.53	19.5	46.5	49.1
S38N25	68.3	0.70	20.5	46.5	42.4
S38N35	95	0.69	17.6	38.5	38.6
S40N15	90	0.59	21.4	45.1	46.7
S40N25	101.7	0.80	25.2	46.6	40.9
S40N35	118.3	0.70	18.3	36.1	36.9
S42N15	105	0.60	21.1	42.2	45.7
S42N25	110	0.71	19.4	38.3	38.4
S42N35	118.3	0.74	16.8	34.9	34.7
FTNM	120	0.70	15.9	32.9	34.3
FFP	80	0.69	20.2	45.8	42.3
Control	0	–	–	–	66.7
SEm (±)		0.06	1.8	2.1	2.1
LSD (0.05)		0.17	5.2	6.2	6.3

Source: Ghosh et al [44]

Different tillage methods had a substantial impact on potassium apparent recovery, but FIRB had the highest potassium apparent recovery, followed by zero tillage plots. This was because FIRB produced significantly higher nitrogen and phosphorus apparent recovery. Since all of these nutrients have a synergistic effect on one another, different tillage techniques increased the availability of nutrients for FIRB, resulting in a much higher apparent recovery (percent of nutrients applied). Targeted yield-based nitrogen management among precision N management techniques produced much higher apparent recovery (N, P & K) [50-53].

5. CONCLUSION

Farm management must employ more environmentally friendly agricultural practises if it is to increase crop output efficiency and lower environmental dangers associated with agriculture. These requirements can be met through precision agriculture. As is commonly agreed, it can assist farmers in applying the proper input in the proper quantity, at the proper location, at the proper time, and in the proper manner. When making decisions on the production of wheat crops, the within-field variability is the main source of uncertainty. At both the geographical and temporal scales, variability must be understood and controlled. In order to produce variable wheat N fertilisation, novel experimental methods, distant and proximal sensing, and crop simulation models will increasingly be used to analyse field variability at a cheap cost.

Looking at the research findings we can conclude that when it comes to growth, productivity and NUE wheat grown in FIRB performs better while economically speaking zero tillage fared better than other tillage methods. Precise nitrogen management involving LCC, SPAD, NE, GS, and targeted yield not only increase growth, yield and NUE but also turned out more profitable than a blanket application of N.

REFERENCES

1. Shiferaw B, Smale M, Braun HJ. Crops that feed the world: Past successes and future challenges to the role played by wheat in global food security. *Food Security*. 2013;5:291–317.
2. Toung TP, Bhuiyan SI. Innovations towards improving water-use efficiency in rice. In: Paper Presented at the World Bank's 1994 Water Resource Seminar, VA, USA; 1994.
3. Saharawat YS, Singh B, Malik RK, Ladha JK, Gathala M, Jat ML, Kumar V. Evaluation of alternative tillage and crop establishment methods in a rice-wheat rotation in north-western IGP. *Field Crops Research*. 2010;116:260-267.
4. Das TK, Bhattacharyya R, Sharma AR, Das S, Saad AA, Pathak H. Impacts of conservation agriculture on total soil organic carbon retention potential under an irrigated agro-ecosystem of the western Indo-Gangetic Plains. *European Journal of Agronomy*. 2013;51:34-42.

5. Kassam AH, Stoop W, Uphoff N. Review of SRI modifications in rice crop and water management and research issues for making further improvements in agricultural and water productivity. *Paddy Water Environment*. 2011;9:163-180.
6. Bhattacharyya R, Pandey SC, Bisht JK, Bhatt JC, Gupta HS, Tuti MD, Mahanta D, Mina BL, Singh RD, Chandra S, Srivastva AK, Kundu S. Tillage and irrigation effects on soil aggregation and carbon pools in the Indian sub-Himalayas. *Agronomy Journal*. 2013;105:101-112.
7. Verhulst N, Govaerts B, Verachtert E, Castellanos-Navarrete A, Mezzalama M, Wall P, et al. "Conservation agriculture, improving soil quality for sustainable production systems, in *Advances in Soil Science: Food Security and Soil Quality*. 2010;3:137–208.
8. Lampurlanés J, Cantero-Martínez C. Hydraulic conductivity, residue cover and soil surface roughness under different tillage systems in semiarid conditions. *Soil Tillage Res*. 2006;85:13–26.
9. Page KL, Dang YP, Dalal RC, Reeves S, Thomas G, Wang W. et al. Changes in soil water storage with no-tillage and crop residue retention on a Vertisol: Impact on productivity and profitability over a 50 year period. *Soil Tillage Res*. 2019;194: 104319.
10. Jat HS, Datta A, Choudhary M, Sharma PC, Yadav AK, Choudhary V. et al. Climate Smart Agriculture practices improve soil organic carbon pools, biological properties and crop productivity in cereal-based systems of North-West India. *CATENA*. 2019;181:104059.
11. Somasundaram J, Salikram M, Sinha NK, Mohanty M, Chaudhary RS, Dalal RC et al. Conservation agriculture effects on soil properties and crop productivity in a semiarid region of India. *Soil Res*. 2019;57:187–199.
12. Montgomery DR. Soil erosion and agricultural sustainability. *Proc. Natl. Acad. Sci. U.S.A.* 2007;104:13268–13272.
13. Thierfelder C, Baudron F, Setimela P, Nyagumbo I, Mupangwa W, Mhlanga B et al. Complementary practices supporting conservation agriculture in southern Africa. A review. *Agron. Sustain Dev*. 2018;38:16.
14. Pradhan A, Chan C, Roul PK, Halbrendt J, Sipes B. Potential of conservation agriculture (CA) for climate change adaptation and food security under rainfed uplands of India: a transdisciplinary approach. *Agric. Syst*. 2018;163:27–35.
15. Baveye PC, Rangel D, Jacobson AR, Laba M, Darnault C, Otten W, et al. From dust bowl to dust bowl: soils are still very much a frontier of science. *Soil Sci. Soc. Am. J*. 2011;75:2037–2048.
16. Derpsch R. *Frontiers in conservation tillage and advances in conservation practice*. 10th International Soil conservation Organization: Sustaining the Global Farm, Purdue University. 1999:248-54.
17. Hobbs PR, Sayre K, Gupta R. The role of conservation agriculture in sustainable agriculture. *Biological Sciences*. 2008; 363(1):543–555.
18. Valbuena D, Erenstein O, Tui SH, Abdoulaye T, Claessens L, Duncan AJ, Gérard B, Rufino MC, Teufel N, van Rooyen A, van Wijk MT. Conservation agriculture in mixed crop–livestock systems: scoping crop residue trade-offs in Sub-Saharan Africa and South Asia. *Field Crops Research*. 2012;132:175-84.
19. Hobbs PR, Govaerts B. How conservation agriculture can contribute to buffering climate change. *Climate Change and Crop Production*. 2010;5:177-199.
20. Gebbers R, Adamchuk VI. Precision agriculture and food security. *Science*. 2010;327:828- 831.
21. Robert PC. Precision agriculture: A challenge for crop nutrition management. *Plant and Soil*. 2002;247:143-149
22. Singh, Bijay, Thind HS, Gupta RK. Need-based nitrogen management using the chlorophyll meter and leaf colour chart in rice and wheat in South Asia: A review. *Nutrient Cycling in Agroecosystems*. 2010;88:361–380.
23. Schirrmann M, Domsch H. Sampling procedure simulating on-the-go sensing for soil nutrients. *Journal of Plant Nutrition and Soil Science* 2011;174:333-343.
24. Boggs JL, Tsegaye TD, Coleman TL, Reddy KC, Fahsi A. Relationship between hyperspectral reflectance, soil nitrate-nitrogen, cotton leaf chlorophyll, and cotton yield: A step towards precision agriculture. *Journal of Sustainable Agriculture*. 2003;22:5-16.
25. Furuya S. Growth diagnosis of rice plants by means of leaf color. *Japanese Agricultural Research Quarterly*. 1987;20: 147-153.
26. Gawdiya, Sandeep. Precision nutrient management in conservation agriculture-

- based wheat production, nutrient use efficiency, soil health and profitability in western Uttar Pradesh. M.Sc. Thesis. SVPUAT, Meerut; 2020.
27. Biradar DP, Aladakatti YR, Rao TN, Tiwari KN. Site-Specific Nutrient Management for maximization of crop yields in Northern Karnataka. *Better Crops*. 2006;90(3): 33-35.
 28. Sulochna, Alam, Md, Ali N, Ali, Singh SK. Nitrogen Management by Using Optical Sensor in Wheat in Jharkhand. *Current Journal of Applied Science and Technology*. 2018;31:1-5.
 29. Budhar MN. Leaf Colour Chart based nitrogen management in direct seeded puddled rice (*Oryza sativa* L.). *Fertilizer News*. 2005;50(3):41-44.
 30. Mohanty SK, Singh AK, Jat SL, Parihar CM, Pooniya V, Sharma S, Sandhy CV, Singh B. Precision nitrogen-management practices influences growth and yield of wheat (*Triticum aestivum*) under conservation agriculture. *Indian Journal of Agronomy*. 2015;60(4):617-621.
 31. Mahajan NC. Improving wheat (*Triticum aestivum* L.) and soil productivity through precision nitrogen management practices and efficient planting system. M.Sc. Thesis. SVPUAT, Meerut; 2018.
 32. Khurana HS, Phillips SB, Singh B, Alley MM, Dobermann A, Sidhu AS, Singh Y, Peng S. Agronomic and economic evaluation of site-specific nutrient management for irrigated wheat in northwest India. *Nutrient Cycling in Agroecosystem*. 2008;82:15-31.
 33. Singh RC, Lenka S, Singh CD. Conservation tillage and manure effect on soil aggregation, yield and energy requirement for wheat (*Triticum aestivum* L.) in vertisols. *Indian Journal of Agricultural Sciences*. 2014;84(2): 267–271.
 34. Hussain F, Zia MS, Akhtar ME, Yasin M. Nitrogen management and use efficiency with chlorophyll meter and leaf colour chart. *Pakistan Journal of Soil Science*. 2003;22:1-11.
 35. Kyaw KK. Plot-specific N fertiliser management for improved N-use efficiency in rice-based systems of Bangladesh. In *Ecology and Development Series No. 12* (P.L.G. Vlek, M. Denich, C. Martius and N.V.D. Giesen, Eds), Cuvillier Verlag Göttinge; 2003.
 36. Maiti D, Das DK, Karak T, Banerjee M. Management of nitrogen through the use of leaf color chart (LCC) and soil plant analysis development (SPAD) or chlorophyll meter in rice under irrigated ecosystem. *The Science World Journal*. 2004;4:838–846.
 37. Huan TTN, Tan PS, Hiraoka H. Optimum application rates of nitrogen fertiliser for high- yielding rice cultivars based on growth diagnosis in the Mekong Delta. *Japan International Research Center for Agricultural Sciences*. 2002;26:3-9.
 38. Balasubramanian R. Response of hybrid rice to levels and time of application of nitrogen. *Indian Journal of Agronomy*. 2002;47:203-204.
 39. Huang J, He F, Cui K, Buresh RJ, Xu B, Gong W, Peng S. Determination of optimal nitrogen rate for rice varieties using a chlorophyll meter. *Field Crops Research*. 2008;105:70-80.
 40. Takebe M, Okazaki K, Karasawa T, Watanabe J, Ohshita Y, Tsuji, H. Leaf colour diagnosis and nitrogen management for winter wheat “Kitanokaori” in Hokkaido. *Journal of Soil Science and Plant Nutrition*. 2006;52:577-577.
 41. Uppal RS, Singh J, Kler DS. Effect of NPK levels on growth and yield of durum wheat. *Indian J Environ Ecolplan* 2000;3(5):255-64.
 42. Tayebbeh, Abedi, Abass, Alemzadeh, Abdolreza, Kazemeini Seyed. Effect of organic and inorganic fertilizers on grain yield and protein banding pattern of wheat. *Australian Journal of Crop Science*. 2010;4(6):384-389.
 43. Alam MM, Ladha JK, Foyjunnessa, Rahman Z, Khan SR, Harun-ur-Rashid, Khan AH, Buresh RJ. Nutrient management for increased productivity of rice-wheat cropping system in Bangladesh. *Field Crops Research*. 2006;96:374-386.
 44. Alam MM, Ladha JK, Khan SR, Foyjunnessa, Harun-ur-Rashida, Khana AH, Buresh RJ. Leaf color chart for managing nitrogen fertiliser in lowland rice in Bangladesh. *Agronomy Journal*. 2005;97:949-959.
 45. Ghosh M, Swain DK, Jha MK, Tewari VK. Chlorophyllmeter-based nitrogen management of wheat in Eastern India. *Experimental Agriculture*. 2017;54(3):349-62.

46. Mohanty S, Jat, Shankar, Parihar CM, Singh, Aditya, Sharma, Savita, Saveipune and Kuri, B R. Precision nitrogen management practices in wheat influences nutrient uptake and their use efficiencies and fertility status of soil under conservation agriculture. *Agric. Res.* 2016;37:282-289.
47. Meena, Rekha, Vashisth, Ananta, Das TK, Meena S. Effect of tillage practices on productivity of wheat (*Triticum aestivum* L.). 12th Ann. Agric. Res. 2018;39(1):12-19.
48. Roberts DF, Adamchuk VI, Shanahan JF, Ferguson RB, Schepers JS. Optimization of crop canopy sensor placement for measuring nitrogen status in corn. *Agronomy Journal.* 2009;101:140-149.
49. Verma, Reena, Dhyani, Vipin and Chaturvedi, Sumit. Growth, Yield and Nitrogen Use Efficiency in Wheat as Influenced by Leaf Colour Chart and Chlorophyll Meter Based Nitrogen Management. *Int. J. Curr. Microbiol. App. Sci.* 2017;6(12):1696-1704.
50. Darvishzadeh R, Atzberger C, Skidmore AK. Hyperspectral vegetation indices for estimation of leaf area index. In ISPRS Commission VII Mid-term Symposium. *Remote Sensing.* 2006;3(4):8-11.
51. Hasan MDN, Khaliq QA, Mia MDAB, Bari MDN, Islam MDR. Chlorophyll Meter-Based Dynamic Nitrogen Management in Wheat (*Triticum aestivum* L.) Under Subtropical Environment. *Curr Agri Res.* 2016;4(1).
52. Kong AYY, Fonte SJ, van Kessel C, Six J. Transitioning from standard to minimum tillage: trade-offs between soil organic matter stabilization, nitrous oxide emissions, and N availability in irrigated cropping systems. *Soil and Tillage Research.* 2009;104:256-262.
53. Kumar V, Singh AK, Jat SL, Parihar CM, Pooniya V, Sharma S, Singh B. Influence of site-specific nutrient management on growth and yield of maize (*Zea mays*) under conservation tillage. *Indian Journal of Agronomy.* 2014;59(4): 657-660.

© 2022 Shukla et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/89523>