



A Comparative Study on Growth, Flowering and Corm Production by Growth Regulators & Cut Corms of Gladiolus (Gladiolus grandiflorus L.)

Angel Mariyam Shaji ^{a*}, Samir Ebson Topno ^a and V. M. Prasad ^a

^a *Department of Horticulture, Naini Agriculture Institute, Sam Higginbottom University of Agriculture Technology and Sciences, Prayagraj (U.P.), 211007, India.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2022/v34i2131298

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/89635>

Original Research Article

Received 06 May 2022
Accepted 15 July 2022
Published 19 July 2022

ABSTRACT

The present investigation was carried out during cool season at Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture Technology and Science Prayagraj (U.P.), India. The experiment was laid out in randomized block design with three replications of 9 treatments. Treatments consisted of Full corm, 1/2, 1/3 corms (cut corms) and growth regulators GA3 @150ppm (full corm), 200ppm (1/2 corm, 250ppm (1/3 corm), 300ppm (full corm), 350ppm (1/2, corm), 400ppm (1/3 corm). Among the different treatments, T3 (GA3@ 150ppm full corm) gave the best performance in terms of good growth, flowering and corm production which includes Plant Height, Number of leaves, Days to spike emergence, Days to opening of the first floret, First florets durability (days), Rachis length(cm), Number of florets/ spike, Number of spike/plant, Spike yield/ha, Number of cormlets/planted corm, Corm yield/ha, No.of corms/plant, Weight of Corms/plot.

Keywords: Cut corms; full corms; GA3; Gladiolus; ppm.

1. INTRODUCTION

Gladiolus (Gladiolus spp.) is a significant bulbous crop that is a member of the South African-

originating Iridaceae family. Due of its widespread cultivation among the bulbous ornamentals, it is known as the "Queen of Bulbous Ornaments." This blossom has an its

*Corresponding author: E-mail: angelmariyamshaji123@gmail.com;

beauty and elegance have both financial and aesthetic worth. Excellent are the protracted blossom spikes. when placed in vases as cut flowers for decoration.

Gladiolus is cultivated as a flower bed in gardens and used to create premium bouquets and floral arrangements for interior decorating [1]. On an area of 9.37 thousand acres, it is commercially grown in India's West Bengal, Himachal Pradesh, Sikkim, Karnataka, Uttar Pradesh, Tamil Nadu, Punjab, and Delhi, producing 707 million spikes per year. The crop is currently in high demand on both domestic and foreign markets [2,3]. Any effort to promote cut flower production in the area not only helps consumers and florists obtain fresh and high-quality cut flowers on a regular basis, but it also aids the small and marginal farmers in the area in improving their financial standing [4].

Gladiolus corms are divided into many parts during propagation in order to increase plant material. One of the most cost-effective ways to boost the yield of corms and cormels in this case is to divide the corms [5-7]. According to [8], corm division is mostly determined by the cormlets size and the presence of buds on the corms. Depending on the amount of buds, medium corms are reportedly divided into two or three portions. Each division ought to have a blossom and some root growth. According to [9], the use of various plant growth regulators stimulates early flowering and improves plant development in terms of plant height, flower quantity, and corm yield in gladiolus. In mature propagated plants, where GA3 concentrations are highest, dormancy in both buds and seeds can be overcome. Gibberellic acid breaks dormancy, promotes growth, and postpones senescence. This study was taken to find out the suitable corm size (full and cut) and growth regulators on growth, flowering, spike and corm yield.

2. MATERIALS AND METHODS

The experiment was conducted at Experimental Research Field, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology & Sciences, Prayagraj (UP) during 2021 during rabi season in India.

Some of selected large corms were cut in to sections according to treatment, retaining a bud with each sections, Bavistin was applied to whole

corm to prevent fungus then the segments are treated with GA3 with respect to treatments. The experiment was laid out in randomized block design with three replications consisting of 9 treatments Corms were planted in November at a spacing of 30 cm x 30 cm and a depth of 5–6 cm. Statistical analysis of variance was performed on the data collected throughout the experiment. The observation were recorded for plant height, number of leaves, Days to spike emergence, Days to opening of first floret, Spike length, Rachis length, Number of florets/ spikes, Spike yield, Number of spikes/ha, cormlets yield, Weight of Corm/ plot (g). The significance of the treatments was determined using the 'F' test at a level of significance of 5%.

3. RESULTS AND DISCUSSION

3.1 Growth Parameters

The data on growth parameters in different treatment combinations was recorded (Table 1). Plant height was increased significantly by plant growth regulator application compared to control in the analysis of data. The maximum plant height was recorded in T3 (fullcorm +GA3@150 ppm) (115.16) being significantly superior while minimum was in T8 (1/3 Corm + GA3 @ 400ppm) (91.2). Growth regulators significantly maximum number of leaves in GA3 @150ppm (6.53) was recorded in treatment T3(Full Corm+GA3 @ 150 ppm) which was statistically at par value. Whereas minimum Number of leaves (4.63) was recorded in treatment T8 (1/3 Corm + GA3 @ 400ppm). Maximum Spike emergence (78.46) was found to be in treatment T3 (Full Corm + GA3 @150 ppm) and T0 (Full corm) No chemical (77.96) which was statistically at par. Whereas minimum Spike emergence (61.63) was recorded in treatment T8 (1/3 Corm + GA3 @ 400 ppm). Minimum days to opening the first floret (76.03) was recorded in treatment T8 (1/3 Corm)+GA3 @ 400ppm). Whereas maximum days to opening of the first floret (93.89) was found to be in the treatment T3 with (Full corm+GA3 @150 ppm) which was statistically at par value. Maximum first floret durability (4.00) was found to be in treatment T3(Full Corm+GA3 @ 150 ppm) which was statistically at par value. Whereas minimum first floret durability (3.00) was recorded in treatment T8 (1/3 Corm+GA3 @ 400ppm).

GA3 increased height of the plant over control which may be due to the growth promotion effect of GA3 in stimulating and accelerating cell

division, increasing cell elongation and enlargement or both (Hartmann et al., 1990).

Number of leaves is negatively correlated to peroxidase activity [10] and probably due to these hormone treatments there is suppression of either peroxidase production or inactivation resulting in an increase in number of leaves. Also, GA3 induces active cell division in apical meristem and is also helpful in elongation of individual cells which results in improved vegetative growth.

GA3 induces active cell division in apical meristem and is also helpful in elongation of individual cells which results in spike emergence. Similar results were also obtained by (Barman D et al., 2006).

Floret durability were increases significantly by apply GA3 and this occurs because of GA3 encourage vegetative growth, enhances the photosynthetic and metabolic activities, causing taller plant that provide taller spike and rachis Bhande et al; [11] gladiolus to which spike length and rachis length increased thereby allowing florets to grow larger Neha et al., [12].

3.2 Flowering Parameters

The data on flowering parameters in different treatment combinations was recorded (Table2). Rachis length was increased significantly by plant growth regulator application compared to control in the analysis of data Maximum rachis length (42.83 cm) was found to be in treatment T3 (GA3@ 150ppm+Full Corm) (42.3) which was statistically at par value. Whereas minimum rachis length (40.5) was recorded in treatment T8(1/3 Corm+GA3@ 400ppm). Maximum No. of florets per spike (14.43) was found to be in treatment T3(Full CormGA3+ @ 150 ppm) followed by T0(Fullcorm+No chemical) (13.8) which was statistically at par value. Whereas minimum No. of florets per spike (7.96) was recorded in treatment T8 (1/3 Corm+GA3@ 400ppm). Maximum Spike length (68.97cm) was found to be in treatment T3(Full Corm+GA3 @ 150 ppm) followed by T0(Fullcorm+ No chemical) (67.67) which was statistically at par value. Whereas minimum spike length (61.00) was recorded in treatment T8 (1/3 Corm+GA3 @ 400ppm). Maximum floret diameter (7.96 cm) was found to be in treatment T3 with (Full Corm+GA3 @ 150 ppm) followed by T0(Fullcorm+ No chemical) (7.93) which was statistically at par value. Whereas minimum floret

diameter (7.3) was recorded in treatment T4 (1/3 Corm+GA3 @ 400ppm). Maximum floret length (11.9 cm) was found to be in treatment T3(Full Corm+GA3 @ 150 ppm) followed by T0(Full) No chemical (11.7) which was statistically at par value. Whereas minimum floret length (10.26) was recorded in treatment T8 (1/3 Corm+GA3 @ 400ppm).

Rachis length were increases significantly by apply GA3 and this occurs because of GA3 encourage vegetative growth, enhances the photosynthetic and metabolic activities, causing taller plant that provide taller spike and rachis Bhande et al. [11] gladiolus to which spike length and rachis length increased thereby allowing florets to grow larger Neha et al., [12].

Gibberellic acid promotes the growth of auxiliary buds and their flowering increase in number of florets with the application of GA3 (Panwar et al; Asil et al., [13]).The increased spike length with GA3 might be due to rapid internodal elongation as a result of increased cell division and cell elongation in the intercalary meristem.

Plant growth regulators such as GA3 promotes vegetative growth and increases the photosynthetic and metabolic activities resulting in more transport and utilization of photosynthetic products which might have resulted in increased spike length. Similar results were also reported by Kumar and Singh [14].

Increase in diameter due to low concentrations of GA3 application and decrease due to higher concentration as observed in the present study is in conformity with the findings of Bhattacharjee (1984) who observed that concentrations of GA3 as it increased from 10 to 1000 ppm decreased the flower diameter correspondingly. Increase in flower diameter by GA3 application was also reported by Prakash and Jha (1998).

3.3 Yield Parameters

The data on yield parameters in different treatment combinations was recorded (Table 2). No of spike/plant was increased significantly by plant growth regulator application compared to control in the analysis of data. Maximum No. of spike per plant (2.3) was found to be in treatment T3(Full Corm+GA3 @ 150 ppm) followed by T0 (Fullcorm+ No chemical) (2.03) which was statistically at par value. Whereas minimum No. of spike per plant (1.26) was recorded in treatment T8(1/3 Corm+GA3 @ 400ppm).

Maximum spike yield (207000) was found to be in treatment T3 (Full Corm+GA3 @ 150 ppm). Whereas minimum spike yield (113400) was recorded in treatment T8 (1/3 Corm+GA3 @ 400ppm). Maximum No. of cormlets /planted corm (3.17) was found to be in treatment T3(Fullcorm+GA3@150 ppm) followed by T0(Fullcorm+No chemical) (3.06) which was statistically at par value. Whereas minimum No. of cormlets /planted corm. (1.16) was recorded in treatment T8(1/3 Corm+GA3 @ 400ppm). Maximum Corm yield/ha (1199.35) was found to be in treatment T3 (Full Corm+GA3 @ 150 ppm). Whereas minimum Corm yield/ha (450) was recorded in treatment T8 (1/3 Corm+GA3@ 400ppm). Maximum No. of Corms/plant (2.17) was found to be in treatment T3 with (Full Corm) GA3 @ 250 ppm. Whereas minimum No. of Corms/plant. (1.57) was recorded in treatment T8 (1/3 Corm+ GA3 @ 400ppm). Maximum Weight of Corm / plot (g) (36.00) was found to be in treatment T3 with (Full Corm) GA3 @ 250 ppm. Whereas minimum Weight of Corm/ plot (g). (27.83) was recorded in treatment T8 (1/3 Corm+GA3 @ 400ppm).

Growth regulators like GA3, which promotes the crop growth by enhanced cell division and enlargement, leading to proper shoot and root growth that ultimately favours of maximum number of spikes and increased spikes yield. Results are in consonance with the findings of (Pathak, G. 2007) and [15].

GA3 causes more splitting and cell division than increasing the size of corms [16] in gladiolus. Khan et al. [17] found that higher concentrations of GA3 enhanced multiple shooting and accelerated corm production in gladiolus. The result is in conformity with the work of Raju [18] in lilies and Rajaram et al. [19] in gladiolus.

GA3 promoted the sink activity of developing corms and cormels at the expense of flower spike, this might be the reason for increase in number of corms and cormels and quality flower spikes. Similar results were also observed by Tawar et al. [20] in gladiolus.

Table 1. Effect of cut corms and different growth regulators on growth parameters of gladiolus

Treatment	Plant height(cm) 90 DAS	No. of leaves per plant 90 DAS	Days to spike emergence	Days to opening of first Floret	First floret Durability
T0 Full corm (no chemical)	111	6.53	68.36	90.80	3.67
T1 ½ corm (no chemical)	104.03	6.33	61.63	82.37	3.20
T2 1/3 corm (no chemical)	100.66	5.8	72	79.83	3.17
T3 GA3 @ 150 ppm (full corm)	115.16	6.03	68.33	93.89	4.00
T4 GA3 @ 200 ppm (1/2corm)	109.16	5.5	74.4	86.20	3.23
T5 GA3@ 250 ppm (1/3corm)	98.16	6.33	65.06	77.03	3.13
T6 GA3 @ 300 ppm (full corm)	107.83	5.43	77.96	82.50	3.10
T7 GA3 @ 350 ppm (1/2corm)	93.83	5.23	75.2	76.70	3.07
T8 GA3 @ 400 ppm (1/3 corm)	91.2	6.63	78.46	76.03	3.00
F- test	S	S	S	S	S
S.Em.(±)	0.63	0.11	1.03	0.52	0.07
C.D. at 5 %	1.85	0.33	3.02	1.54	0.20
C.V	1.41	4.37	3.34	1.47	4.87

Table 2. Effect of cut corms and different growth regulators on flowering and yield parameters of gladiolus

Treatment	Flowering parameters						Yield parameters				
	Rachis length (cm)	No. of florets/spike	Spike length (cm)	Floret Diameter (cm)	Floret Length (cm)	No. of spike/ha (lakh)	No. of Cormlet/plant	No. of corms/plant	Weight of Corms/plot (g)	Corm Yield/ha (kg)	
T0	42.6	13.8	67.67	7.93	11.7	182700	11.83	2.07	35.53	790	
T1	41.93	9.7	64.50	7.63	10.96	155700	10.50	1.83	31.57	700	
T2	41.9	9.43	64.17	7.56	10.8	146700	9.83	1.73	29.60	690	
T3	42.83	14.43	68.97	7.96	11.9	207000	14.00	2.17	36.00	1200	
T4	42.3	13.53	66.90	7.8	11.4	171000	11.50	1.97	33.87	900	
T5	41.83	9.33	62.33	7.5	10.66	144000	8.50	1.67	29.00	600	
T6	41.96	11.46	66.30	7.66	11.16	140400	11.10	1.90	33.20	800	
T7	41.16	8.8	62.00	7.33	10.4	122400	7.83	1.60	28.00	590	
T8	40.5	7.96	61.00	7.3	10.26	113400	7.50	1.57	27.83	450	
F- test	S	S	S	S	S	S	S	S	S	S	
S.Em. (±)	0.25	0.22	0.49	0.03	0.08	1.82700	0.29	0.06	0.42	0.22	
C.D. at 5 %	0.75	0.67	1.46	0.11	0.23	1.55700	0.84	0.18	1.23	0.65	
C.V	1.41	4.84	1.77	1.13	1.68	1.46700	6.40	7.93	3.07	2.52	

Under peer review

4. CONCLUSION

From the present investigation, it is concluded that, treatment T3(GA3@150ppm+full corm) gave the best performance in terms of growth, flowering and corm production.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Lepcha B, Nautiyal MC, Rao VK. Variability studies in gladiolus under mid hill conditions of uttarakhand. *Journal of Ornamental Horticulture*. 2007;10(3):169-172.
- Mckay E, Byth DE, Tommerup J. The effect of corms size and division of the cormlets in gladiolus. *Australian Journal of Experimental Agriculture*. 1981;21(110):343-348.
- Candyman SA, Langthasa DN, Hazarika B, Gautam P, Goswami RK. Influence of GA₃ and BA on morphological, phenological and yield attributes in gladiolus cv. Red IOSR *Journal of Agriculture and Veterinary Science*. 2015;8(6):37-42.
- Naresh S, Dorajee RAVD, Vijaya B, V, Uma KK, Paratpara RM. Evaluation of gladiolus hybrids under coastal Andhra Pradesh conditions. *Plant Archives*. 2015;(1):451-454.
- Tameshwar Prasad VM, Narayanan Lal. Effect of plant growth regulators on gladiolus cv. jester. *Trends in Biosciences*. 2017;10(20):3982- 3985. ISSN 0974-8431.
- Rashid MHA. Influence of size and plant growth regulators on corm and cormel production of gladiolus (*Gladiolus grandiflorus* L.). *Progressive Agriculture*. 2018;29(2).
- Reshma VS, Panchbhai DM, Kumar P, Adarsh MN. Effect of GA3 spray on gladiolus (*Gladiolus* spp.) Varieties under Dry Conditions of Vidharba Region, *International Journal. Pure App. Biosci*. 2017;5(3):123-129.
- Gromov AN. Propagation of Gladiolus corms and cormels. *The world of the gladiolus*. NAGC, USA. 1972;98-102.
- Singh A, K, Kumar R, Sisodia A. Effect of GA3 on growth and flowering attributes of gladiolus cultivars. *Annals of Agriculture Research New Series*. 2013;34:315-19.
- Galston AW, Davies PJ. Hormonal regulation in higher plants. *Science*. 1969; 163:1288–97.
- Bhande SS, Ambade RB, Shinde DV, Ambade SB, Patil SA, Naushad M, Mane RS, Alothman ZA, Lee SH, Han SH. Improved photoelectrochemical cell performance of tin oxide with functionalized multiwalled carbon nanotubes–cadmium selenide sensitizer. *ACS Applied Materials & Interfaces*. 2015; 7(45):25094-104.
- Neha Chopde, Ashwini Patil, Bhand MH. Growth, yield and quality of gladiolus as influenced by growth regulators and methods of application. *Plant Archives*. 2015;15(2):691-694.
- Asil MH, Roen Z, Abbasi J. Response of tuberose (*Polian tuberosa* L.) to gibberellic acid and benzyl adenine. *Horticulture, Env and Biotech*. 2011;52:46-51.
- Kumar MD, Singh OP. Virtual water in global food and water policy making: is there a need for rethinking?. *Water resources management*. 2005;19(6):759-89.
- Singh BN, Connolly SJ, Crijns HJ, Roy D, Kowey PR, Capucci A, Radzik D, Aliot EM, Hohnloser SH. Dronedaron for maintenance of sinus rhythm in atrial fibrillation or flutter. *New England Journal of Medicine*. 2007;357(10):987-99.
- Baskaran R, Cullen R, Colombo S. Estimating values of environmental impacts of dairy farming in New Zealand. *New Zealand Journal of Agricultural Research*. 2009;52(4):377-89.
- Khan FN, Rahman MM, Hossian MM. Effect of benzyl adenine and gibberellic acid on dormancy breaking, growth and yield of gladiolus corms over different storage periods. *Journal of Ornamental Horticulture*. 2012;3:59-71.
- Raju TS. Terminal sugars of Fc glycans influence antibody effector functions of IgGs. *Current opinion in immunology*. 2008;20(4):471-8.
- Rajaram PK. Humanitarianism and Representations of the Refugee. *Journal of refugee studies*. 2002;15(3):247-64.

20. Tawar U, Bansal S, Shrimal S, Singh M, Tandon V. Nuclear condensation and free radical scavenging: a dual mechanism of bisbenzimidazoles to modulate radiation damage to DNA. *Molecular and cellular biochemistry*. 2007;305(1):221-33.

© 2022 Shaji et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/89635>