



Evaluation of Fungicides and Bio-Agents against *Ascochyta* Blight (*Ascochyta rabiei*) of Chickpea under Dryland Conditions of Kashmir, India

Sabiya Bashir^{a*}, Mohammad Najeeb Mughal^b, Shabeena Majeed^a, Zahida Rashid^a, Sabeena Naseer^a, Zahoor A. Dar^a, S. A. Hakeem^a, Faisal Rasool^a, Shaheena Nagoo^c, Sabiha Ashraf^c and Shahida Iqbal^a

^a Dryland Agriculture Research Station, Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir, Shalimar, Srinagar-190025, India.

^b Division of Plant Pathology, Faculty of Horticulture, SKUAST-Kashmir, India.

^c College of Temperate Sericulture, SKUAST-Kashmir, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2022/v34i2131567

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/91883>

Original Research Article

Received 20 May 2022
Accepted 26 July 2022
Published 29 July 2022

ABSTRACT

Five non-systemic fungicides: chlorothalonil, carbendazim (12%) + mancozeb (63%), captan 50WP, dodine 65WP and copper oxychloride 50WP and four systemic fungicides viz., difenoconazole 25EC, carbendazim 50WP, hexaconazole 5EC and myclobutanil 10WP were evaluated against *Ascochyta* blight of chickpea caused by *Ascochyta rabiei*. The *in-vitro* evaluation of non-systemic fungicides through poisoned food technique at five different concentrations: 50, 100, 250, 500 and 1000 $\mu\text{g ml}^{-1}$ fungicides showed that carbendazim (12%) + mancozeb (63%) proved most effective and resulted in highest mycelial growth inhibition (82.27%) of the pathogen followed by dodine with mycelial growth inhibition of 67.99%. The least efficacious fungicide was copper oxychloride with only 29.81% mycelial growth inhibition. Among the systemic fungicides evaluated at concentrations of 25, 50, 100, 200 and 500 $\mu\text{g ml}^{-1}$, carbendazim proved to be the most effective and caused highest mycelial growth inhibition of 90.23% followed by difenoconazole (71.24%). Myclobutanil 10WP was least efficacious among the systemic fungicides and resulted in only 52.76% of mycelial growth inhibition. Among the three methods used for evaluation of efficacy of fungicides and bio-

*Corresponding author: E-mail: drsabiyaashir@gmail.com;

agents, revealed that the seed treatment with carbendazim (12%) + mancozeb (63%) proved most effective with lowest disease incidence (15.0%) and disease intensity (4.02%). Among the biological control agents used, seeds treated with *Bacillus subtilis* proved the most effective which resulted in disease incidence and intensity of 32.25 and 20.04%, respectively. Combination treatment comprising of seed treatment with *Bacillus subtilis* at concentration of 10^9 spores ml^{-1} and foliar spray with mancozeb 63% WP+ carbendazim 12%WP at the concentration of 2.5% was most efficacious and resulted in lowest disease incidence (7.80%) and disease intensity of (4.82%) while seed treatment with *Bacillus subtilis* at the concentration 10^9 spores ml^{-1} alone was least efficacious and resulted in highest disease incidence of 53.80% with disease intensity of 37.25%. It was, however superior than control where disease incidence and intensity was 80.00% and 51.34 %, respectively.

Keywords: *Ascochyta*; *bio-agents*; *chickpea*; *disease incidence*; *fungicides*; *seed treatment*.

1. INTRODUCTION

Chickpea (*Cicer arietinum* L.) is an important pulse crop belongs to family *Fabaceae* grown world-wide and is ranked third after beans and dry peas [1]. India is the largest producer, consumer and importer of the chickpea in world, contributing 71 % in area and 71.95% in production [2]. Chickpea contributes nearly 48 to 52% of total pulse production in India [3]. The total area under its cultivation in India is about 105.61 lakh hectares with annual production of 112.29 lakh tonnes and an average productivity of 1063 kg ha^{-1} [4]. The land placed under production of chickpea in Jammu and Kashmir during the year 2012-13 was 2000 hectare with average production of 1000 tonnes ha^{-1} and yield of 542 kg ha^{-1} [5].

Chickpea crop is affected by several diseases, pests and poor management practices resulting in heavy yield losses. Amongst them, diseases play important role in reducing yield potential of chickpea [6]. About 50 diseases have been reported to affect chickpea with huge economic importance. These include wilt [*Fusarium oxysporum* f. sp. *ciceri* (Padwick) Synd & Hans.], black root rot [*Fusarium solani*(Mart.)Sacc.], wet root rot (*Rhizoctonia solani* Kuhn.), dry root rot [*Rhizactonia bataticola* (Taub.) Butler], Aschochyta blight [*Aschochyta rabiei* (Pass.) Lab.], Pythium root and seed rot (*Pythium ultimum* Trow.) and collar rot (*Sclerotium rolfsii* Sacc.). Ascochyta blight (*Ascochyta rabiei*) is one of the most destructive diseases of chickpea and is wide spread in its distribution [7]. Lesions can occur on leaves, leaflets, stems, pods and seeds. Symptoms begin to appear after 4-6 days after the spores infect the plant. Conidia are exuded from fruiting body called pycnidia in a sticky spore mass and are spread to the nearby plants during rain-splashes [8]. The pathogen

survives on diseased crop debris and on diseased seeds in the form of pycnidia containing conidia. Primary infection takes place through conidia and causes necrotic spots which develop into conidia at the centre. These conidia work as inoculum for secondary infection during rains [9]. Ascochyta blight is mainly controlled with fungicides and bio-agents [10].

Fungicides with novel chemistry are being introduced and evaluated before their application can be recommended to farmers. Therefore, *in-vitro* and *in-vivo* evaluation of these fungicides is of paramount importance. There is constant need to compare and evaluate new fungicides along with some compatible bio-agents for controlling fungal diseases such as ascochyta blight. Use of fungicides create environmental pollution, but bio-agents are environment friendly. Hence, the present study was carried out to evaluate non-systemic and systemic fungicides and some bio-control agents against the ascochyta blight of chickpea.

2. MATERIALS AND METHODS

Five non-systemic fungicides: chlorothalonil, carbendazim (12%)+mancozeb (63%), captan 50WP, dodine 65WP and copper oxychloride 50WP and four systemic fungicides: difenoconazole 25EC, carbendazim 50WP, hexaconazole 5EC and mycobutanil 10WP were evaluated against aschochyta blight of chickpea using poisoned food technique [11] where potato dextrose agar was medium. The non-systemic fungicides were evaluated at different concentrations of 50, 100, 250, 500 and 1000 $\mu\text{g ml}^{-1}$ and systemic fungicides at 25, 50, 100, 200 and 500 $\mu\text{g ml}^{-1}$. Different concentrations of fungicides were prepared by adding appropriate quantity of fungicides to sterilized molten potato dextrose agar (PDA) medium in conical flasks.

On sterilized petri plates, 25 ml of PDA medium was aseptically poured. After cooling, 5mm diameter mycelial disc of *Ascochyta* blight pathogen was aseptically placed in the centre of each petri plate. Petri plates containing only PDA media amended with equal amount of sterilized water and inoculated with mycelial disc of the pathogen (5mm diameter) served as check. Each treatment was replicated four times and incubated at 28±2°C for next 4 days. The comparative efficacy of each fungicide was calculated as per cent inhibition of mycelial growth of the test pathogen in each treatment as compared to check by the formula described by Vincent [12]:

$$\text{Per cent mycelial growth inhibition} = \frac{C - T}{C} \times 100$$

Where

C = Radial mycelial growth (mm) in check

T = Radial mycelial growth in the treatment (mm)

The three methods: seed treatment, soil treatment and foliar application were used for evaluation of efficacy of fungicides and bio-agents against *Ascochyta* blight of chickpea in three different plots. Field trials were conducted during Rabi season of 2018-19 at Dryland Agriculture Station (DARS), SKUAST-Kashmir, Rangreth. Field was equally divided into beds with plot size of 2m×1m, with each plot having 50 stands of chickpea plants. Three replications for each treatment were maintained in completely randomized blocks designs. In first method (seed treatment), the chickpea seeds of susceptible variety (Local channa) were sown after treatment with fungicides and bio-agents. Similarly in the second method (soil treatment), soil was treated with fungicides and bio-agents at recommended concentrations and the seeds of susceptible variety was sown. The third method involving sowing, susceptible chickpea seeds and the crop was sprayed with fungicides and bio-agents after 15 days of seed emergence. Disease incidence and intensity for each treatment were recorded after 30 days after seed emergence.

The bio-control agent and the fungicide which proved most effective in *in-vitro* were further evaluated under field conditions for their efficacy in controlling the disease. Further integrated management trials were conducted during Rabi season 2019-20 at Dryland Agriculture Station, SKUAST-K Rangreth. Field was equally divided into beds with plot size of 2m×1m with each plot having 50 stands of chickpea plants. Three replications for each treatment were maintained in completely randomized blocks designs. The

chickpea seeds of susceptible variety (Local channa) were sown after being dressed with different combination of bio-agent and fungicides as seed treatment. The foliar sprays were done 15 days after seed emergence. The plots where only water was sprayed served as check. Observations on disease incidence and intensity were recorded 30 days after seed emergence.

3. RESULTS AND DISCUSSION

In *in-vitro* evaluation of non-systemic and curative fungicides revealed that carbendazim (12%) + mancozeb (63%) proved most effective in causing the highest mycelial growth inhibition (82.27%) of the pathogen followed by dodine with mycelial growth inhibition of 67.99%. The least efficacious fungicide among tested fungicide was copper oxychloride with 29.81% mycelial growth inhibition (Table 1), while among the systemic fungicides (Table 2), carbendazim proved the most effective in causing the mycelial growth inhibition of 90.23% followed by difenoconazole with 71.24% mycelial growth inhibition. The least effective fungicide was myclobutanil with 52.76% mycelia growth inhibition (Table 2). Further, the different methods of application of fungicides, seed treatment was found effective in controlling the disease followed by foliar spray. The data revealed that the seed treatment with carbendazim (12%) + mancozeb (63%) proved most effective with lowest disease incidence (15.0%) and disease intensity (4.02%) followed by carbendazim with 16.50% and 4.82% disease incidence and disease intensity. The least efficacious fungicide was copper oxychloride with 60.25% and 40.32% disease incidence and intensity which is still superior over check with disease incidence and intensity of 83.33% and 71.11%, respectively. The foliar spray also showed better results than soil application method, where disease incidence and disease intensity was 20.25% and 6.24% when sprayed with carbendazim (12%) + mancozeb (63%) followed by carbendazim with 25.00% and 7.15% disease incidence and intensity. Among the biological control agents used, seeds treated with *Bacillus subtilis* proved most effective and resulted in disease incidence and intensity of 32.25% and 20.04%, respectively, followed by *Pseudomonas fluorescens* with 42.00% and 25.34% disease incidence and intensity. Similarly, foliar spray with *Bacillus subtilis* also showed promising results as disease incidence and intensity was recorded 39.50% and 26.15%, respectively (Table 3).

Table 1. *In-vitro* evaluation of non-systemic fungicides against *Ascochyta rabiei* causing blight of chickpea during Rabi 2018-19

| S. No. | Treatment | % inhibition of radial mycelial growth at concentration ($\mu\text{g ml}^{-1}$) | | | | | Mean |
|--------|-----------------------------------|---|---------------|---------------|---------------|----------------|-------|
| | | 50 | 100 | 250 | 500 | 1000 | |
| 1. | Chlorothalonil | 44.03 (41.03) | 53.26 (46.87) | 53.26 (46.87) | 64.16 (53.23) | 80.20 (63.62) | 58.98 |
| 2 | Carbendazim (12%) +mancozeb (63%) | 61.66 (51.75) | 75.33 (60.17) | 81.43 (64.48) | 92.96 (74.67) | 100.00 (90.00) | 82.27 |
| 3 | Captan 50 WP | 35.76 (36.76) | 43.13 (41.05) | 43.16 (41.06) | 51.83 (46.04) | 60.43 (51.02) | 46.86 |
| 4 | Dodine 65 WP | 44.40 (41.77) | 70.06 (56.84) | 71.86 (57.98) | 72.20 (58.20) | 81.43 (64.48) | 67.99 |
| 5 | Copper oxychloride 50WP | 25.90 (30.59) | 28.33 (32.11) | 27.06 (31.26) | 29.56 (32.93) | 38.20 (38.17) | 29.81 |
| | Mean | 42.35 | 54.02 | 55.35 | 62.14 | 72.05 | |

CD ($p=0.05$); Non-systemic fungicide = (2.1); Concentration = (1.9); Fungicide \times Concentration = (4.70)*Figures within parentheses are arc sign transformed value

Table 2. *In-vitro* evaluation of systemic fungicides against *Ascochyta rabiei* causing blight of chickpea during Rabi 2018-19

| S. No. | Treatment | % inhibition of radial mycelial growth at concentration ($\mu\text{g ml}^{-1}$) | | | | | Mean |
|--------|-----------------------|---|---------------|---------------|----------------|----------------|-------|
| | | 25 | 50 | 100 | 200 | 500 | |
| 1. | Difenoconazole 25 EC, | 53.80 (41.17) | 61.50 (51.64) | 66.60 (54.69) | 74.30 (59.53) | 100.00 (90.00) | 71.24 |
| 2 | Carbendazim 50 WP | 82.00 (64.89) | 84.56 (66.86) | 84.60 (66.89) | 100.00 (90.00) | 100.00 (90.00) | 90.23 |
| 3 | Hexaconazole 5EC | 48.66 (44.23) | 56.33 (48.63) | 64.10 (53.19) | 66.63 (54.71) | 66.70 (54.74) | 64.56 |
| 4 | Mycobutanil 10 WP | 43.53 (41.28) | 51.23 (45.70) | 51.20 (45.68) | 56.33 (48.63) | 61.50 (51.64) | 52.76 |
| | Mean | 56.99 | 63.40 | 66.62 | 74.31 | 82.05 | |

CD ($p=0.05$); Systemic Fungicide = (1.27); Concentration = (1.25); Fungicide \times Concentration = (2.33); *Figures within parentheses are arc sign transformed value

Table 3. Evaluation of different fungicides and bio-control agents against *Ascochyta* blight of chickpea through seed treatment, soil treatment and foliar application during Rabi 2019-20

| S. No. | Treatment | Con. (%) | Seed treatment | | Soil treatment | | Foliar spray | |
|--------|----------------------------------|----------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|
| | | | Disease incidence (%) | Disease intensity (%) | Disease incidence (%) | Disease intensity (%) | Disease incidence (%) | Disease intensity (%) |
| 1. | Chlorothalonil | 0.30 | 35.50(20.79) | 10.75(19.13) | 49.25(44.57) | 17.37(24.63) | 43.25(41.12) | 13.24(21.33) |
| 2 | Carbendazim (12%)+mancozeb (63%) | 0.25 | 15.00(22.78) | 4.02(11.56) | 21.75(27.79) | 7.80(16.21) | 20.25(26.74) | 6.24(14.46) |
| 3. | Carbendazim 50 WP | 0.05 | 16.50(23.96) | 4.82(12.68) | 28.75(32.42) | 9.23(17.68) | 25.00(30.00) | 7.15(15.50) |
| 4. | Difenoconazole 25EC | 0.05 | 23.25(28.82) | 8.52(16.97) | 36.50(37.16) | 15.54(23.21) | 27.25(46.26) | 10.14 (18.56) |
| 5. | Hexaconazole 5 EC | 0.03 | 37.25(37.61) | 12.15(12.15) | 51.50(45.85) | 18.75(25.65) | 45.50(42.41) | 15.11(22.87) |

| S. No. | Treatment | Con. (%) | Seed treatment | | Soil treatment | | Foliar spray | |
|--------|--------------------------------|--------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|
| | | | Disease incidence (%) | Disease intensity (%) | Disease incidence (%) | Disease intensity (%) | Disease incidence (%) | Disease intensity (%) |
| 6. | Captan 50 WP | 0.30 | 44.00(41.55) | 27.88(31.87) | 60.25(50.91) | 42.27(40.55) | 48.25(43.99) | 30.12(33.28) |
| 7. | Dodine 65 WP | 0.06 | 34.50(35.97) | 10.25(18.67) | 47.32(25.60) | 16.09(23.64) | 41.25(39.96) | 12.76(20.92) |
| 8. | Myclobutanil 10 WP | 0.10 | 38.25(22.48) | 13.42(21.48) | 52.00(46.14) | 19.13(25.93) | 45.75(42.56) | 16.32(23.82) |
| 9. | Copper oxychloride | 0.30 | 60.25(50.91) | 40.32(39.41) | 65.50(54.02) | 51.34(45.76) | 61.15(51.44) | 45.14(42.21) |
| 10. | <i>Trichoderma viridi</i> | 0.2 | 44.25(41.69) | 31.13(33.91) | 52.15(46.23) | 37.11(37.53) | 47.25(43.42) | 33.42(35.31) |
| 11. | <i>Bacillus subtilis</i> | 0.2 | 32.25(34.60) | 20.04(26.59) | 35.75(36.72) | 28.02(31.96) | 39.50(38.93) | 26.15(30.75) |
| 12. | <i>Pseudomonas fluorescens</i> | 0.2 | 42.00(40.39) | 25.34(34.42) | 50.75(45.42) | 35.23(36.40) | 45.15(42.21) | 25.75(30.49) |
| 13. | Check (Untreated) | 83.33(65.90) | | 71.11(57.48) | | | | |

Table 4. *In-vivo* field evaluation of the bio-agent and fungicide against the Ascochyta blight of Chickpea during 2020-21

| S. No | Treatment | Mean disease Incidence (%) | Mean Disease Intensity (%) |
|-------|--|----------------------------|----------------------------|
| T1 | Seed treatment <i>Bacillus subtilis</i> @10 ⁹ spores ml ⁻¹ | 53.80(41.17) | 37.25(37.61) |
| T2 | Seed treatment with mancozeb 63% WP+ carbendazim12%WP @ 2g/kg of seed | 36.50(37.16) | 20.25(26.74) |
| T3 | Seed treatment with carbendazim 50WP 1g/kg of seed | 44.25(41.69) | 34.50(35.97) |
| T4 | Seed treatment with <i>Bacillus subtilis</i> @10 ⁹ spores per ml and foliar spray with same @10 ⁹ | 21.75(27.79) | 16.50(23.96) |
| T5 | Seed treatment with <i>Bacillus subtilis</i> @10 ⁹ spores ml ⁻¹ and foliar spray with mancozeb 63%WP + carbendazim12%WP @ 2.5% | 7.80(16.21) | 4.82(11.56) |
| T6 | Seed treatment with <i>Bacillus subtilis</i> @10 ⁹ spores ml ⁻¹ foliar spray with carbendazim 50WP @ 2.5% | 17.37(24.63) | 8.25(16.97) |
| T7 | Foliar with <i>Bacillus subtilis</i> @10 ⁹ spores ml ⁻¹ | 51.50(45.85) | 31.13(33.91) |
| T8 | Foliar spray with mancozeb 63% WP+ carbendazim12%WP @ 2.5% | 20.25(26.74) | 13.42(21.48) |
| T9 | Foliar spray with carbendazim @ 0.5% | 18.75(25.65) | 9.23(17.68) |
| T10 | Control | 80.00(64.89) | 51.34(45.76) |

CD ($p \leq 0.05$) 0.34; Drenching was done after 15 days of emergence

Integrated disease management experiment conducted during Rabi 2020-21 to study the effect of some promising bio-agents and fungicides under field conditions as seed treatment and foliar spray against the ascochyta blight of chickpea at DARS, Rangreth revealed that carbendazim (12%) + mancozeb (63%) was most effective systemic fungicide (carbendazim 50 WP) and *Bacillus subtilis* was the most effective bio-agent for management of the disease. The results on disease incidence and intensity (Table 4) revealed that all the seed treatments and foliar spray were superior over the check in controlling the disease incidence and intensity. In the experiment the mean disease incidence ranged from 7.80% to 80.00% and the mean disease intensity ranged from 4.82% to 51.34%. Lowest disease incidence of 7.80% was recorded in treatment combination of seed treatment with *Bacillus subtilis* applied at concentration of 10^9 spores ml^{-1} and foliar spray with carbendazim (12%) + mancozeb (63%) applied at concentration of 2.5%. Highest disease incidence of 80.00% was observed in check plots. Seed treatment with *Bacillus subtilis* applied at 10^9 spores ml^{-1} and foliar spray with carbendazim 50WP applied at concentration of 2.5% resulted in the disease incidence and disease intensity of 17.37% and 8.25% followed by foliar spray with carbendazim applied at concentration of 0.5% with disease incidence and disease intensity of 18.75% and 9.23%, respectively. Among the different seed treatments, least disease incidence and intensity of 36.50% and 20.25% was observed when the seeds were treated with mancozeb 63% WP+ carbendazim 12%WP applied at the concentration of 2g/kg of seed.

Under field conditions, during Rabi 2020-2021 integrated disease management option that involved seed dressing with *Bacillus subtilis* applied at the concentration of 10^9 spores ml^{-1} and foliar spray with mancozeb (63%WP) + carbendazim (12%WP) applied at the concentration of 2.5% at 15 days after seed emergence proved effective in managing this disease. Our results are in conformity with those reported by Abdel Kader et al. [13]; Ahmad et al. [14] Chongo et al. [15], Gurdip and Maninder [16]. The fungicides and bio-agents which can effectively inhibit the test fungus in laboratory conditions are supposed to be effective against the same fungal pathogen under field conditions. Generally, all the treatments and combinations check the activities of the inoculated fungus, *Ascochyta rabiei* and hence promote the healthy

conditions of chickpea crop. Although with the increase in concentrations, few fungicides completely inhibited the growth of *Ascochyta rabiei* [17]. In the present investigation, *Bacillus subtilis* proved most effective in laboratory conditions and these findings were completely in agreement with many workers who found many isolates/strains of *Bacillus* isolated from the rhizosphere regions of host crop plants effective in managing the disease [18-22].

In this approach of integrating fungicides with bio-agents eliminates the effect of *Ascochyta rabiei* through double action. The fungicides eliminates the seed and soil borne inoculums, while the bio-agents takes care of soil-borne inoculums and increases crop productivity by improving nutrient status and soil health [23-25].

4. CONCLUSIONS AND RECOMMENDATIONS

It is important to integrate all available methods into a management package for *Ascochyta rabiei* under chickpea production. The use of seed free of contaminating pathogens will reduce the probability of transmitting seed-borne disease to the seedlings. The use of fungicide and biological agents could be two key measures for a rational integrated management of ascochyta blight of chickpea in sustainable cropping systems in Kashmir. Inclusion of fungicides and bio-agents in the integrated management of *Ascochyta rabiei* under chickpea production in Kashmir.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Shifa S, Sabiya B, Najeeb M, Reyaz RM, Bhat FA, Shah TA. Identification of Sources of Resistance in Chickpea (*Cicer arietinum*) against Wilt (*Fusarium oxysporum* f. sp. ciceri) under Temperate Agro-Climatic Conditions of Kashmir. International J Current Microbio and App Sci. 2018;7(9):190-194.
2. Samriti, Subhash S, Ravinder S, Ankit Pathania .Trends in Area, Production, Productivity and Trade of Chick Pea in India .Eco Affairs.2020;65(2):261-265.

3. Anonymous. Pulses Revolution- From Food to Nutritional Security. Ministry of Agriculture and Farmers Welfare (DAC & FW), Govt. of India. 2018;3-4.
4. FAOSTAT. Food and Agriculture Organization. Statistical Database of the United Nation Food and Agriculture Organization Statistical Division .2019; Rome.
5. Anonymous. E-pulses Data book(Station wise), Directorate of Economics and statistics, Krishi Bhawan, New Delhi, powered by Indian Institute of Pulse Research Institute, Kanpur; 2016. Available: <http://www.indiaagristat.com>
6. Nene YL. A review of *Ascochyta* blight of chickpea. *Int J Pest Manage.* 1982;28:61–70.
7. Ahmad S, Khan MA, Sahi ST, Ahmad R. Evaluation of chickpea germplasm against *Ascochyta rabiei* (Pass). *J Animal & Plant Sci.* 2013;23(2):440-443.
8. Reddy MV, Singh KB. Relationship between *Ascochyta* blight severity and yield loss in chickpea and identification of resistant lines. *Phytopathol. Mediterr.* 1990;29:32–38.
9. Shtienberg D. Applications of epidemiology in the management of *ascochyta* blight in chickpea and lentil. In: *Compendium of Chickpea and Lentil Diseases and Pests.* 2010; (Chew, W., Muehlbauer, F.J., and Sharma, H. C. eds.). APSPress, St Paul, Minnesota, USA.
10. Salman A, Muhammad AK, Irfan A, Zafar I, Ejaz A, Muhammad AYA, Saima N. Efficacy of fungicides, plant extracts and bio-control agents against *Ascochyta* blight (*Ascochyta rabiei*) of chickpea (*Cicer arietinum*) under field conditions. *Plant. Sci Today.*2012;8(2):255-262.
11. Carpenter JB. A toximetric study of some eradicant fungicides. *Phytopathol.* 1942;32 :845.
12. Vincent JH. Distortion of fungal hyphae in presence of certain inhibitors. *Nature.* 1947;15:850.
13. Abdel Kader DA, EW Kil A, Tohami MR, Ghoniem MI. Effect of some agricultural practices and chemical control on the incidence of *ascochyta* blight of chickpea plants. *Egyptian J Phytopathol.*1989;21 (1):31-44.
14. Ahmad HU, Hwang SF, Gossen BD, Chang KF, Howard RJ, Burke D. Chemical control of *Ascochyta* blight (*Ascochyta rabiei*) of Chickpea. *Canadian J Plant Patho.* 2008;30(2):367-367.
15. Chongo G, Buchwaldt L, Gossen BD, Lafond GP, May WE, Johnson EN, Hogg T. Foliar fungicides to manage *ascochyta* blight (*Ascochyta rabiei*) of chickpea in Canada. *Canadian J Plant Patho.* 2003; 25(2):135-142
16. Gurdip S, Maninder S. Chemical control of *Ascochyta* blight of chickpea. *Indian Phytopathol.*1990; 43(1):59-63.
17. Sabiya Bashir, Najeeb M, Sher AD, Seerat, Shafeeq H, Wani RA, Baba JA, Mehfooza H. In vitro efficacy of fungicides and bio-control agents against wilt of chickpea caused by *Fusarium oxysporum* f.sp.ciceris (Foc). *International J Current Microbio and App Sci.* 2017;6(11):1392-1399
18. Bais HP, Fall R, Vivanco JM. Bio-control of *Bacillus subtilis* against infection of *Arabidopsis* roots by *Pseudomonas syringae* is facilitated by biofilm formation and surfactin production. *Plant Physio* .2004;134:307–319.
19. Branda SS, Gonzalez-Pastor JE, Ben-Yehuda S, Losick R, Kolter R. Fruiting body formation by *Bacillus subtilis*. *Proc. Natl. Acad. Sci. U.S.A.* 2001;98:11621–11626.
20. Chen Y, Cao S, Chai Y, Clardy J, Kolter R, Guo JH, et al. A *Bacillus subtilis* sensor kinase involved in triggering biofilm formation on the roots of tomato plants. *Mol Microbio.* 2012;85:418–430.
21. Gong AD, Li HP, Yuan QS, Song XS, Yao W, He W, Jing BZ, Yu CL. Antagonistic mechanism of iturin A and plipastatin A from *Bacillus amyloliquefaciens* S76-3 from wheat spikes against *Fusarium graminearum*. *PLoS ONE.* 2015;10(2): e0116871. Editor; Beom Seok Kim, Korea university, KOREA
22. Stein T. *Bacillus subtilis* antibiotics: structures, syntheses and specific functions. *Mol Microbio.* 2005;56:845–857.
23. Bretag TW, Keane PJ, Price TV. The epidemiology and control of *ascochyta* blight in field peas: a review. *Australian J Agri Res* .2006;57:883–902.
24. Warkentin TD, Rashid KY, Xue AG. Fungicidal control of *ascochyta* blight of field pea. *Canadian J Plant Sci.* 1996; 76:67–71.

25. Warkentin TD, Xue AG, Mcandrew DW. Canadian J Plant Sci. 2000;80:403–406.
Effect of mancozeb on the control of mycosphaerella blight of field pea.

© 2022 Bashir et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/91883>