



Effect of Irrigation Methods, Nutrient Management and Intercropping System on Grain Yield, Maize Equivalent Yield, Protein Content and Economics of Maize (*Zea mays* L.)

Devendra Singh ^a, Mritunjay Kumar ^b, Devilal Birla ^{a*}, Gajanand ^a
and Sameer Shrivastava ^a

^a Department of Agronomy, Dr. Rajendra Prasad Central Agricultural University, India.

^b Department of Agronomy and Project Incharge of Centre for Start Up Facilitation, Dr. Rajendra Prasad Central Agricultural University, Bihar, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2022/v34i232539

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/92999>

Original Research Article

Received 09 September 2022
Accepted 11 November 2022
Published 18 November 2022

ABSTRACT

This study aims to determine Effect of Irrigation Methods, Nutrient Management and Intercropping System on Grain Yield, Maize Equivalent Yield, Protein Content and Economics of Maize (*Zea mays* L.). An experiment was conducted during the *rabi* season of 2020-21 at Tirhut College of Agriculture, Dholi (Dr. Rajendra Prasad Central Agricultural University, Pusa, Bihar). To study the effect of irrigation methods, nutrient management and intercropping system on grain yield, MEY, protein content and economics of maize (*Zea mays* L.). The growing of maize under drip irrigation recorded significantly higher grain yield (9.42 t/ha), MEY (11.67 t/ha), protein content (8.7%), gross returns (215942 ₹/ha), net returns (146679 ₹/ha) and B: C ratio (2.11) over furrow and surface irrigation. Application of nutrients as per STCR, recorded higher grain yield (9.24 t/ha), MEY (11.15 t/ha), protein content (8.57%), gross returns (206343 ₹/ha) and net returns (134893 ₹/ha) as compared to RDF and SSNM whereas, highest B: C ratio was noticed under RDF (1.94). Increment in grain yield (5.50 % and 12.14 %), MEY (18.16% and 18.84%), protein content (4.58 % and

^oM. Sc. Research Scholar;

[#]Professor;

*Corresponding author: E-mail: devilalbirla130596@gmail.com;

7.87%), gross returns (15.85% and 15.37%), net returns (19.8% and 21.59%) and B: C ratio (13.10% and 19.41%) were observed under vegetable pea over lentil and rajmash intercropping respectively. Overall results revealed that irrigation applied through drip method, nutrient management through STCR and maize + vegetable pea intercropping system is a promising option for higher productivity, quality and profitability of maize.

Keywords: Drip irrigation; furrow irrigation; soil test crop response; site-specific nutrient management; maize; intercropping.

1. INTRODUCTION

Globally, the area, production and productivity of maize are 184.8 million hectares (Mha), 1,070 million tonnes (Mt) and 5.62 tonnes/ha (t/ha), respectively (FAOSTAT, 2020).

In India, maize covers a large area of almost 9.86 Mha, with production and productivity of around 28.5 Mt and 2.89 t/ha, respectively [1]. In India, maize is the third most important crop after rice and wheat and contributes nearly 9% to the national food basket [2]. Maize is called the 'queen of cereal' as it is grown throughout the year due to its photo thermo- insensitive character and highest genetic yield potential among the cereals. In India, maize is cultivated for various purposes including grain, feed, fodder, green cobs, sweet corn, baby corn, popcorn and industrial products. For diversifying agricultural production, maize is considered a viable option owing to its wider adaptability in multiple seasons in different ecologies [3]. Maize is emerging as one of the potential crops in rice-based systems that can help to resolve challenges like food and nutritional stability, climate change, water shortage, and other industrial needs [4]. Increased production and profitability are the targets of integrating grain legumes into cereal-based cropping systems to achieve food and nutritional stability as well as long-term sustainability. In addition, legumes play a well-known role in soil fertility restoration and they are also essential sources of protein-rich food, feed, and fodder. Although the intercropping with leguminous crops (cereal + legume) has become common as a way to protect monocropping under rainfed conditions from crop failure. The main purpose of intercropping is to assure a more consistent and long-term yield [5]. Legumes are intercropped with cereals, which helps cereals to benefit from legume nitrogen fixation [6]. The ability of cereal-legume intercropping to provide nitrogen is determined by crop densities, light interception, crop species, and nutrients [7].

In India, generally traditional flood irrigation methods (surface and furrow) are used to irrigate the crops, wherein the entire soil surface is almost flooded without considering the actual consumptive requirements of the crops. These practices have created the problem of waterlogging and salinity and reduction in the overall irrigation efficiency. Therefore, it is needed to adopt modern efficient irrigation methods like drip. The drip irrigation method offers several advantages over surface and furrow irrigation methods including higher crop yields, saving water, water use efficiency and efficient fertilizer application.

Maize is an exhaustive crop, the nutrient requirement cannot be supplied only through native nutrient reserves, and hence the additional nutrient requirements have to be met from fertilizers. Among the several soil test-based fertilizer application techniques, site-specific nutrient management (SSNM) and soil test crop response (STCR) are cost-effective and plant need-based approaches with specific yield targets. The SSNM and STCR approach not only aim to reduce or increase fertilizer use and also the effective tools for supplying crop nutrients as and when needed to achieve higher yield, besides this they also aim to increase system nutrient use efficiency, leading to more net returns per unit of fertilizer invested. In the view of above, the present study was undertaken to assess the effect of irrigation methods, nutrient management and intercropping system on grain yield, maize equivalent yield, protein content and economics of maize.

2. MATERIALS AND METHODS

The experiment was carried out at the Trihut College of Agriculture (TCA) Dholi, Muzaffarpur (Dr. Rajendra Prasad Central Agricultural University, Pusa, Samastipur, Bihar) during the *rabi* season 2020-21. The research farm is located on the banks of the Burhi Gandak River in North Bihar, at a height of 51.2 meters above mean sea level, at 25°59' N latitude and 85°75'

E longitude. The coldest months were December and January, with an average temperature of 11°C, while the hottest months were April and May, with an average temperature of 37°C. The total rainfall during the trial was 4.28 mm. The highest relative humidity was 94 % and the minimum was 73%. The soil status of the field experiment was low in organic carbon (0.43%), available nitrogen (182.46 kg/ha), available potassium (93.29 kg/ha) and medium in phosphorus (21.70 kg/ha), with pH 8.2 which was slightly alkaline. The field experiment was set up in a split-split plot design with three replications. The main plot treatments consisted of three methods of irrigation *i.e.*, surface irrigation, drip irrigation and furrow irrigation, subplot consisted of three doses of fertilizer *i.e.*, recommended dose of fertilizer (RDF) (150:75:50 NPK kg/ha), site-specific nutrient management (SSNM) (120:78:166 NPK kg/ha) and soil test crop response (STCR) (216:140:99 NPK kg/ha) and sub-sub plot consisted three intercrop treatments with maize *i.e.*, lentil, rajmash and vegetable pea.

Protein content in the maize grain was determined by the nitrogen content in the grain and multiplying with the correction factor (6.25). To compute the gross returns, biological yield of maize and multiplied by the price of the produce. The cost of cultivation was subtracted from the actual gross returns to calculate the net return. Maize equivalent yield in (MEY) (t/ha) was calculated by using the following formula:

$$MEY = \text{Maize yield} + \frac{\text{Grain yield of pulse} \left(\frac{t}{ha}\right) \times \text{Price of pulse} (\text{₹/t})}{\text{Price of maize} (\text{₹/t})}$$

3. RESULTS AND DISCUSSION

3.1 Effect on Grain Yield, MEY and Protein Content

The grain yield, MEY and protein content in maize were significantly influenced by irrigation methods, nutrient management and intercropping system (Table 1). The highest grain yield (9.42 t/ha), MEY (11.67 t/ha) and protein content (8.76%) was recorded under drip irrigation which was significantly higher over both furrow and surface irrigation respectively. The maximum yield under drip irrigation might be due to adequate soil moisture availability throughout the crop growth stages and precise water delivery ensures minimal weed growth which resulted in

improved crop growth and yield contributing features, and eventually grain output [8]. On the other hand, lower grain yield is obtained under furrow and surface irrigation due to a large amount of water applied at a time, resulting in leach down of nutrients from the root zone decreasing the growth of the crop and finally reducing grain yield. The highest average maize grain yield was obtained from full irrigation treatment using the drip irrigation method [9] while applications of excessive irrigation water did not increase grain yields [10].

The maximum grain yield (9.24 t/ha), MEY (11.15 t/ha) and protein content (8.57%) was obtained under soil test crop response (STCR) which was significantly higher than the RDF and SSNM. It's possibly due to the higher grain output under STCR being attributable to the use of more fertilizer. As we supplied nutrients as per STCR, the balanced and required quantity application of nutrients according to the crop's needs may have boosted the crop's growth and development, consequently enhancing grain production. However, RDF and SSNM nutrients were found to be inadequate in quantity to fully exploit maize's production potential, as RDF and SSNM yields were much lower than STCR yields [11-13].

Intercropping of maize with vegetable pea recorded higher grain yield (9.12 t/ha), MEY (11.87 t/ha) and protein content (8.51%) which was statistically at par with lentil and significantly superior with rajmash. Grain yield of maize higher under maize + vegetable pea intercropping because vegetable pea harvest (green pods) earlier, after (80-85 DAS) and after harvesting of a vegetable pea, maize growth and development increases due to there is no competition for light, space, water and nutrient resulting enhanced yield attributes and finally increasing grain yield. On other hand, maize + rajmash intercropping significantly lower grain yield as compared to maize + vegetable pea intercropping. This could be because rajmash does not fix atmospheric nitrogen due to non-functional root nodules seen in rajmash, and it is harvested later (110-115 DAS) than a vegetable pea, resulting in lower maize grain yields under maize + rajmash intercropping. A similar response was also reported in *Cajanus Cajan* [14,15].

Table 1. Effect of irrigation methods, nutrient management and intercropping system on yield and quality of maize

Treatments	Grain yield (t/ha)	MEY (t/ha)	Protein content (%)
Irrigation management			
Surface irrigation	7.95	9.72	7.25
Drip irrigation	9.42	11.67	8.76
Furrow irrigation	8.53	10.50	8.18
SEm±	0.22	0.22	0.13
CD (P=0.05)	0.88	0.88	0.51
CV (%)	13.45	10.89	8.28
Nutrient management			
RDF	8.44	10.53	8.06
SSNM	8.22	10.21	7.84
STCR	9.24	11.15	8.57
SEm±	0.21	0.20	0.10
CD (P=0.05)	0.64	0.62	0.32
CV (%)	12.40	9.78	6.78
Intercropping			
Lentil	8.65	9.98	8.12
Rajmash	8.14	10.04	7.84
Vegetable pea	9.12	11.87	8.51
SEm±	0.20	0.20	0.11
CD (P =0.05)	0.56	0.57	0.34
CV (%)	11.76	9.68	7.58

(MEY - Maize Equivalent Yield)

Table 2. Effect of irrigation methods, nutrient management and intercropping system on the economics of maize

Treatments	C.O.C (₹/ha)	Gross returns (₹/ha)	Net returns (₹/ha)	Benefit: Cost ratio
Irrigation management				
Surface	67266	179782	112516	1.67
Drip	69263	215942	146679	2.11
Furrow	70885	194258	123373	1.73
SEm±	-	4123.7	4123.7	0.05
CD (P=0.05)	-	16191.7	16191.7	0.23
CV (%)	-	10.89	16.80	16.64
Nutrient management				
RDF	66226	194805	128579	1.94
SSNM	69738	188834	119096	1.70
STCR	71450	206343	134893	1.88
SEm±	-	3704.7	3704.7	0.05
CD (P=0.05)	-	11415.4	11415.4	0.16
CV (%)	-	9.78	15.09	15.33
Intercropping				
Lentil	66104	184710	118606	1.79
Rajmash	69755	185763	116008	1.66
Vegetable pea	71555	219509	147954	2.06
SEm±	-	3666.6	3666.6	0.05
CD (P=0.05)	-	10516.4	10516.4	0.15
CV (%)	-	9.68	14.94	15.28

(₹ - The Indian rupee)

3.2 Effect on Economics

The economics of maize like gross returns, net returns and B: C ratio were significantly influenced by irrigation methods, nutrient management and intercropping system (Table 2).

The maximum gross returns (215942 ₹/ha), net returns (146679 ₹/ha) and B: C ratio (2.11) were obtained under drip irrigation which was significantly superior over both furrow and surface irrigation respectively. The highest cost of cultivation (70885 ₹/ha) under furrow irrigation followed by drip irrigation (69263 ₹/ha) and the lowest under surface irrigation (67266 ₹/ha) due to the higher number of labors require for bed making and also in drip irrigation higher cost of cultivation because of the initial investment is higher for drip installation but this can be offset by the higher yield of crops resulted in maximum net returns.

Application of nutrient as par STCR, obtained highest gross returns (206343 ₹/ha) and net returns (134893 ₹/ha) which was significantly higher than the RDF and SSNM respectively. However, the highest B: C ratio (1.94) was recorded under RDF which was statistically at par with STCR. This was due to the balanced quantity of nutrients applied through STCR, as we calculate the required quantity of nutrients based on previous history, target yield and soil and climatic conditions.

The highest gross returns (219509 ₹/ha), net returns (147954 ₹/ha) and B: C ratio (2.06) were obtained under maize + vegetable pea intercropping which was significantly superior over both maize + lentil and maize + rajmash intercropping system respectively. The highest cost of cultivation in vegetable pea (71555 ₹/ha) and the lowest in lentil (66104 ₹/ha) intercropping. This could be due to more price of seed, but this was offset by higher crop yield, resulting in greater gross returns, net returns and B: C ratio.

4. CONCLUSION

The results of the present investigation, clearly concluded that among the different Irrigation management practices in maize, drip irrigation, among the nutrient management practices STCR (216:140:99 NPK kg/ha), and among the intercrops vegetable pea was found most effective and remunerative in terms of grain yield, maize equivalent yield, protein content and economics of maize.

ACKNOWLEDGEMENTS

This paper and the research behind it would not have been possible without the exceptional support of my supervisor, Dr. M. Kumar. His enthusiasm, knowledge and exacting attention to detail have been an inspiration and kept my work on track. I thank to and Dr. Rajendra Prasad Central Agricultural University for providing resources for completion my research work.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. FAOSTAT F. Food and Agriculture Organization statistical database; 2020.
2. Dass A, Suri VK, Choudhary AK. Site-specific nutrient management approaches for enhanced nutrient-use efficiency in agricultural crops. *Res Rev J Crop Sci Technol.* 2014;3(3):1-6.
3. Satyanarayana T, Majumdar K, Pampolino M, Johnston AM, Jat ML, Kuchanur P et al. Nutrient Expert™: A tool to optimize nutrient use and improve productivity of maize. *Better crops-South Asia.* 2013;97(1):21-4.
4. Timsina J, Jat ML, Majumdar K. Rice-maize system in South Asia: current status, future prospects and research priorities for nutrient management. *Plant Soil.* 2010;335(1-2):65-82.
5. H. Seran TH, Brintha I. Review on maize-based intercropping. *J Agron.* 2010;9(3):135-45.
6. Ahmad AA, Radovich TJK, Hue NV. Effect of intercropping three legume species on growth and yield of sweet corn (*Zea mays*) in Hawaii. *J Crop Improv.* 2015;29(3): 370-8.
7. Francis CA. Biological efficiencies in multiple-cropping systems. *Adv Agron.* 1989;42:1-42.
8. Fanish SA, Muthukrishnan P, Santhi P. Effect of drip fertigation on field crops-a review. *Agric Rev.* 2011;32(1): 14-25.
9. Yazar A, Sezen SM, Gencil B. Drip irrigation of corn in the Southeast Anatolia Project (GAP) area in Turkey. *Irrig and Drain.* 2002;51(4):293-300.

10. Yildirim E, Kodal S. Effect of irrigation water on corn grain yield in Ankara conditions. Turk J Agric For. 1998;22(6): 65-70.
11. Kumar A, Rana KS, Rana DS, Bana RS, Choudhary AK, Pooniya V. Effect of nutrient- and moisture-management practices on crop productivity, water-use efficiency and energy dynamics in rainfed maize (*Zea mays L.*) + soybean (*Glycine max*) intercropping system. Indian J Agron. 2015;60(1):152-6.
12. Patiram B. Soil Test Crop Response: an effective approach for integrated plant nutrient supply for maize-potato cropping system in inceptisols of Raigarh District of Chhattisgarh plains. Int J Agric Sci. 2018; 10(9):5958-61.
13. Sangtam S, Gohain T, Kikon N. Assessment of nitrogen doses and planting densities for optimizing growth and yield performance of rainfed maize (*Zea mays L.*). Indian J Agric Res. 2017;51(5): 473-7.
14. Krishnaprabu S. Influence on intercropped *Cajanus Cajan* Caused by Land Configuration, Post-Monsoon Irrigation, and P Fertilization. International Journal of Innovative Technology and Exploring Engineering. 2019;46(4):457-29.
15. Pandey IB, Singh SK, Tiwari S. Integrated nutrient management for sustaining the productivity of pigeon pea (*Cajanas cajan L.*)-based intercropping system under rainfed condition. Indian J Agron. 2013; 58(2):192-7.

© 2022 Singh et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/92999>