



Effectiveness of Micronutrients and Seed Coating Polymer Application on Seed Yield and Yield Attributing Parameters of Pigeonpea (*Cajanus cajan* L.)

Mallikarjun G. Handiganoor^{1*}, S. B. Patil² and S. N. Vasudevan¹

¹Department of Seed Science and Technology, College of Agriculture, University of Agricultural Sciences, Raichur – 584 104, Karnataka, India.

²Department of Seed Science and Technology, College of Agriculture, Vijayapur, University of Agricultural Sciences, Dharwad– 580 005, Karnataka, India.

Authors' contributions

This work was carried out in collaboration between all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2018/42985

Editor(s):

(1) Dr. Alejandro Hurtado Salazar, Professor, Departamento de Produccion Agropecuaria, Universidad de Caldas, Colombia.

Reviewers:

(1) Kaouthar Bayahi, University of Tunis el Manar, Tunisia.
(2) Aibuedefe Osagie Eremwanarue, University of Benin, Nigeria.
(3) Bharati Soogareddy, Switzerland.

Complete Peer review History: <http://www.sciencedomain.org/review-history/27421>

Original Research Article

Received 27 June 2018
Accepted 20 November 2018
Published 26 November 2018

ABSTRACT

Aims: To study the Effectiveness of Micronutrients and Seed coating polymer application on seed yield and yield attributing parameters of pigeonpea (*Cajanus cajan* L.)

Place of Study: Field experiment was conducted during *kharif* 2014 at Main Agricultural Research Station, College of Agriculture, University of Agricultural Sciences, Raichur.

Methodology: The different micronutrients viz., Potassium molybdate (2 or 4 ml per kg of seed), ZnSO₄ (2, 4 ml per kg of seed) and boron (2, 4 ml per kg of seed) as per the treatment either individually or in combination were coated with polymer at the rate of 6 ml per kg of seed using rotary seed coating machine, such coated seeds were properly dried under shade. after imposition of treatments, the seeds before sowing were treated with *Rhizobium* and *Trichoderma harizianum* culture. In addition, two foliar sprays as per the treatments either individually or in combination at an

*Corresponding author: E-mail: arjun4726@gmail.com;

interval of 10 days during flowering stage (75 and 85 DAS) were given [Potassium molybdate (0.1%), (Zinc sulphate (0.5 %) in EDTA form, Borax (0.2 %)]. Various observations on seed yield and yield attributing parameters were recorded, analysed statistically to study the effectiveness of Micronutrients and Seed coating polymer application on seed yield and yield attributing parameters of pigeonpea (*Cajanus cajan* L.).

Results: Seed coating polymer (@ 6 ml/kg) of pigeonpea seeds with the combination of potassium molybdate + ZnSO₄ + boron (each @ 2g / kg) of seed along with two foliar sprays of potassium molybdate (0.1 %) + zinc sulphate (0.5 %) in EDTA form + borax (0.2 %) at an interval of 10 days during flowering stage (75 and 85 DAS) recorded significantly maximum number of pods per plant (193.0), maximum number of seeds per pod (3.67), higher seed yield per plant (62.80 g) and finally, highest seed yield (16.30 q) per hectare as compared to control.

Conclusion: Micronutrients (*viz.*, Potassium molybdate, ZnSO₄, boron) and seed coating polymer at optimum doses can stimulate the physiological functions, resulting in early germination, improved seedling vigour with better stand establishment lead to increased productivity of pigeonpea. Hence, It could be advocated for the better establishment of seedlings and higher seed yield.

Keywords: Micronutrient; Pigeonpea; Seed coating polymer; Seed yield.

1. INTRODUCTION

Pigeonpea (*Cajanus cajan*) is a deep rooted hardy crop grown either as sole crop or intercrop in combination with a variety of crops. Initial slow growth habits, wider row spacing and long maturity period of pigeonpea make it ideal as a component crop with most of the cereal or millets, pulses, oilseed and vegetables in rainfed uplands. Its deep root systems help tap plant nutrients from deeper layers allowing the base food crop to feed at top layers of the soil. The pigeonpea based intercropping systems give greater relative yield advantage under stress and can provide useful buffer against low yields. Pigeonpea is one of the important pulse crops of India and 91 per cent of the world's pigeonpea is produced in India. The productivity of pigeonpea in India (799 kg ha⁻¹) is far below the average productivity of world (848 kg ha⁻¹) as per the reports of Ministry of Agriculture. In Karnataka, pigeonpea occupies an area of 0.66 million hectares with an average productivity of 555 kg per hectare [1]. Pigeonpea is grown in almost all the states, and larger portion of the area is in the states like Maharashtra, Uttar Pradesh, Madhya Pradesh, Karnataka and Gujarat. Andhra Pradesh and Tamil Nadu together occupy 87.89 per cent of the area and contribute 86.10 per cent to the total production. Pigeonpea is cultivated for grain purpose as *dhal* which is a major source of protein for poor farmers. It has three times protein as compared to cereals. Tender green seeds are used as vegetable, crushed seeds are used as animal feed, green leaves as fodder, stem is used as fuel wood and to thatch huts, for basket making and fencing and also used to culture the lac insects. It is also

often used as a live fence around small farms. It is grown across the mountain slopes to reduce soil erosion. The low yield of pigeonpea is mainly attributed to their cultivation on poor soils with inadequate and imbalanced nutrient application without the application of macronutrients and micronutrients like boron, zinc and iron.

Seed coating (polymerisation) is a technique wherein any substance applied to the seed does not obscure or change its shape. The film is readily water soluble (hydrophilic) so as not to impede seed germination [2]. In this method small quantity of chemicals or micronutrients are needed as compared to soil application or foliar spray. Seed polymerisation is one of the most economical approaches for improving seed performance. Film coating helps to smoothen the seed surface which improves ability of flow and helps in mechanised planting [2]. The improvements in crop establishment, growth and yield due to coating or pelleting have been reported in several agricultural, horticultural and tree crops [2-3].

The use of micronutrients in pulses production is one of the ways to boost up the productivity [4]. Zinc plays an important role in the formation of chlorophyll and growth hormones and is associated with the uptake of water [5]. Boron plays an important role in calcium metabolism, its uptake and efficient use, which is the important constituent of cell wall [6]. Molybdenum plays the key role in process of atmospheric nitrogen fixation besides role in enzyme systems [7]. The yield and yield attributing characters are directly related to the productivity and hence the micronutrients like zinc, boron and molybdenum

are important in the production of legume crops [8].

Pigeonpea is an important pulse crop of rainfed agriculture and a principal source of protein in the Indian diet. Rainfed soils are generally degraded with poor native fertility [9]. Mineral nutrient deficiencies limit nitrogen fixation by the legume-rhizobium symbiosis, resulting in low legume yields. Nutrient limitations to legume production result from deficiencies of not only major nutrients but also micronutrients such as molybdenum (Mo), zinc (Zn), boron (B) and iron (Fe) [10]. Inadequate nodulation of pigeonpea can be associated with low plant available Mo. Increase in flower numbers, pod set improvement, and reduction in days to flowering are influenced by Mo [10]. Application of micronutrients along with recommended doses of fertilizers (RDF) to pigeonpea is essential for higher yield under rainfed conditions. The micronutrients may be supplied to the plants through soil application, foliar spray or applied to seed through seed treatment [8]. Although the required amount of micronutrients can be supplied by any of these methods, foliar sprays have been more effective in yield improvement and grain enrichment [11]. Application of micronutrients through foliar spray is taken up at later growth stages when crop stands are already established [8]. Hence, an alternative acceptable method of supplying micronutrients during the early stage of seedling establishment is the need of the hour.

Very little work has been carried out regarding the response of pigeonpea to the application of seed coating polymer and foliar spray with

micronutrients. Hence, the present investigation entitled "Effectiveness of Micronutrients and Seed coating polymer application on seed yield and yield attributing parameters of pigeonpea (*Cajanus cajan* L.) was undertaken.

2. MATERIALS AND METHODS

The experiment was conducted during *kharif* 2014 at Main Agricultural Research Station, College of Agriculture, University of Agricultural Sciences, Raichur, India. Geographically, the station is situated in the North-Eastern dry zone (Zone- 2) of Karnataka State at 16° 15' North latitude and 77° 20' East longitude and at an altitude of 389 meter above mean sea level (Table 1.). The soil of the experimental site was deep black, clay in nature and 8.3 pH [12 &13] (Table 2.). The experiment consisted of 16 different treatments viz., T₁: Potassium molybdate @ 2g per kg of seed, T₂: Potassium molybdate @ 4g per kg of seed, T₃: ZnSO₄ @ 2g per kg of seed, T₄: ZnSO₄ @ 4g per kg of seed, T₅: Boron @ 2g per kg of seed, T₆: Boron @ 4g per kg of seed, T₇: Potassium molybdate + ZnSO₄ (each @ 2g / kg of seed), T₈: Potassium molybdate + ZnSO₄ (each @ 4g / kg of seed), T₉: ZnSO₄ + Boron (each @ 2g / kg of seed), T₁₀: ZnSO₄ + Boron (each @ 4g / kg of seed), T₁₁: Potassium molybdate + Boron (each @ 2g / kg of seed), T₁₂: Potassium molybdate + Boron (each @ 4g / kg of seed), T₁₃: Potassium molybdate + ZnSO₄ + Boron (each @ 2g / kg of seed), T₁₄: Potassium molybdate + ZnSO₄ + Boron (each @ 4g / kg of seed), T₁₅: Only polymer and T₁₆: Absolute control laid out in randomized block design with three replications.



Fig. 1. Pigeonpea seeds coated with polymer and micronutrients

Table 1. Monthly meteorological data for the experimental year 2014-2015 and mean of the last 81 years (1932-2013) at Meteorological Observatory, Main Agricultural Research Station, University of Agricultural Sciences, Raichur

Month	Rainfall (mm)		Temperature (°C)				Relative humidity (%)	
	1932-2013	2014-15	Mean maximum		Mean minimum		1932-2013	2014-15
			1932-2013	2014-15	1932-2013	2014-15		
April	70.70	19.40	39.9	39.32	22.6	24.77	77.00	40.17
May	71.50	93.00	39.7	39.08	22.5	24.68	80.00	47.76
June	182.70	48.40	35.3	37.76	22.3	24.68	82.00	52.60
July	62.50	123.10	33.4	33.74	20.5	23.08	79.00	68.24
August	21.20	372.90	32.9	32.70	19.1	22.71	79.00	71.00
September	4.00	102.70	32.2	31.01	16.2	22.48	76.00	75.37
October	1.20	50.60	31.5	31.99	16.8	21.32	77.00	66.39
November	1.10	12.00	31.3	30.90	18.5	17.50	62.00	62.57
December	44.30	1.80	30.5	29.41	22.6	15.11	56.00	64.45
January	12.95	10.00	31.3	30.25	24.4	17.23	53.00	58.84
February	42.90	0.00	32.5	32.65	25.3	18.29	60.00	57.21
March	113.80	24.90	36.5	35.25	23.3	21.81	79.00	50.66
Total	628.85	875.30	-	-	-	-	-	-

Table 2. Physical and chemical properties of the soil in experimental site

Properties	Value obtained	Method adopted
I. Physical properties		
1. Particle size analysis		
Sand (%)	20.18	International Pipette method (Piper, 1966, Jena et al., 2013)
Silt (%)	23.02	
Clay (%)	56.80	
Texture	Clay	
2. Bulk density (Mg m ⁻³)	1.28	Core Sampler method (Dastane, 1967)
II. Chemical properties		
Soil reaction (pH)	8.30	Potentionmeter (Jackson, 1967)
Electrical conductivity (dS m ⁻¹)	0.25	Conductivity bridge (Jackson, 1967)
Organic carbon (%)	0.60	Wet Oxidation method (Jackson, 1967)
Available nitrogen (kg ha ⁻¹)	218.00	Alkaline permanganate method (Subbaiah and Asija, 1956)
Available phosphorus (kg ha ⁻¹)	32.00	Olsen's method (Jackson, 1967)
Available potassium (kg ha ⁻¹)	180.67	Flame photometry method (Jackson, 1967)

Table 3. Influence of seed coating polymer and micronutrients with foliar spray on seed yield and yield attributing parameters of pigeonpea

Treatment	No. of pods/plant	No. of seeds/pod	Seed yield (g)/plant	Seed yield (q)/ha
T ₁	179.0	3.10	54.88	14.20
T ₂	183.0	3.21	57.50	14.46
T ₃	174.2	2.98	53.60	14.10
T ₄	181.0	3.16	56.12	14.15
T ₅	174.3	3.00	54.00	14.10
T ₆	181.7	3.10	56.33	14.30
T ₇	186.0	3.26	59.05	14.95
T ₈	185.0	3.22	58.80	14.92
T ₉	184.3	3.18	57.93	14.86
T ₁₀	184.0	3.21	57.50	14.20
T ₁₁	187.2	3.32	59.60	15.17
T ₁₂	187.6	3.30	59.40	15.06
T ₁₃	193.0	3.67	62.80	16.30
T ₁₄	188.0	3.32	60.08	15.20
T ₁₅	173.0	2.97	53.42	14.03
T ₁₆	169.0	2.93	53.20	13.86
Mean	181.9	3.17	57.14	14.62
S.Em.±	0.96	0.04	0.64	0.14
CD (P = 0.05)	2.77	0.13	1.84	0.40

The seeds of pigeonpea variety TS 3R were sown with a spacing of 90 cm between rows and 30 cm between plants with the plot size was 9.0 x 4.8 m². The fresh seeds of pigeonpea were treated with different micronutrients as per above treatments through seed coating polymer at the rate of 6 ml per kg of seed which was standardised from laboratory experiment using rotary seed coating machine and the coated seeds were properly dried under shade (Fig. 1). In addition to seed coating polymer, two foliar sprays of micronutrients viz., zinc sulphate (0.5%), potassium molybdate (0.1%) in EDTA form and borax (0.2%) in non EDTA form were given during flowering stage (75 and 85 DAYS) at an interval of 10 days either individually or in combination as per the treatments.

The various observations viz., number of pods per plant, number of seeds per pod, seed yield per plant and seed yield per hectare were recorded. The research data was statistically analysed by adopting the procedure prescribed by Panse and Sukhatme [14].

3. RESULTS AND DISCUSSION

Though micronutrients are required in small amounts but play a key role in biochemical activities in plant. Seed coating polymer with

micronutrients and foliar spray significantly influenced the seed yield and yield attributing parameters (Table 3). The number of pods per plant is a major yield attributing component contributing to the final seed yield. In the present investigation among the different treatments of seed coating polymer with micronutrients and foliar spray, the treatment T₁₃ [Potassium molybdate + ZnSO₄ + boron (each @ 2g / kg seed)] along with two foliar sprays of potassium molybdate (0.1%) + zinc sulphate (0.5%) in EDTA form + borax (0.2%) recorded significantly maximum number of pods per plant (193.0). Whereas, minimum number of pods (169.0) per plant was recorded in the control. This increase in number of pods per plant might be due to higher number of primary and secondary branches per plant and also due to higher values for various physiological growth parameters registered by this treatment [15]. Similar results were reported by Dixit and Elamathi [16] in cowpea due to seed fortification and pelleting with the combination of ZnSO₄ (0.2%) + MnSO₄ (0.2%) + Na₂Mo₄ (0.1%) per kg seeds. John et al. [17] also reported the same in green gram due to foliar application of DAP (2%) + NAA (40 ppm) + B (0.2 %) + Mo (0.05 %). As a result of stimulated physiological functions due to imposed treatments and foliar spray the number of seeds per pod was also differed significantly. However, T₁₃ recorded maximum number of

seeds per pod (3.40) as compared to control (2.93) i.e. T₁₆.

Seed coating polymer with micronutrients and foliar spray significantly influenced the seed yield per plant and seed yield per hectare. The treatment T₁₃ [Potassium molybdate + ZnSO₄ + boron (each @ 2g / kg seed)] along with two foliar sprays of potassium molybdate (0.1%) + zinc sulphate (0.5%) in EDTA form + borax (0.2%) produced higher seed yield per plant (62.80 g) and seed yield per hectare (16.30 q) which were found to be superior over all the treatments. Whereas, the lowest seed yield per plant (53.20 g) and seed yield per hectare (13.86 q/ha) was recorded in the control (T₁₆). Polymer present in the coating material might have also helped in higher rate of water uptake, in turn, resulted in the early germination with more seedling vigour and better stand establishment, which might have ultimately led to better growth, plant stand and productivity of pigeonpea and increase in plant height, number of pods per plant and test weight as a consequent of improvement in root growth and nodulation [18]. The reason for the increased yield might also be due to the increased photosynthetic efficiency through stabilisation of chlorophyll, higher production of photosynthesis resulting in increased translocation of nutrients from the source to sink in the treated plants. The increase in seed yield due to combined application of zinc, boron and potassium molybdenum might be attributed to the complementary effect of these nutrients with each other and along with a standard dosage of polymer ultimately, resulting in the higher growth and seed yield. Similar findings were observed by Duingqiong et al. [19] where they observed beneficial effects of micronutrients (zinc, manganese, iron, copper, boron and molybdenum) and polymer together in getting better yield in maize. As Mo being essential for Nitrogen fixation, which has ensured better Nitrogen supply to the crop while Zn being an activation of several enzyme systems might have improved physiological functions of the plant and conversion of flower to pod which resulted in better growth and yield of the crop. These results are in accordance with the findings of Sharma et al. [20] in groundnut who observed similar increase in the yield due to seed treatment with Zn+Mo at the rate of 8 g per kg of seed. Similarly, boron is directly linked with the process of pollen producing the capacity of anther, viability of pollen grains, pollen germination and pollen tube growth and

fertilisation [21]. Dinesh [22] also reported a significant increase in seed yield and its attributes when seeds were pelleted with boron alone, Zinc + Boron + arappu leaf powder. Shil et al. [23] observed that combined application of both boron and molybdenum were superior to their single application in chickpea. In the same line [22] also noticed an increased yield of soybean due to the combined application of Zn, B and Mo at 2 kg, 1.0 kg and 0.5 kg respectively with *Bradyrhizobium* biofertiliser.

However, from the present study it is clear that there was a significant increase in the yield and yield attributing characters due to seed coating polymer with micronutrients i.e. zinc, boron and potassium molybdate sole or in combination but the sole treatment of polymer at the rate of 6 ml per kg was found to be similar with T₁₆ (control) in terms of number of seeds per pod, seed yield per plant and per hectare. These results are in conformity with the findings of Vinodkumar et al. [24] in pigeonpea where they recorded statistically similar values for plant height, number of primary and secondary branches, number of pods per plant, pod weight per plant, seed yield per plant and seed yield per hectare with seed treatment of polymer at the rate of 5 ml per kg of seed as compared to control. Hence, in general it is important to note that the polymer is more effective when used in combination with micronutrients, biofertilisers and fungicides than sole. Similar findings were reported by [25] in sunflower.

4. CONCLUSION

Seed coating polymer and micronutrients with foliar spray in combination resulted in better seed yield and yield attributing parameters. Seed coating polymer (6 ml per kg of seed) with the combination of micronutrients namely, potassium molybdate + ZnSO₄ + boron each at 2g per kg of seed with two foliar sprays (0.1 % + 0.5 % + 0.2 % respectively, potassium molybdate and ZnSO₄ in EDTA form) at an interval of 10 days during flowering stage (75 and 85 DAS) can be recommended as the optimum dose for pigeonpea cultivation during *kharif* season.

ACKNOWLEDGEMENT

We thankfully acknowledge the contribution of the reviewers.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Anonymous. All India area, production and yield of total pulses. 2010-11. Ministry of Agriculture, Govt. of India; 2010. Retrieved from www.indiastat.com
2. Punithavathi M. Seed hardening and pelleting studies in ragi Cv. Co-13. M. Sc (Agri.) Thesis, Tamil Nadu Agric. Univ. Coimbatore (India); 1997.
3. Malla Reddy M, Padmaja B, Malathi S, Jalapathi Rao L. Effects of micronutrients on growth and yield of pigeonpea. Journal of SAT Agricultural Research. 2016;14:1-6.
4. Gowda CL, Samineni S, Gaur PM, Saxena KB. Enhancing the productivity and production of pulses in India; 2013.
5. Mousavi SR, Mohammad Galavi M, Rezaei M. The interaction of zinc with other elements in plants: A review. Intl J Agri Crop Sci. 2012;4(24):1881-1884.
6. Naghii MR, Samman S. The role of boron in nutrition and metabolism. Prog Food Nutr Sci. 1993;17(4):331-49.
7. Weisany W, Raei Y, Allahverdipoor KH. Role of some of mineral nutrients in biological nitrogen fixation. Bulletin of Environment, Pharmacology and Life Sciences. 2013;2(4):77-84.
8. Farooq M, Wahid A, Siddique KH. Micronutrient application through seed treatments: A review. Journal of Soil Science and Plant Nutrition. 2012;12(1): 125-142.
9. Bhuiyan MAH, Khanam D, Ali MY. Chickpea root nodulation and yield as affected by micronutrient application and rhizobium inoculation. International Chickpea and Pigeonpea Newsletter. 1999;6:28-29.
10. Prasad M, Pillai A, Faber G, Field S, Dowling T, Prasad M, Pillai A, Faber G, Field S, Dowling T. Molybdenum response of pigeonpea on Ferruginous Latosols in Fiji. International Chickpea and Pigeonpea Newsletter. 1998;5:55-56.
11. Arif M, Chohan MA, Ali S, Gul R, Khan S. Response of wheat to foliar application of nutrients. Journal of Agricultural and Biological Science. 2006;1(4):30-34.
12. Jena RK, Jagadeeswaran R, Sivasamy R. Analogy of soil parameters in particle size analysis through laser diffraction techniques. Indian Journal of Hill Farming. 2013;26(2):78-83.
13. Jankowski SJ, Freiser H. Flame Photometric Methods of Determining the Potassium Tetraphenylborate. Analytical Chemistry. 1961;33(6):773-775.
14. Panse VG, Sukhatme PV. Statistical Methods for Agricultural Workers. ICAR-New Delhi (India). 1985;167-174.
15. Ponnuswamy AS, Vijaya J. Effect of seed fortification and pelleting on crop growth and yield in cowpea. Madras Agric. J. 1997;84(7):395-431.
16. Pradeep Mohan Dixit and Elamathi, S. Effect of foliar application of DAP, micronutrients and NAA on growth and yield of green gram (*Vigna radiata* L.). Legume Res. 2007;30(4):305-307.
17. Sherin Susan John, Bharati, Natesan P, Raja K, Seed film coating technology for maximizing the growth and productivity of maize. Karnataka. J. Agric. Sci. 2005; 18(2):349-356.
18. Geetha KN, Shankar AG, Shivashankar K. Effect of Mo, Zn and Ca on productivity of groundnut. J. Oilseeds Res. 1995;13(2): 167-172.
19. Duyingqiong Q, Liao Xingrong, He Jianghua, Hiang Zhoyao, Zhou Xiaohong. Effect of B and Mo on the growth, development and yield of peanut. Plant nutri. Fert. Sci, 2002;8(2):233-35.
20. Sharma A, Nakul HT, Jelgeri BR, Ashok Surwenshi. Effect of micronutrients on growth, yield and yield components in pigeonpea (*Cajanus cajan* (L.) Millsp.). Res. J. Agric. Sci. 2010;1(2):142-144.
21. Masuthi DA, Vyakaranahal BS, Deshpande VK. Influence of pelleting with micronutrients and botanical on growth, seed yield and quality of vegetable cowpea. Karnataka J. Agric. Sci. 2009;22: 898-900.
22. Dinesh Chandra R. Effect of biofertilizer, zinc, boron and molybdenum on nodulation, growth and yield of soybean. M. Sc. (Agri.) Thesis, Bangladesh Agricu. Univ. Mymensingh (Bangladesh); 2014.
23. Shil NC, Noor S, Hossain MA. Effect of boron and molybdenum on the yield of chickpea. J. Agric. Rural Dev. 2007;5(1-2): 17-24.

24. Vinodkumar SB, Vyakarnahal BS, Dhananjaya P, Yegappa Hipparagi, Asha AM. Effect of seed polymer coating on field performance and quality of pigeonpea (*Cajanus cajan* L.). Environ. Eco. 2013;31(1):43-46.
25. Shakuntala NM, Vyakaranahal BS, Shankergoud I, Deshpande VK, Pujari BT, Nadaf HL. Effect of seed polymer coating on growth and yield of sunflower hybrid - RSFH-130. Karnataka J. Agric. Sci. 2010;23(5):708-711.

© 2018 Handiganoor et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sciencedomain.org/review-history/27421>