



Effect of Molybdenum Enzyme Activity of Green Gram (*Vigna radiata*) Grown in Low Land Acid Soils of Pudukottai Region of Tamil Nadu

J. Karpagam^{1*} and Rajesh²

¹Department of Soil Science and Agriculture Chemistry, N. S. College of Agriculture, Markapur 523329, Acharya N. G. Ranga Agriculture University, Lam 522034, Andhra Pradesh, India.

²Department of Agronomy, Tamil Nadu Agricultural University, Coimbatore-641003, Tamil Nadu, India.

Authors' contributions

This work was carried out in collaboration between both authors. Author JK designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Author Rajesh managed the literature searches. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2018/44298

Editor(s):

(1) Dr. Francisco Cruz-Sosa, Department of Biotechnology, Metropolitan Autonomous University Iztapalapa Campus, Av. San Rafael Atlixco 186 México City, México.

Reviewers:

(1) V. Vasanthabharathi, Annamalai University, India.

(2) Júnior Melo Damian, University of São Paulo, Brazil.

(3) Miguel Aguilar Cortes, Universidad Autonoma Del Estado De Morelos, Mexico.

Complete Peer review History: <http://www.sciencedomain.org/review-history/27505>

Original Research Article

Received 01 September 2018
Accepted 13 November 2018
Published 30 November 2018

ABSTRACT

The role of molybdenum (Mo) in greengram in regulating the activities of enzymes such as nitrate reductase (NR) and glutamine synthetase (GS) in particular to nitrogen assimilation was investigated in acid soils of Semi-arid Tropics (SAT) of Tamil Nadu. Incubation studies were taken up with two sources of molybdenum namely sodium molybdate (Na_2MoO_4) and ammonium molybdate ($(\text{NH}_4)_6\text{Mo}_7\text{O}_{24}$) in which sodium molybdate proved to be a better performer and recommended as Mo fertiliser in SAT acid soils. Application of Mo increases root, shoot length and biomass. It also increases the number of nodules and rhizobium population in green gram. In particular to microbial aspect the rhizobium population count serial dilution technique was followed. The anatomy of the nodule was imaged using high-resolution imaging Scanning Electron Microscope (FEI SEM Quanta 250, Netherlands).

Keywords: Nitrate reductase; glutamine synthetase; greengram; molybdenum.

*Corresponding author: E-mail: karpagam13@yahoo.com;

1. INTRODUCTION

Pulses are the important protein source for human diet. In India pulses can be cultivated with limiting resources which is less expensive than animal protein. Pulses are less expensive when compared to other vegetables which can be grown in rainfed conditions as intercrop as well as a mixed crop in particular of giving better returns. Apart from consumption pulses crop cultivation improves soil fertility, the physical structure of soil and also act as nutrition fodder for cattle. The pulse availability per adult per day at present is only 36 which is against the minimum requirement of 50g/adult/day. The recommended dietary allowances (RDA) for adult male and female is 60 g and 55g per day. Greengram in Indian soils considered an excellent source of high quality and digestible proteins but deficient in important micronutrients viz., Mo, Zn, Fe. Pulses are very exact in their requirement like neutral soil pH (liming acid soils), phosphate and sulphur manuring, use of Rhizobium culture and molybdenum for seed treatment for better nodule activity to achieve maximum yield. Pulses are cultivated in two seasons viz., Kharif and Rabi season with a production area of 143.41 lakh hectare, the productivity of 91.16 lakh tonnes and yield of 636 Kg per hectare [1]. Even India increased output of pulse production to 22.95 million tonnes from 16.35 million tonnes, need arises to import 50.8 lakh tonnes from China, Canada, Australia and Myanmar which costs about Rs 17 280 crores [2].

Pulses contribute 11 percent of total gross cropped area. Among all pulses, the area under gram is 4% arhar 2% and other pulses in about 5% of the gross cropped area. The net irrigated area in the country is 47% while the remaining falls under rainfed conditions. The pulses under irrigation cultivated about 37% of the area while 63% of pulses are grown under rainfed conditions. India's outstanding contribution towards total global acreage and production of pulses at 35% & 25% respectively.

The biological importance of Mo in plants is due to its highly beneficial action in the fixation of nitrogen from the atmosphere by the nitrogen-fixing bacterium (*Azotobacter chroococum*). Some of the crops considered most sensitive to Mo deficiency are clover, cauliflower, broccoli, rape, beet, spinach, lettuce and alfalfa. Among the micronutrients, Mo is readily translocated,

and its deficiency symptoms generally appear on the whole plant. The deficiency for other micronutrients appears on the young leaves at the top of the plant because of their inability to translocate within the plant. Mo deficiency emerges as general yellowing and stunting of the plant, interveinal mottling and cupping of the older leaves followed by necrotic spots at leaf tips and margins. Mo deficiency reduces the availability of Fe and P and the utilisation of N, increases the contents of S, Cu and Mn in plants [3]. During the growth of pulses, atmospheric nitrogen get fixed in their root nodules by Rhizobium. Mo helps in localisation and regulation of nitrogen assimilation and amino acid accumulation via regulating enzymes [4], and also is a co-factor of nitrate reductase (NR) enzyme. NR is required for the reduction of nitrates for protein synthesis. Mo also stimulates the activity of glutamine synthetase [5]. Glutamine synthetase (GS) converts ammonia into protein via GS-GOGAT cycle. Thus Mo deficiency reduces NR and GS activities. Mo deficiency occurs mainly in acidic soils of temperate climatic conditions or in hilly acidic soils where the soil is usually acidic. In contrast, in some of the laterite low land acidic soils of Pudukottai and Sivagangai Districts of Tamil Nadu of southern India under SAT (semi-arid tropical) climatic conditions, Mo deficiency is very common. This study was undertaken to investigate the physiological response of greengram on the application of Mo fertiliser in acidic low land soil and the pathways of N-accumulation and utilisation for the same.

2. MATERIALS AND METHODS

Pudukottai District covers a geographical area of 4,56,449 ha receiving rainfall of 685mm from northeast monsoon season falls under the temperature range of 32.8°C to 17.3°C. Pudukottai extends across the north latitude of 8° 30' to 10° 40' Eastern latitude of 78°24' to 79°40'. Pudukottai District consists mainly of Cretaceous type of geological distribution which consists of sand, clay, sandy-clay lying over the crystalline rocks of Archean age in parts of Pudukottai district. The soil is classified taxonomically as Paralithic Haplustalf, In particle size analysis as per international pipette method the percentage of sand (56%), silt (15%) and clay (28%). The pH (5.61) is acidic in nature, organic carbon content (0.36) is low and calcium carbonate (3.22%) is non-calcareous.

2.1 Incubation Experiment

A laboratory experiment was conducted to study the solubility of molybdenum fertilisers namely sodium molybdate and ammonium molybdate at varying doses. There were five levels of Mo (0, 1, 2, 3, 4 and 5 kg Mo ha⁻¹) replicated four times in completely randomised design (CRBD). The factors used for the incubation study were:

- i) Sources of Mo : sodium molybdate and ammonium molybdate
- ii) Levels of molybdenum (Six) : 0, 1, 2, 3, 4 and 5 kg Mo ha⁻¹

Sodium molybdate of 0.64 gram was dissolved in 1000 ml double distilled water which gives 640 mg L⁻¹. From the stock solution of 640 mg L⁻¹, 1, 2, 3, 4, 5 ml was pipetted out and made up the volume to 500 ml in order to mix the solution with the soil sample taken for incubation study. 0.55 gram of ammonium molybdate was dissolved in 1000ml double distilled water which gives 550 mg L⁻¹. From the stock solution of 550 mg L⁻¹, 1, 2, 3, 4, 5 ml was pipetted out and made up the volume to 500 ml to mix the solution with the soil sample taken for incubation study (Table 1).

2.1.1 Incubation containers

Clear white plastic jars (14.0 cm height, 12.0 cm diameter and 1.0 kg capacity) fitted with tight lids were selected for conducting the experiment.

2.1.2 Incubation study

The incubation process was done physically by taking 1 kg of soil into plastic cups and two molybdenum sources such as sodium molybdate and ammonium molybdate applied at the rate of 1, 2, 3, 4 and 5 kg Mo ha⁻¹. Sodium and ammonium molybdate were thoroughly mixed and brought to the required moisture level with deionised water to get the moist friable condition of the soil. The plastic cups having fertiliser were covered with tight lids and kept in the laboratory at room temperature for thirty days.

To facilitate the normal chemical process, the soil – molybdenum fertiliser sources were stirred with a glass rod once a day. To maintain sufficient moisture level in each cup throughout the incubation period, the plastic cups were weighed every three days and the weight was recorded and the difference between the last and current weight was compensated by adding deionised water.

Soil samples were drawn at the beginning, 1st, 2nd, 3rd and 4th week of incubation for analysis of molybdenum. After sampling, the cups were weighed in order to add water to keep the moisture level constant in each cup.

2.2 Basic Studies

A pot culture experiment was conducted at the Department of Soil Science and Agricultural Chemistry in order to study the N-fixation process in greengram plants. Soil samples drawn from NPRC, Vamban was used for the study. Treatments composed of 5 levels of molybdenum (0, 200, 400, 600, 800 and 1000 g Mo ha⁻¹) replicated 5 times in a completely randomised design. During the experimentation, nodules count, Rhizobium population, N-fixation process and high-resolution imaging of nodules were taken up.

2.2.1 Nodulation number

At 40 DAS, plants were uprooted, washed thoroughly and the total nodule count was taken up. The data were presented as a number of nodules plant⁻¹.

2.2.2 Rhizobium population (Serial Dilution technique)

One gram of the soil sample was transferred aseptically to a sterile water blank of 9 ml to get 10⁻¹ dilution. One ml of this dilution was transferred to 9 ml sterile blank to get 10⁻² dilution. Similarly, the sample was serially diluted up to 10⁻⁹ dilution and one ml of this dilution was used for plating.

Table 1. Different dose levels of sodium molybdate and ammonium molybdate

Treatments	Sodium molybdate kg ⁻¹ soil	Ammonium molybdate kg ⁻¹ soil
1 kg Mo ha ⁻¹	1.25 mg (L ₁)	0.925 mg (M ₁)
2 kg Mo ha ⁻¹	2.50 mg (L ₂)	1.85 mg (M ₂)
3 kg Mo ha ⁻¹	3.75 mg (L ₃)	2.775 mg (M ₃)
4 kg Mo ha ⁻¹	5.0 mg (L ₄)	3.700 mg (M ₄)
5 kg Mo ha ⁻¹	6.25 mg (L ₅)	4.625 mg (M ₅)

2.3 Nitrate Reduction Activity

Nitrate reductase enzyme was determined after 30th day of the vegetative stage by following the method developed by Hageman et al. [6]. 0.5 ml phosphate buffer (pH 7.5) was pipetted out in a test tube. Then 0.2 ml of potassium nitrate solution, 0.4 ml NADH solution and 0.7 ml of water was added. The reaction was initiated by the addition of 0.2 ml enzyme extract, similarly, a control was set up in the same way but it should be done with water instead of enzyme extract. Incubation was done at 30°C for 15 minutes. The reaction was terminated by the rapid addition of 1 ml of sulphaniilamide followed by 1ml of naphthyl ethylenediamine reagent. It should be kept for 90 minutes. The absorbance was measured at 540 nm. A standard graph was prepared with potassium nitrite. Different known aliquots of potassium nitrite standard solution were pipette into a series of test tubes and the volume was made up in each tube to 2 ml by adding water.

2.4 Glutamine Synthetase

Glutamine synthetase enzyme was determined after 30th day of vegetative stage by following the method developed by Pateman [7]. The reagent (ml) was pipetted out as mentioned in the order below : Glutamine 2.0 ml, Sodium arsenate 0.5 ml, MnCl₂ 0.3 ml, Hydrxylamine 0.5 ml, ADP 0.5 ml, enzyme extract 0.2 ml.

In order to set a blank 2 ml of 20 mM Tris –HCl was added instead of Glutamine. The reaction mixture was incubated for 30 minutes at 37°C. The reaction was stopped by adding 1 ml of ferric chloride reagent. The brown colour was measured at 540 nm. A range of standards containing 100-500 µg γ-glutamyl hydroxamate in 4 ml buffer solution was prepared and the colour was developed by adding 1 ml of ferric chloride.

2.5 Scanning Electron Microscope (SEM)

The scanning electron microscope (SEM) is a type of electron microscope that images the sample surface by scanning it with a high-energy beam of electrons in a raster scan pattern. Here, a wide range of magnifications is possible, from about 10 times (about equivalent to that of a powerful hand-lens) to more than 500,000 times.

Nodules were removed from the roots of green gram plant. The nodules were washed with distilled water for 4 to 5 times to remove the soil particles. Then the moisture was removed from

the nodule with the help of tissue paper. Then the cross section of the nodule was taken and loaded on the sample stage and images were taken in a Scanning Electron Microscope (FEI SEM Quanta 250, Netherlands).

2.6 Statistical Analysis

The data collected were subjected to statistical analysis in ANOVA [8]. Whenever the treatment difference was found significant (F test), critical difference was worked out at 5 per cent probability level and the values were furnished. If there is no significant difference between treatments, it was denoted by the symbol NS.

3. RESULTS AND DISCUSSION

3.1 Mo Release Pattern from Two Sources of Mo Fertilisation

In order to study the Mo release pattern from two sources of Mo fertilisers sodium molybdate and ammonium molybdate and release of Mo were periodically monitored for a duration of 4 weeks. An incubation experiment was conducted for a period of 4 weeks. The results showed that the AB-DTPA extractable Mo showed an increasing trend with the progression of the incubation period (Table 2 and Fig. 1). The values increased progressively with an advancement incubation period regardless of sources of Mo fertiliser. However, the values were 10 to 15 per cent higher for sodium molybdate than ammonium molybdate at all stages of observations. Mo fertilisation at the rate of 1000 g ha⁻¹ registered highest values 4 to 5 times higher than control and such response continued in the incubation period. At the last week of incubation, the available Mo status of M5 applied at the rate of 1000 g Mo ha⁻¹ registered 79.3 and 62.3 µg g⁻¹ for sodium and ammonium molybdate, respectively. The available Mo value was nearly doubled in fertilised soils in comparison to an initial value in the experimental site. The basic solubility had shown that ammonium Mo solubility is enhanced when the medium gets heated up. The sodium molybdate is soluble under room temperature and therefore it is ideal as a recommended source of Mo fertiliser under field conditions. Performance of sodium molybdate (Na₂MoO₄) was better than ammonium molybdate [(NH₄)₂MoO₄] in terms of available Mo. Na₂MoO₄ is readily soluble at room temperature because its mineralisation occurs at a faster rate than (NH₄)₂MoO₄. So, Na₂MoO₄ is recommended as Mo fertiliser source.

Table 2. Molybdenum release pattern from two sources of Mo fertilisers

Treatments	0 DAS		1 st week		2 nd week		3 rd week		4 th week	
	NaMoO ₄	NH ₄ MoO ₄	NaMoO ₄	NH ₄ MoO ₄	NaMoO ₄	NH ₄ MoO ₄	NaMoO ₄	NH ₄ MoO ₄	NaMoO ₄	NH ₄ MoO ₄
Mo (0 g Mo ha ⁻¹)	0.04	0.03	0.06	0.04	0.08	0.06	0.1	0.08	0.13	0.09
M1 (200 g Mo ha ⁻¹)	10.8	9.2	14.3	12.3	19.0	16.2	24.7	22.3	28.7	26.3
M2 (400 g Mo ha ⁻¹)	15.1	13.3	21.3	19.5	25.3	23.4	30.3	28.1	34.5	32.7
M3 (600 g Mo ha ⁻¹)	19.2	17.2	26.3	24.8	32.4	30.3	37.6	34.6	40.2	37.3
M4 (800 g Mo ha ⁻¹)	35.5	28.5	42.6	33.4	46.7	37.5	50.1	43.7	53.7	47.3
M5 (1000 g Mo ha ⁻¹)	44.8	37.6	48.5	41.3	59.3	44.5	65.3	58.7	79.3	62.3
Average	20.95	17.66	25.54	21.92	31.69	25.36	35.93	31.27	40.41	34.37
SEd	0.17	0.02	0.02	0.01	0.02	0.01	0.01	0.01	0.02	0.01
CD(0.05)	0.36	0.05	0.04	0.03	0.03	0.02	0.03	0.03	0.04	0.03

Table 3. Effect of Mo fertilisation on Nitrate Reductase and Glutamine Synthetase (GS) activities in greengram (*Vigna radiata*)

Treatments	Nitrate reductase activity ($\mu\text{g of NO}_2 \text{ g}^{-1}$ of leaf hour ⁻¹)	Glutamine synthetase ($\text{mol min}^{-1} \text{ mg}^{-1}$ protein)
Mo (0 g Mo ha ⁻¹)	93.8	45.8
M1 (200 g Mo ha ⁻¹)	98.7	49.7
M2 (400 g Mo ha ⁻¹)	100.9	51.2
M3 (600 g Mo ha ⁻¹)	106.7	53.7
M4 (800 g Mo ha ⁻¹)	110.4	55.8
M5 (1000 g Mo ha ⁻¹)	123.7	56.3
SEd	0.24	0.02
CD(0.05)	0.5	0.04

Table 4. Effect of different methods of Mo fertiliser application on growth parameters in greengram (*Vigna radiata*)

Treatments	Shoot		Root	
	Plant Height (cm)	Drymatter Product (DMP) (g plant ⁻¹)	Drymatter Product (DMP) (g plant ⁻¹)	Length (cm)
T1 (control)	38	12.52	0.65	13.5
Seed Treatment				
T2 (2 g Mo kg ⁻¹)	41.8	17.3	1.63	28.2
T3 (4 g Mo kg ⁻¹)	44.5	17.66	1.73	29.3
Foliar spray				
T4 (0.025 %)	53.4	19.86	1.94	32.3
T5 (0.050 %)	57.1	20.28	2.3	34.3
Soil application				
T6 (500 g Mo ha ⁻¹)	48.2	21.04	1.87	27.3
T7 (1000 g Mo ha ⁻¹)	49.7	21.88	1.94	29.3
SEd	1.02	1.3	0.04	0.8
CD (0.05)	2.09	2.7	0.07	1.7



Fig. 1. Overall view of the incubation experiment

3.2 Effect of Molybdenum on Nitrate Reductase and Glutamine Synthetase

The key enzymes involved in N-assimilation namely nitrate reductase and glutamine synthetase. The nitrate reductase activity increased correspondingly with incremental levels of Mo. Molybdenum fertilisation with graded levels increased the nitrate reductase and glutamine synthetase activities (Table 3). Nitrate reductase is a substrate inducible enzyme which closely coincides with the availability of nitrate nitrogen in the soil. Mo fertilisation through soil, seed and foliar spray favoured the availability of Mo in the soil vis-a vis availability of nitrogen. The highest nitrate reductase activity of $123.7 \mu\text{g NO}_2 \text{g}^{-1}$ of leaf hour^{-1} registered in M5 which is 24.2 per cent higher than control. Since Mo fertilised greengram plants are adequately nourished with Mo and nitrogen it is understandable that N-assimilation has improved. Similar trend of response was seen for glutamine synthetase also. The glutamine synthetase activity registered the highest in M5 which is 18.8 per cent higher than control. Further, another N-assimilating enzyme glutamine synthetase facilitates assimilation of ammonium into amino acid glutamine. It is quite evident that N-assimilatory pathway has modified or improved due to Mo fertilisation. Our data are in agreement with the findings of [9]. The correlation study indicated that available Mo positively correlated with nitrate reductase ($r = 0.95$) and glutamine synthetase ($r = 0.85$).

3.3 Effect of Mo Fertilisation on Growth Parameters of Green Gram

Plant parameters such as plant height, shoot dry matter product and root dry matter product characteristics were measured at 40 DAS Molybdenum fertilisation done through seed, foliar sprays and soil had significantly increased the plant height irrespective of doses (Table 4). The highest values were recorded in foliar spray treatments followed by soil and seed treatment. The lowest value was registered in control. In this experiment, Mo applied by foliar application of T4 (0.025 per cent) followed by T5 (0.050 per cent) favourably enhanced shoot and root characteristics. The correlation study indicated that available Mo positively correlated with shoot mass ($r = 0.958$) and root mass ($r = 0.901$). Our data are in agreement with the findings of [10,11] and [12].

3.4 Effect of Molybdenum on Dry Matter Production (DMP) on Shoot and Root

Application of Mo through seed, foliar and soil had increased the individual plant dry matter significantly, the lowest value was registered in control 12.52 g. Among the method of Mo fertilisation, soil application of Mo applied at the rate of 500 or 1000 g ha^{-1} increased the dry matter production nearly 1.5 to 1.7 times in comparison to control. The dry matter Mo produced in soil treatments was comparable to foliar spray treatments. These values were significantly higher than seed treatment. Root measurements

namely length and dry matter were measured at the harvest stage of green gram. The control had the lowest values for all the three parameters measured. The foliar spray of Mo at the rate of 0.025 per cent or 0.050 per cent had increased all the 3 root parameters followed by soil application of Mo and seed treatment. Foliar spray of Mo had increased the dry matter by 3 to 4 times, root length by 2.5 times and volume by 2 times indicating the importance of foliar spray of Mo in root proliferation and dry matter product. This data is in accordance with the findings of [13].

3.5 Effect of Mo Fertilisation on Nodulation and Rhizobium Population

Application of Mo at incremental levels increased the number of nodules linearly and the highest number registered in M5 where Mo was applied at the rate of 1000 g ha⁻¹ (25 nodules plant⁻¹)

(Table 5 and Fig. 2). The number of nodules registered in the best treatment was 2.5 times than control (10 nodules plant⁻¹). The increase in nodule numbers was higher with the higher level of Mo application. The rhizobium population of Mo fertilised treatments at the rate of 1000 g ha⁻¹ had the highest value of 8 CFU ml⁻¹ while control registered the lowest value of 2 CFU ml⁻¹. Molybdenum fertilisation with graded levels increased the Rhizobium population which in turn increases the number nodulation which resulted in the effective N-fixation process. The highest Rhizobium population (8 CFU ml⁻¹) was registered in M5 (1000 g Mo ha⁻¹) and the nodulation number of 25. Our data are in agreement with the findings of [14,15] and [16]. The correlation study indicated that available Mo positively correlated with nodulation number ($r = 0.901$). Our data have unequivocally demonstrated Mo fertilisation assist in nodulation and rhizobium population that promote N-fixation in greengram plants.




Fig. 2. Comparison of nodulation in greengram plant fertilised with (1000 g Mo ha⁻¹) and without Mo application (Control)

Table 5. Effect of Mo fertilisation on nodulation and *Rhizobium* population in greengram (*Vigna radiata*)

Reatments	No of nodules plant ⁻¹	Rhizobium population (CFU/ ml)
Mo (0 g Mo ha ⁻¹)	10	2
M1 (200 g Mo ha ⁻¹)	13	3
M2 (400 g Mo ha ⁻¹)	18	4
M3 (600 g Mo ha ⁻¹)	20	5
M4 (800 g Mo ha ⁻¹)	22	6
M5 (1000 g Mo ha ⁻¹)	25	8
SEd	0.61	0.45
CD (0.05)	1.25	0.92

SEM IMAGE OF GREENGRAM NODULES




Fig. 3a. Without Mo application-Control
100 µm Magnification




Fig. 3b. With Mo application- 1000 g ha⁻¹
100 µm Magnification

3.6 SEM View of Effective Nodules

An experiment was conducted to undertake basic anatomical features of nodules developed in greengram at 45 DAS. The SEM view clearly indicated that the treatments that received Mo fertilisation were found to contain a large number of Bacteroides in comparison to control (Fig. 3a and Fig. 3b). In addition to the number of bacteroids, the peri bacterial membrane shown to be thicker and widespread as a result of Mo fertilisation. The best treatment was soil applied at the rate of 1000 g ha⁻¹ had higher number of active bacteroids involved in N-fixation process. On the other hand, the control had less number of Bacteroides besides many of them appear to be non-functional. On the other hand, the control had less number of Bacteroides besides many of them appear to be non-functional. Our data are in agreement with the findings of [17] and [18].

4. CONCLUSION

Overall, the data clearly indicated that the molybdenum deficiency is closely associated with acidic soil condition. Molybdenum application @ 1000 g ha⁻¹ in the form of sodium molybdate or foliar spray of Mo @ 0.05 per cent at pre-flowering and 15 days later found to enhance the yield of green gram to the tune of 30-35 per cent. Besides, Mo improves the nodulation in green gram which resulted in enhanced available N content of the soil.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Agriculture Statistics at a glance; 2017. Available:<http://www.agricoop.nic.in/recentinitiatives/pocket-book-agricultural-statistics-2017>
2. Economic Survey. A national market for agricultural commodities- some issues and the way forward; 2018. Available:<http://www.indiabudget.nic.in/es2017-18/echapvol1-08.pdf>
3. Chatterjee C, Shukla S, Khurana N. Effect of sulphur deficiency on metabolism of green gram (*Phaseolus radiatus*). Indian Journal of Agricultural Sciences. 1992; 62(7):445-449.
4. Hristozkova M, Geneva M, Stancheva I, Georgiev G. Nitrogen assimilatory enzymes and amino acid content in inoculated foliar fertilized pea plants grown at reduced molybdenum concentration. Journal of Plant Nutrition. 2007;30:1409-1419.
5. Kevresan S, Petrovic N, Popovic M, Kandrac J. Effect of heavy metals on nitrate and protein metabolism in sugar beet. Plant Physiol. 1998;13(12):716-724.

6. Hageman RH, Reed AJ, Part C, ed. Anthony san methods in enzymology. 1980;69.
7. Pateman JA. Regulation of synthesis of glutamate dehydrogenase and glutamine synthetase in microorganism. Biochemical Journal. 1969;15(4):769-775.
8. Ranganathan P, Pramesh CS, Agarwal R. Common pitfalls in statistical analysis: New York, Marcel Dekker. 1990;331-350.
9. Vieira RF, Vieira C, Cardoso EJBN, Mosquim PR. Foliar application of molybdenum in common bean. II. Nitrogenase and nitrate reductase activities in a soil of low fertility. Journal of Plant Nutrition. 1998;21(10):2141—2151.
10. Adkine PM, Mankar DD, Khandait VM, Bhandare VL, Nawlakhe SM. Journal of Soils and Crops. 2011;21(1):116-123.
11. Kovacs B, Anita P, Preszner, Huzsvai L, Levai, Laszlo, Eva B. Effect of molybdenum treatment on molybdenum concentration and nitrate reduction in maize seedlings. Plant Physiology and Biochemistry. 2015;96:38-44.
12. Qin SY, Sun XC, Hu CX, Zhao XH. Uptake, transport and distribution of molybdenum in two oilseed rape (*Brassica napus* L.) cultivars under different nitrate/ ammonium ratios. Journal of Zhejiang University Science. 2017;18(6):512-521.
13. Steiner F, Tiago Z, Alan MZ, Patticia PM, Jardel Z, Andre Z. Foliar application of molybdenum enhanced quality and yield of crisphead lettuce (*Lactuca sativa* L., cv Grand Rapids). Acta Agron. 2018;77(1):73-78.
14. Tripathi SK, Edward JC. Response of rhizobium culture inoculation, zinc and molybdenum application on rhizosphere and phyllosphere microbial population of soyabean (*Glycine max* Merrill) [India]. Current Sciences. 1978;47:503-504.
15. Dey BK, Ghosh A. Effect of molybdenum on nitrogen fixation in rhizosphere soils of rice. Journal of Indian Society of Soil Science. 1986;34:493-497.
16. Bhuiyan MAH, Khanam D, Hossain MF, Ahmed MS. Effect of Rhizobium inoculation on nodulation and yield of chickpea in calcareous soil. Bangladesh Journal of Agriculture Research. 2008; 33(3):549-554.
17. Roth and Stacey. Bacterium release into host cells of nitrogen- fixing soybean nodules: the symbiosome membrane comes from three sources. Eur. J. Cell Biology. 1989;49(1):13-23.
18. Benson HP, Eric Boncompagni, Mary Lou Guerinot. An iron uptake operon required for proper nodule development in the *Bradyrhizobium japonicum*-Soybean Symbiosis. 2005;18(9):950-959.

© 2018 Karpagam and Rajesh; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sciencedomain.org/review-history/27505>