



# Effect of Zinc Levels on Growth and Yield of Cowpea (*Vigna unguiculata* L.) Varieties

Abhishek Chauhan<sup>a++\*</sup>, Biswarup Mehera<sup>b#</sup>  
and Abhishali<sup>a++</sup>

<sup>a</sup> Department of Agronomy, Naini Agricultural Institute, SHUATS, Prayagraj-211007, Uttar Pradesh, India.

<sup>b</sup> Faculty of Agriculture, Naini Agricultural Institute, SHUATS, Prayagraj, Uttar Pradesh, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJPSS/2023/v35i82894

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/98170>

Original Research Article

Received: 27/01/2023

Accepted: 29/03/2023

Published: 07/04/2023

## ABSTRACT

The field experiment was conducted at Crop Research Farm (CRF), Department of Agronomy, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj (U.P.) during *Zaid* season of 2022 on Cowpea crop. The treatments Consists of 3 levels of Zinc [20kg/ha, 0.6% and (10kg/ha + 0.3%)] and 3 different varieties (GOMTI, HIMANI-06, KASHI-KANCHAN) and a Control. The experiment was laid out in Randomized Block Design with 10 treatments replicated thrice. The application of Zinc (10kg/ha + 0.3%) + GOMTI variety recorded higher plant height (83.96 cm), more number of nodules per plant (9.20), maximum plant dry weight (12.55 g/plant), Crop growth rate (7.94 g/m<sup>2</sup>/day), Relative growth rate (0.037 g/g/day) and yield attributes namely Number of Pods per plant (14.44), Number of seeds per pod(18.55), Test weight(91.00 g), Seed yield (1.03 t/ha), Stover yield (5.33 t/ha) and Harvest index (16.12%) was found maximum in treatment 3 that is Zinc (10kg/ha + 0.3%) + GOMTI variety as compared to other treatment combinations. The maximum gross return (INR 112750.00/ha), net return (INR 74582.50/ha) and B:C ratio (1.95) is recorded in treatment 3 Zinc (10kg/ha + 0.3%) + GOMTI variety as well.

<sup>++</sup> M. Sc. Scholar;

<sup>#</sup> Dean;

\*Corresponding author: E-mail: 21msagro119@shiats.edu.in, abhishek.ak6547@gmail.com;

**Keywords:** Cowpea; zinc; varieties; growth parameters; yield attributes; yield and economics.

## 1. INTRODUCTION

“Cowpea (*Vigna unguiculata* L.) is a native to Central Africa and belongs to the family Leguminaceae with subfamily Papilionaceae” [1]. “The common names are Black-eye-pea, Southern pea, China pea, Marble pea. Cowpea is annual herbaceous plant known for its drought hardy nature with large tap root system and alternate trifoliate leaves with ovate leaflets. It also has the useful ability to fix atmospheric nitrogen through its root nodules. In India, during 2022-23 the production of pulses was 27.69 metric ton. Cowpea is grown over an area about 58000ha with an average production of 4.8 lakh tons and average productivity is 8.44 t/ha. During 2022-23 the total coverage under cowpea in Uttar Pradesh 16900 ha with a production will be around 113200 tones and the productivity 6.70t/ha” GOI [2]. It consists of high-quality protein for human consumption and it is rich in protein for livestock fodder.

“Zinc is an essential micronutrient for plants, due its role as a functional, structural or regulatory cofactor in a large number of enzymes. Zinc is necessary to activate many enzymes, enzymes that are activated by zinc are Tryptophan synthetase superoxide dismutase and dehydrogenases. The plant with lack of zinc is poor in amount of protein. Foliar spraying of micronutrients for the growth of Cowpea and its quality in industry views is necessary for growth and quality of Cowpea”. Sirisha et al [1] showed that “zinc fertilizer application causes root and shoot growth during the growing season and therefore, lead to increased seed yield. Spraying the leaves with the nutrient elements is one of the methods of plant supply”. “Application of Higher levels of zinc to cowpea leads to increase in the higher auxin activity in plant which promoted growth attributes and higher biomass accumulation in plants. This resulted in higher plant dry matter. The results were in resonance with” Kumar and Bohra [3]. The involvement of zinc in different physiological processes like enzyme activation, electron transport, chlorophyll formation, stomatal regulation, etc. With the increase in levels of zinc the plant height gradually increased, which might be attributable to greater photosynthetic activity and chlorophyll synthesis due to zinc fertilization resulting into better vegetative growth. Similar results were reported by Kumar et al. [4].

## 2. MATERIALS AND METHODS

The field experiment was conducted at Crop Research Farm (CRF), Department of Agronomy, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj (U.P.) during *Zaid* season of 2022 on Cowpea crop. The soil of the experimental field constituting a part of central Gangetic alluvium is neutral and deep. The soil of experimental plot was sandy loam in texture, nearly neutral in soil reaction (pH 7.4), low in organic carbon (0.49%), available N (219kg/ha), available P (21.3kg/ha) and available K (235.8kg/ha). The treatments Consists of 3 levels of Zinc [20kg/ha, 0.6% and (10kg/ha + 0.3%)] and 3 different varieties (GOMTI, HIMANI-06, KASHI-KANCHAN) and a Control. The experiment was laid out in Randomized Block Design with 10 treatments replicated thrice. T<sub>1</sub>: GOMTI (SPL.) + 20kg/ha ZnSo<sub>4</sub>, T<sub>2</sub>: GOMTI (SPL.) + 0.6% ZnSo<sub>4</sub>, T<sub>3</sub>: GOMTI (SPL.) + [10kg/ha + 0.3%] ZnSo<sub>4</sub>, T<sub>4</sub>: HIMANI-06 + 20kg/ha ZnSo<sub>4</sub>, T<sub>5</sub>: HIMANI-06 + 0.6% ZnSo<sub>4</sub>, T<sub>6</sub>: HIMANI-06 + [10kg/ha + 0.3%] ZnSo<sub>4</sub>, T<sub>7</sub>: KASHI-KANCHAN + 20kg/ha ZnSo<sub>4</sub>, T<sub>8</sub>: KASHI-KANCHAN + 0.6% ZnSo<sub>4</sub>, T<sub>9</sub>: KASHI-KANCHAN + [10kg/ha + 0.3%] ZnSo<sub>4</sub>, T<sub>10</sub>: Control [N:P:K (25:50:25) kg/ha].

**Note:** The collected data was subjected to statistical analysis by analysis of variance method [5].

## 3. RESULTS AND DISCUSSION

### 3.1 Growth Parameters

**Plant height (cm):** The data revealed that significant and higher plant height (83.96 cm) was recorded in treatment 3 that is GOMTI + [10kg/ha + 0.3%] ZnSo<sub>4</sub>, which was significantly superior over rest of the treatments (Table 1). Significant increase of plant height might be due to the increase in involvement of zinc in different physiological processes like enzyme activation, electron transport, chlorophyll formation, stomatal regulation, etc. With the increase in levels of zinc the plant height gradually increased, which might be attributable to greater photosynthetic activity and chlorophyll synthesis due to zinc fertilization resulting into better vegetative growth. Similar results were reported by Kumar et al. [4].

**Number of nodules/plant:** The data revealed that significant and higher number of nodules/plant (9.20) was recorded in Treatment 3 that is GOMTI + [10kg/ha + 0.3%] ZnSo<sub>4</sub>. However Treatment 2 that is GOMTI + 0.6% ZnSo<sub>4</sub> were found to be statistically at par with treatment 3 GOMTI + [10kg/ha + 0.3%] ZnSo<sub>4</sub> (Table 1). Application of zinc increased significantly number of nodules per plant, active nodule per plant and active nodule weight per plant. Significant increase of nodules/plant might be due the maximum number of nodules per plant and active nodule were recorded in 15 kg Zn / ha while minimum in recorded in control. Similar trend was also observed by Singh and Shukla [6] and Wani and Khan [7].

**Dry weight(g):** The data revealed that significant and higher plant dry weight (12.55 g/plant) was recorded in Treatment 3 that is GOMTI + [10kg/ha + 0.3%] ZnSo<sub>4</sub> which was significantly superior over rest of all the treatments (Table 1). Significant increase of plant dry weight might be due to Application of Higher levels of zinc to cowpea leads to increase in the higher auxin activity in plant which promoted growth attributes and higher biomass accumulation in plants. This resulted in higher plant dry matter. The results were in resonance with Kumar and Bohra [3].

**Crop growth rate(g/ m<sup>2</sup>/ days):** The data revealed that significant and maximum crop growth rate (8.23 g/ m<sup>2</sup>/ days) was recorded in treatment 4 that is [HIMANI-06 + 20kg/ha ZnSo<sub>4</sub>], however treatment 1 GOMTI + 20kg/ha ZnSo<sub>4</sub>, treatment 2 GOMTI + 0.6% ZnSo<sub>4</sub>, treatment 3 GOMTI + [10kg/ha + 0.3%] and treatment 5 HIMANI-06 + 0.6% ZnSo<sub>4</sub> were found to be statistically at par with treatment 4 [HIMANI-06 + 20kg/ha ZnSo<sub>4</sub>] (Table 1). The significant increase in crop growth rate observed with the application of [10kg/ha + 0.3%] ZnSo<sub>4</sub> might be due the dry matter yield increase in the Zn treatments over the control suggests that Zn was one of the limiting nutrients in the soils. This indicates that, at this level, the soil Zn was further improved with better Zn nutrition leading to high dry matter production which in turn increases CGR. The similar results were reported by Marnagar and Dawson (2017). [8].

**Relative growth rate (g/g/day):** The data revealed that significant and maximum relative growth rate (0.054 g/g/day) was recorded in treatment 9 that is KASHI-KANCHAN + [10kg/ha + 0.3%] ZnSo<sub>4</sub>, however treatment 4 is found to be statistically at par with treatment 9 KASHI-

KANCHAN + [10kg/ha + 0.3%] ZnSo<sub>4</sub> (Table 1). The significant increase in relative growth rate observed with the application of [10kg/ha + 0.3%] ZnSo<sub>4</sub> might be due the application of micronutrients on growth of cowpea, in terms of dry matter and crop growth rate can be interpreted in terms of the metabolic function of micronutrients in the plant which ultimately increases the relative growth rate. The similar results were reported by Singh et al. [9].

### 3.2 Yield Parameters

**No. of Pods/plant:** The data revealed that significant and higher No. of Pods/plant (14.44) was recorded in Treatment 3 that is GOMTI + [10kg/ha + 0.3%] ZnSo<sub>4</sub>. However treatment 1 GOMTI + 20kg/ha ZnSo<sub>4</sub> is statistically at Par with the treatment 3 GOMTI + 20kg/ha ZnSo<sub>4</sub> (Table 2). Number of pods/plant was significantly increased with application of [10kg/ha + 0.3%] ZnSo<sub>4</sub>. It could possibly be explained by the fact that zinc application increased the realization of flower into pods. The similar results were reported by Kumar et al. [10].

**No. of Seeds/Pod:** The data revealed that significant and higher no. of seeds /pods (18.55) was obtained in treatment 3 that is GOMTI + [10kg/ha + 0.3%] ZnSo<sub>4</sub>. However treatment 2 GOMTI + 0.6% ZnSo<sub>4</sub> is statistically at Par with the treatment 3 GOMTI + [10kg/ha + 0.3%] ZnSo<sub>4</sub> (Table 2). Number of seeds/pod was significantly increased with application of [10kg/ha + 0.3%] ZnSo<sub>4</sub>. Zinc plays a very important role in the metabolism of the plant process by influencing the activity of growth enzymes as well as it is involved in carbohydrate metabolism, maintenance of the integrity of cellular membranes, protein synthesis, and regulation of auxin synthesis and pollen formation. The similar results were reported by Khan et al. [11].

**Test weight (g):** The significant and maximum Test weight (91.00 g) was obtained in treatment 3 that is GOMTI + [10kg/ha + 0.3%] ZnSo<sub>4</sub>, however treatment 2 GOMTI + 0.6% ZnSo<sub>4</sub> is statistically at par with the treatment 3 GOMTI + [10kg/ha + 0.3%] ZnSo<sub>4</sub> (Table 2). Test weight was significantly increased with application of [10kg/ha + 0.3%] ZnSo<sub>4</sub>. Similarly higher yield attributes and yield were noticed with the combined foliar spray of micro nutrients with zinc attributed to optimum availability of nutrients for luxurious crop growth and efficient partitioning of assimilates from source to sink. The similar results were reported by Masih et al. [12].

**Table 1. Effects of zinc levels on growth attributes of cowpea varieties**

Treatment	60 DAS			45-60 DAS	
	Plant height (cm)	No. of nodules/plant	Plant dry weight (g)	Crop Growth Rate (g/m <sup>2</sup> /day)	Relative Growth Rate (g/g/day)
1 GOMTI + 20kg/ha ZnSo <sub>4</sub>	80.98	8.60	11.21	7.54	0.040
2 GOMTI + 0.6% ZnSo <sub>4</sub>	81.96	8.93	11.85	7.65	0.038
3 GOMTI + [10kg/ha + 0.3%] ZnSo <sub>4</sub>	83.96	9.20	12.55	7.94	0.037
4 HIMANI-06 + 20kg/ha ZnSo <sub>4</sub>	77.68	7.40	10.20	8.23	0.053
5 HIMANI-06 + 0.6% ZnSo <sub>4</sub>	78.80	7.80	10.38	7.70	0.046
6 HIMANI-06 + [10kg/ha + 0.3%] ZnSo <sub>4</sub>	79.48	8.20	11.19	7.28	0.039
7 KASHI-KANCHAN + 20kg/ha ZnSo <sub>4</sub>	72.97	6.40	8.71	6.25	0.044
8 KASHI-KANCHAN + 0.6% ZnSo <sub>4</sub>	74.95	6.67	9.15	5.92	0.038
9 KASHI-KANCHAN + [10kg/ha + 0.3%] ZnSo <sub>4</sub>	76.57	7.00	10.27	6.33	0.036
10 Control [N:P:K (25:50:25) kg/ha]	68.45	5.87	8.37	6.82	0.054
<b>F test</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>
<b>SEm(±)</b>	<b>0.38</b>	<b>0.10</b>	<b>0.17</b>	<b>0.31</b>	<b>0.00</b>
<b>CD (5%)</b>	<b>1.14</b>	<b>0.29</b>	<b>0.51</b>	<b>0.92</b>	<b>0.007</b>

**Table 2. Effects of zinc levels on yield attributes of cowpea varieties**

S. No.	TREATMENTS	No. of pods/plant	No. of seeds/pod	Test weight (g)	Seed yield (t/ha)	Stover yield (t/ha)	Harvest index (%)
1	GOMTI + 20kg/ha ZnSo <sub>4</sub>	13.44	17.00	86.33	0.88	5.15	14.54
2	GOMTI + 0.6% ZnSo <sub>4</sub>	13.22	17.89	88.00	0.98	5.20	15.80
3	GOMTI + [10kg/ha + 0.3%] ZnSo <sub>4</sub>	14.44	18.55	91.00	1.03	5.33	16.12
4	HIMANI-06 + 20kg/ha ZnSo <sub>4</sub>	11.78	14.67	81.00	0.84	4.67	15.31
5	HIMANI-06 + 0.6% ZnSo <sub>4</sub>	12.22	15.33	84.33	0.87	4.78	15.43
6	HIMANI-06 + [10kg/ha + 0.3%] ZnSo <sub>4</sub>	13.00	16.33	86.67	0.87	4.98	14.90
7	KASHI-KANCHAN + 20kg/ha ZnSo <sub>4</sub>	8.55	13.11	74.00	0.79	4.63	14.65
8	KASHI-KANCHAN + 0.6% ZnSo <sub>4</sub>	10.00	13.22	76.00	0.77	4.64	14.17
9	KASHI-KANCHAN + [10kg/ha + 0.3%] ZnSo <sub>4</sub>	10.89	14.44	79.00	0.84	4.65	15.26
10	Control [N:P:K (25:50:25) kg/ha]	7.89	10.45	72.00	0.74	4.44	14.24
<b>F test</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>
<b>SEm (±)</b>	<b>0.41</b>	<b>0.28</b>	<b>1.21</b>	<b>0.02</b>	<b>0.03</b>	<b>0.34</b>	<b>0.34</b>
<b>CD (5%)</b>	<b>1.22</b>	<b>0.83</b>	<b>3.61</b>	<b>0.06</b>	<b>0.08</b>	<b>1.00</b>	<b>1.00</b>

**Table 3. Effects of zinc levels on economics of cowpea varieties**

S. No.	Treatments	Cost of cultivation (₹/ha)	Gross Return (₹/ha)	Net return (₹/ha)	Benefit cost ratio (B:C)
1	GOMTI + 20kg/ha ZnSo <sub>4</sub>	40632.50	96470.00	55837.50	1.37
2	GOMTI + 0.6% ZnSo <sub>4</sub>	35702.50	106286.30	70583.80	1.94
3	GOMTI + [10kg/ha + 0.3%] ZnSo <sub>4</sub>	38167.50	112750.00	74582.50	1.95
4	HIMANI-06 + 20kg/ha ZnSo <sub>4</sub>	44632.50	92840.00	48208.00	1.08
5	HIMANI-06 + 0.6% ZnSo <sub>4</sub>	39702.50	95992.60	56290.10	1.41
6	HIMANI-06 + [10kg/ha + 0.3%] ZnSo <sub>4</sub>	42167.50	95956.30	53788.80	1.27
7	KASHI-KANCHAN + 20kg/ha ZnSo <sub>4</sub>	42632.50	87340.00	44707.50	1.05
8	KASHI-KANCHAN + 0.6% ZnSo <sub>4</sub>	37702.50	84260.00	46558.00	1.23
9	KASHI-KANCHAN + [10kg/ha + 0.3%] ZnSo <sub>4</sub>	40167.50	92290.00	52122.50	1.29
10	Control [N:P:K (25:50:25) kg/ha]	37528.50	81143.70	43615.20	1.16

**Seed yield (t/ha):** The significant and maximum Seed yield (1.03 t/ha) was obtained in treatment 3 that is GOMTI + [10kg/ha + 0.3%] ZnSo<sub>4</sub>, however treatment 2 GOMTI + 0.6% ZnSo<sub>4</sub> is statistically at par with the treatment 3 GOMTI + [10kg/ha + 0.3%] ZnSo<sub>4</sub> (Table 2). Seed yield was significantly increased with application of [10kg/ha + 0.3%] ZnSo<sub>4</sub>. Zn influenced the synthesis of IAA in plants which indirectly enhanced the growth and development and uptake of nutrient in plants. Application of Zn also influenced the grain yield of cowpea. The similar results were reported by Chavan et al. [13]0.

**Stover yield (t/ha):** The significant and maximum Stover yield (5.33 t/ha) was obtained in treatment 3 that is GOMTI + [10kg/ha + 0.3%] ZnSo<sub>4</sub>, which was significantly superior over all the treatments. stover yield was significantly increased with application of [10kg/ha + 0.3%] ZnSo<sub>4</sub> (Table 2). It might be due to its direct influence on auxin production which in turn enhanced the elongation processes of plant development. The similar results were reported by Masih et al. [12].

**Harvest index (%):** The significant and maximum Harvest index (16.12 %) was obtained in treatment 3 that is GOMTI + [10kg/ha + 0.3%] ZnSo<sub>4</sub>, however treatment 2 GOMTI + 0.6% ZnSo<sub>4</sub>, treatment 4 HIMANI-06 + 20kg/ha ZnSo<sub>4</sub>, treatment 5 HIMANI-06 + 0.6% ZnSo<sub>4</sub> and treatment 9 KASHI-KANCHAN + [10kg/ha + 0.3%] ZnSo<sub>4</sub> is statistically at par with the treatment 3 GOMTI + [10kg/ha + 0.3%] ZnSo<sub>4</sub> (Table 2). Harvest index was significantly increased with application of [10kg/ha + 0.3%] ZnSo<sub>4</sub>. It is found due to the increase in grain yield and straw yield the harvest index increases. The similar results were reported by Khan et al. [11].

**Economics(₹/ha):** Maximum gross return (112750.00 ₹/ha), highest net return (74582.50 ₹/ha) and highest B:C (1.95) was also recorded in treatment 3 with the application of GOMTI + [10kg/ha + 0.3%] ZnSo<sub>4</sub> as compared to control (Table 3). Significantly increase net return and benefit cost ratio due to inorganic sources of nutrient under the present study it can be concluded that, the reproductive efficiency of black gram is significantly influenced by NPK with zinc and iron treatments. The similar results were reported by Kumawat et al. [14].

#### 4. CONCLUSION

In view of the obtained results, it could be concluded that among the studied treatments,

treatment 3 that is GOMTI + [10kg/ha + 0.3%] ZnSo<sub>4</sub> was found to be more desirable that give higher growth parameters, yield attributes, seed yield, stover yield and harvest index. It also recorded maximum gross return, net return and B:C ratio.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Sirisha SV, Umesha C, Singh SN. Influence of phosphorus levels on growth, yield and economics of cowpea (*Vigna unguiculata* L.) Varieties. The pharmajournal; 2022.
2. GOI. Agricultural statistics at a glance; 2022. Agricultural Statistics Division, Directorate of Economics and Statistics, Department of Agriculture, Cooperation Ministry of Agriculture, Government of India. New Delhi. [hips://eands. Available from: daenet.nic.in](https://eands. Available from: daenet.nic.in).
3. Kumar R, Bohra JS. Effect of NPKS and Zn application on growth, yield, economics and quality of baby corn. Arch Agron Soil Sci. 2014;60(9):1193-206.
4. Kumar R, Rathore DK, Singh M, Kumar P, Khippal A. Effect of phosphorous and zinc nutrition on growth and yield of fodder cowpea. Legume Res. 2016;39(2):262-7.
5. Gomez KA, Gomez AA. Three or more factor experiment. (In:) Statistical Procedure for Agricultural. research 2nd ed. 1976;1976:139-41.
6. Singh K, Shukla UC. Evaluation of critical level of soil zinc for predicting response of maize to zinc fertilization in arid soil. Int J Trop Agric. 1986;3:288-92.
7. Wani MR, Khan S. Estimate of genetic variability in mutated populations and scope of selection for yield attributes in *Vigna radiate* (L) Wilezek. Egypt J Biol. 2006;8:1-6.
8. Marnagar E, Dawson J. Effect of biofertilizers, levels of nitrogen and zinc on growth and yield of hybrid maize (*Zea mays* L.). Int J Curr Microbiol Appl Sci. 2017;6(9):3614-22.
9. Singh LB, Yadav R, Abraham T. Studies on the effect of zinc levels and methods of boron application on growth, yield and protein content of wheat (*Triticum aestivum* L.). Bull Environ Pharmacol Life Sci. 2015;4(2):108-13.

10. Kumar R, Singh YV, Singh S, Latore AM, Mishra PK, Supriya. Effect of phosphorus and sulphur nutrition on yield attributes, yield of mungbean (*Vigna radiata* L. Wilczek). J Chem Pharm Res. 2012;4(5):2571-3.
11. Khan A, Hayat Z, Ali Khan A, Ahmad J, Abbas MW, Nawaz H et al. Effect of foliar application of zinc and boron on growth and yield components of wheat. Agric Res Technol Open Access J. 2019; 21(1):3-6.
12. Masih A, Dawson J, Singh RE. Effect of levels of phosphorus and zinc on growth and yield of Greengram (*Vigna radiata* L.). Int J Curr Microbiol Appl Sci. 2020; 9(10):3106-12.
13. Chavan AS. Khaf,M.R.; raj,A.D.; Parmar, R.M. Eff Potassium Znic Yield Protien Content Uptake Micronutrients Cow Pea [ (*Vigna unguiculata* L.) Walp.] Jornal of Agriculture Science Digest. 2012;32(2):175.
14. Kumawat PK, Tiwari RC, Golada SL, Godara AS, Garhwal RS, Choudhary R. Effect of phosphorus sources, levels and biofertilizers on yield attributes, yield and economics of black gram (*Phaseolous mungo* L.). Legume Res Int J. 2013;36(1):70-3.

© 2023 Chauhan et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/98170>