



Economic Factors and Different Growth Phases of Sweet Corn (*Zea mays* L. Var. *Saccharata*) in the South Gujarat Area, India as affected by Intra-Row Spacing and Potassium Levels

Manisha Sharma^{a++}, Manoj Kumar^{a#} and B. M. Dabhi^{a†}

^a Department of Agronomy, College of Agriculture, Junagadh Agricultural University, Junagadh, Gujarat- 362001, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i153074

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/99784>

Original Research Article

Received: 09/03/2023
Accepted: 11/05/2023
Published: 09/06/2023

ABSTRACT

To determine how intra-row spacing and potassium levels affected sweet corn (*Zea mays* L. var. *Saccharata*), a field experiment was conducted in Junagadh (Gujarat) during the Rabi season of 2016–17. Four levels of intra-row spacing (5, 10, 15 and 20 cm) and four potassium levels (0, 20, 40, and 60 kg K₂O/ha) were combined into sixteen different treatment combinations. Three replications of a factorial randomized block design were used to set up the experiment. Based on the data of growth attributes significant and maximum recorded dry matter accumulation (102.46

⁺⁺ PG Scholar;

[#] Assistant Development Officer;

[†] Retd. Research Scientist;

*Corresponding author: E-mail: manojagric@gmail.com;

and 144.54 g plant⁻¹) at 60 DAS and harvest, absolute growth rate (3.07 and 1.41 g day⁻¹) at 45-60 DAS and 60 DAS-harvest, crop growth rate (0.00275 and 0.00126 g m⁻²day⁻¹) at 45-60 DAS and 60 DAS-harvest under the treatment intra-row spacing (20 cm), respectively. According to data on growth attributes, the treatment (K₄) 60 K₂O + 120 N₂O + 60 P₂O₅ kg/ha resulted in significant and maximum dry matter accumulation (96.93 and 142.00 g plant⁻¹) at 60 DAS and harvest, absolute growth rate (3.03 and 1.39 g day⁻¹) at 45-60 DAS and 60 DAS-harvest, and crop growth rate (0.00272 and 0.00124 g m⁻²day⁻¹) respectively. Economic analysis showed that higher net returns and B: C ratio from sweet corn (Sweet-16) can be secured by sowing the crop at 20 cm intra-row spacing + application of 60 kg K₂O/ha.

Keywords: Economics; Spacing; Gujarat; Growth; Potassium.

1. INTRODUCTION

Due to its classification as a C₄ type crop, maize (*Zea mays* L.) is often grown effectively throughout the year. Among the numerous forms of maize, sweet corn is particularly famous for the use of its green cobs all throughout the world. Popular vegetable sweet corn is in second in farm value and fourth in terms of commercial crops. The sweet corn is possible to boost agricultural revenue because of the rise in demand. The major consideration is to maintain stand density in order to increase cob yields. The form and size of plant leaf area are determined by its spatial layout, which in turn affects how well it can absorb solar energy and how quickly its roots can develop and function. Only when plant population permits each plant to reach its full natural potential and maximum production can be anticipated. In order to get the best population density, inter- and intra-row spacing must be modified in connection to other agronomic parameters [1].

For maize, potassium (K) is a macronutrient because the plant absorbs a lot of it during the growth season. K serves as an activator for several enzymes and metabolic processes, including those involved in photosynthesis, protein synthesis, and starch production in grains, even though the plant does not use it as a building block for organic molecules. Potassium has a function in the flow of water, minerals, and carbohydrates inside the plant. It controls how stomata close and open, which affects how much water and gas are exchanged. Moreover, K is crucial for cell wall strength and cellulose formation. Strong cell walls that increase disease resistance and the capacity of the crop to keep firm, robust stalks are linked to high K fertility. For regulating disease incidence and stalk strength as corn output levels rise, it's critical to maintain a balance between nitrogen (N) and potassium (K) levels. The plant's ability to take N

from the soil is constrained when K is a limiting factor, which has an effect on stalk strength, disease resistance, and grain output [2].

2. MATERIALS AND METHODS

The outcomes of a field experiment titled "Study of intra-row spacing and potassium levels on growth, yield, and quality of sweet corn (*Zea mays* L. var. *Saccharata*) under South Saurashtra conditions" carried out at the Farming System Research Centre, Department of Agronomy, College of Agriculture, Junagadh Agricultural University, Junagadh, Gujarat during *Rabi* season of 2016-17.

2.1 Crop Husbandry

Dry matter accumulation: Five plants were randomly selected from border lines of each experimental plot at 30, 45, 60 DAS and harvest. After chopping, plant samples were placed separately in perforated paper bags and oven dried at 65°C till a constant weight is obtained. Later, these were weighted and dry matter was expressed as g plant⁻¹. **Relative growth rate (RGR):** The values for relative growth rate were calculated for the stage between 30 DAS and 45 DAS, 45 DAS and 60 DAS and then between 60 DAS and at harvest with the help of following formula. $RGR = \frac{\log_e w_2 - \log_e w_1}{(t_2 - t_1)}$ g g⁻¹ day⁻¹. Where, $\log_e w_1$ = Loge of dry weight of plant at time interval t_1 . $\log_e w_2$ = Loge of dry weight of plant at time interval t_2 . **Crop growth rate (CGR):** The values for crop growth rate were calculated for the stage between 30 DAS and 45 DAS, 45 DAS and 60 DAS and then between 60 DAS and harvest with the help of following formula: $CGR = \frac{1}{p} \frac{(w_2 - w_1)}{(t_2 - t_1)}$ g m⁻² day⁻¹. Where, w_1 = weight of dry matter of plant at time t_1 . w_2 = weight of dry matter of plant at time t_2 . p = ground area (m²).

Absolute growth rate: The values for absolute growth rate were calculated for the stage

between 30 DAS and 45 DAS, 45 DAS and 60 DAS and then between 60 DAS and harvest with the help of following formula $AGR = (w_2 - w_1) / (t_2 - t_1)$ g day⁻¹. Where, w_1 = dry weight of plant at time t_1 . w_2 = dry weight of plant at time t_2 .

2.2 Crop Economics

Cost of cultivation: The expenses incurred for all the routine operations from preparatory tillage to harvesting including threshing, cleaning as well as the cost of inputs viz. seeds, fertilizers, pesticides, irrigation etc. applied to each treatment were calculated on the basis of prevailing local charges and then cost of cultivation was calculated. **Gross returns:** The gross realization in terms of rupees per hectare was worked out separately for each treatment considering the green cob and green fodder yields from each treatment and local market prices. **Net returns:** The total cost of cultivation was deducted from the gross realization to work out the net income for each treatment combinations and was recorded accordingly. **Benefit: cost ratio (B:C):** The Benefit: Cost ratio (B:C) ratio was calculated with the help of following formula. $B:C = \text{Gross returns (₹/ha)} / \text{Total cost of cultivation (₹/ha)}$.

2.3 Crop Statistical Analysis

By using the appropriate analysis of variance as suggested by Gomez and Gomez [3] the data was subjected to statistical analysis. The critical difference (CD) values were generated for each instance when the F values were determined to be significant at the 5% level of probability in order to compare the treatment means.

3. RESULTS AND DISCUSSION

3.1 Growth Attributes

Different intra-row spacing levels had no discernible impact on the accumulation of dry matter at 30 and 45 DAS, according to an analysis of the data (Table 1). Nevertheless, intra-row spacing of 20 cm (S_4) recorded considerably greater dry matter accumulation at 60 DAS and harvest (102.46 g and 144.54 g, respectively), which was deemed statistically comparable to treatment S_3 (intra-row spacing of 15 cm).

Due to the consideration of various intra-row spacing values, absolute growth rate between 30-45 DAS was not significantly impacted.

Nevertheless, during 45-60 DAS, intra-row spacing of 20 cm (S_4) recorded considerably the higher absolute growth rate (3.07 g/day) which was remained statistically at par with treatment S_3 (15 cm). Nevertheless, between 60 DAS-harvest and treatment S_4 (20 cm), which was discovered statistically at par with treatment S_3 , absolute growth rate continued to drop and was recognized to be greater (1.41 g/day) (15 cm). Crop growth is a well-established consequence of environmental interaction. In the current study, it was revealed that intra-row spacing had a significant impact on crop production for each plant, indirectly dictating the amount of rivalry among plants for different growth inputs as well as the availability of different growth nutrients to individual plants in the community.

Also, between 45 and 60 DAS, the treatment with a 20 cm intra-row spacing (S_4) recorded statistically equivalent results to treatment S_3 but with a much greater crop growth rate (0.00275 g m⁻² day⁻¹) (15 cm). Nevertheless, between 60 DAS-harvest and treatment S_4 (20 cm), which was found statistically at par with treatment S_3 , the crop growth rate started to drop and was recorded higher (0.00126 g m⁻² day⁻¹) (15 cm).

It is common knowledge that N, P, and K are key nutrients for crop growth and development. The greatest levels of N, P, and K in the crop's plant portion at the recommended intra-row spacing (S_4) of 20 cm may have aided in the promotion of plant development through active cell division and elongation. Under 20 cm of spacing, there appears to be a larger accumulation of photosynthates and ultimately a higher accumulation of dry matter by individual plants as a result of the enhanced nutritional state. In comparison to limited intra-spacing, higher values of growth characteristics were seen with broader intra-row spacing. The increase seen with greater intra-row separation may be related to a reduction in plant competition for nutrients and light under equidistant spatial arrangement [4]. The result is in close accordance with findings of Bozorgi et al. [5], Gozubenli and Konuskan [6], Hamni and Dadari [7] and Paradkar [8].

Application of treatment K_4 (60 K₂O + 120 N₂O + 60 P₂O₅ kg/ha) resulted in considerably larger dry matter accumulation at 60 DAS and harvest than treatment K_3 (40 K₂O + 120 N₂O + 60 P₂O₅ kg/ha) (96.93 g and 142.00 g respectively). Significantly reduced levels of dry matter accumulation (73.11 g and 108.14 g) were observed at 60 DAS and harvest, respectively,

Table 1. Effect of intra-row spacing and potassium levels on dry matter accumulation on relative growth rate, absolute growth rate and crop growth rate of sweet corn

Treatments	Dry matter accumulation				Crop growth rate			Relative growth rate			Absolute growth rate		
	at (g plant ⁻¹)			Harvest	at (g m ⁻² day ⁻¹)			at (g g ⁻¹ day ⁻¹)			at (g/day)		
	30 DAS	45 DAS	60 DAS			30-45 DAS	45-60 DAS	60 DAS- Harvest	30-45 DAS	45-60 DAS	60 DAS- Harvest	30-45 DAS	45-60 DAS
Intra-row spacing (cm)													
S ₁ : 5 Intra-row + 45 Inter-rows	24.37	51.02	70.72	106.33	0.00163	0.00231	0.00100	0.0535	0.0497	0.0135	1.81	2.58	1.11
S ₂ : 10 Intra-row + 45 Inter-rows	25.84	52.75	82.90	122.73	0.00165	0.00237	0.00106	0.0561	0.0505	0.0139	1.84	2.64	1.18
S ₃ : 15 Intra-row + 45 Inter-rows	25.85	55.75	92.96	133.62	0.00166	0.00259	0.00124	0.0562	0.0525	0.0140	1.85	2.89	1.38
S ₄ : 20 Intra-row + 45 Inter-rows	27.48	57.94	102.46	144.54	0.00168	0.00275	0.00126	0.0609	0.0571	0.0141	1.88	3.07	1.41
S.Em.±	0.75	2.14	3.42	4.27	0.00005	0.00010	0.00004	0.0020	0.0023	0.0003	0.06	0.11	0.05
C.D. at 5%	NS	NS	9.87	12.32	NS	0.00029	0.00013	NS	NS	NS	NS	0.32	0.14
Potassium levels (kg/ha)													
K ₁ : 0 K ₂ O + 120 N ₂ O + 60 P ₂ O ₅ kg/ha	24.95	51.01	73.11	108.14	0.00163	0.00229	0.00102	0.0535	0.0478	0.0136	1.82	2.55	1.14
K ₂ : 20 K ₂ O + 120 N ₂ O + 60 P ₂ O ₅ kg/ha	25.31	53.48	85.10	122.34	0.00165	0.00241	0.00107	0.0561	0.0517	0.0139	1.83	2.69	1.20
K ₃ : 40 K ₂ O + 120 N ₂ O + 60 P ₂ O ₅ kg/ha	26.55	56.25	93.89	134.74	0.00166	0.00260	0.00122	0.0576	0.0534	0.0140	1.85	2.90	1.36
K ₄ : 60 K ₂ O + 120 N ₂ O + 60 P ₂ O ₅ kg/ha	26.72	56.73	96.93	142.00	0.00168	0.00272	0.00124	0.0595	0.0569	0.0141	1.87	3.03	1.39
S.Em.±	0.75	2.14	3.42	4.27	0.00005	0.00010	0.00004	0.0020	0.0023	0.0003	0.06	0.11	0.05
C.D. at 5%	NS	NS	9.87	12.32	NS	0.00029	0.00013	NS	NS	NS	NS	0.32	0.14
Interaction (S × K)													
S.Em.±	1.51	4.27	6.84	8.53	0.00010	0.00020	0.00009	0.0039	0.0045	0.0006	0.12	0.22	0.10
C.D. at 5%	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
C.V. %	10.08	13.62	13.57	11.65	10.76	13.91	13.36	11.99	14.99	7.80	10.85	13.91	13.36

with treatment K₁ (0 K₂O + 120 N₂O + 60 P₂O₅ kg/ha). The application of 60 K₂O + 120 N₂O + 60 P₂O₅ (K₄) between 45 and 60 DAS also recorded a considerably greater absolute growth rate (3.03 g/day), which was statistically comparable to treatment K₃ (40 K₂O + 120 N₂O + 60 P₂O₅ kg/ha). However, between 60 DAS-harvest and treatment K₄ (60 K₂O + 120 N₂O + 60 P₂O₅ kg/ha), which was found statistically at par with treatment K₃ (40 K₂O + 120 N₂O + 60 P₂O₅ kg/ha), the absolute growth rate continued to drop and was recorded higher (1.39 g/day). When treatment K₃ (40 K₂O + 120 N₂O + 60 P₂O₅ kg/ha), which was statistically comparable to treatment K₄, applied 60 K₂O + 120 N₂O + 60 P₂O₅ kg/ha, it considerably increased crop growth rate (0.00272 g m⁻² day⁻¹) between 45 and 60 DAS. Nevertheless, between 60 DAS-harvest and treatment K₄ (60 K₂O + 120 N₂O + 60 P₂O₅ kg/ha), which was found statistically at par with treatment K₃ (40 K₂O + 120 N₂O + 60 P₂O₅ kg/ha), the crop growth rate started to fall and was recorded higher (0.00124 g m⁻² day⁻¹).

Since potassium has a favourable influence on growth and enhances cell division and cell expansion, it has a beneficial effect on growth. Potassium's impact on the production of phytohormones plays a significant part in meristematic growth. Cytokinin, one of several

plant hormones, is crucial for the development of tillers and buds. High spikelet fertility is a result of improved pollen germination in the florets due to potassium feeding. Such an increase may also be attributable to the root systems of plants receiving enough potassium from the soil, which increases photosynthesis and the production of metabolites and enzymes in plants [9]. According to Kumar et al. [10] potassium boosts the plant's potential resistance to illnesses and insect pests. [11,12] both indicated that K had positive impacts on growth.

3.2 Economics

According to a review of the data (Table 2), treatment S₄ (intra-row spacing of 20 cm) considerably produced greater gross returns, which were ₹ 92734/ha, whereas treatment S₁ (intra-row spacing of ₹ 65575/ha) significantly produced lower gross returns (5 cm). Net return considerably was secured with an intra-row spacing of 20 cm (S₄), which stayed on the same bar as treatment S₃ (15 cm), and the lower net returns of ₹ 39719/ha were accumulated under S₁ (5 cm). Due to greater availability of nutrients, moisture, solar radiation, and room for growth and development, population maintenance at intra-row spacing of 20 cm (S₄) provided greatest net returns of ₹ 67408/ha and BCR 3.7.

Table 2. Effect of intra-row spacing and potassium levels on economics of sweet corn

Treatments	Gross returns (₹/ha)	Cost of cultivation (₹/ha)	Net returns (₹/ha)	BCR
Intra-row spacing (cm)				
S ₁ : 5 Intra-row + 45 Inter-rows	65575	25856	39719	2.5
S ₂ : 10 Intra-row + 45 Inter-rows	75340	25502	49838	2.9
S ₃ : 15 Intra-row + 45 Inter-rows	87995	25384	62611	3.5
S ₄ : 20 Intra-row + 45 Inter-rows	92734	25326	67408	3.7
S.Em.±	2152	-	2152	-
C.D. at 5%	6214	-	6214	-
Potassium levels (kg/ha)				
K ₁ : 0 K ₂ O + 120 N ₂ O + 60 P ₂ O ₅ kg/ha	66242	24841	41401	2.7
K ₂ : 20 K ₂ O + 120 N ₂ O + 60 P ₂ O ₅ kg/ha	73614	25292	48323	2.9
K ₃ : 40 K ₂ O + 120 N ₂ O + 60 P ₂ O ₅ kg/ha	88276	25743	62533	3.4
K ₄ : 60 K ₂ O + 120 N ₂ O + 60 P ₂ O ₅ kg/ha	93513	26193	67319	3.6
S.Em.±	2152	-	2152	-
C.D. at 5%	6214	-	6214	-
Interaction (S × K)				
S.Em.±	4303	-	4303	-
C.D. at 5%	NS	-	NS	-
C.V. %	9	-	14	-

Also, S₄ with an intra-row spacing of 20 cm acquired the greatest benefit cost ratio of 3.7, while S₁ with an intra-row spacing of 25 cm accumulated the lowest benefit cost ratio of 2.5 (5 cm). It is clear from the data (Table 2) that the application of 60 K₂O + 120 N₂O + 60 P₂O₅ (K₄), which remained equivalent to treatment K₃ (40 K₂O + 120 N₂O + 60 P₂O₅), was responsible for the higher gross returns of ₹ 93513/ha. Treatment K₁ (0 K₂O + 120 N₂O + 60 P₂O₅ kg/ha) had lower gross returns, with a total of ₹ 66242/ha. When potassium was applied at a rate of 60 K₂O + 120 N₂O + 60 P₂O₅ kg/ha (K₄), the cost of culture was found to be greatest (₹ 26193/ha), whereas treatment K₁ (0 K₂O + 120 N₂O + 60 P₂O₅ kg/ha) recorded the lowest cost of cultivation (₹ 24841/ha) [13].

4. CONCLUSIONS

Sweet corn (Sweet-16) may be planted with an intra-row spacing of 20 cm and an inter-row spacing of 45 cm, and it can be fertilised with 60 kg/ha of K₂O, 120 kg/ha of N₂O, and 60 kg/ha of P₂O₅ in addition to the recommended amounts of N and P₂O₅.

CONFERENCE DISCLAIMER

Some part of this manuscript was previously presented in the conference: 3rd International Conference IAAHAS-2023 "Innovative Approaches in Agriculture, Horticulture & Allied Sciences" on March 29-31, 2023 in SGT University, Gurugram, India. Web Link of the proceeding: <https://wikifarmer.com/event/iaahas-2023-innovative-approaches-in-agriculture-horticulture-allied-sciences/>

ACKNOWLEDGEMENT

Department of Agronomy, College of Agriculture, Junagadh Agricultural University, Junagadh, Gujarat- 362001, India for providing me the necessary resources and support for carrying out this study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Manishaben Rathod, Bavalgave VG, Patil VA, Deshmukh SP. Growth, yield and

quality of sweet corn (*Zea mays L. var. Saccharata*) as influenced by spacing and INM practices under South Gujarat condition. International Journal of Economic Plants. 2018;5(4):170-173.

2. Silva G. Nutrient removal rates by grain crops. Michigan State University Extension; 2017. Available:https://www.canr.msu.edu/news/nutrient_removal_rates_by_grain_crops.

3. Gomez KA, Gomez AA. Statistical procedures for agricultural research. An International Rice Research Institute Book. John Wiley and Sons. New York. 1984;680.

4. Shekhawat PS, Gautam RC. Effect of row spacing and weed control methods on the growth attributes and grain yield of maize under tilled and untilled conditions. Annals of Agricultural Research. 2002;23(4):626-629.

5. Bozorgi HR, Dousta HZ, Sadeghi SM, Keshavarz A, Faraji A. Study effect of plant density and nitrogen fertilizer on yield and yield components of maize. World Applied Sciences Journal. 2011;13:147-151.

6. Gozubenli H, Konuskan O. Nitrogen dose and plant density effects on popcorn grain yield. African Journal of Biotechnology. 2010;9(25):3828-3832.

7. Hamni H, Dadari SA. Growth and yield analysis of irrigated popcorn (*Zea mays everta L.*) grown in Kadawa as affected by sowing date and intra-row spacing, using correlation coefficient. ASSET Series. 2003;4:63-70.

8. Paradkar VK. Effect of N levels and plant spacing on performance of baby corn at Chindwara. Annual Progress Report, Directorate of Maize Research, IARI, New Delhi. 2004:A-77.

9. Jat G, Majumdar SP, Jat NK, Majumdar SP. Potassium and zinc fertilization of wheat in western arid zone of India. Indian Journal of Agronomy. 2013;58(1):67-71.

10. Kumar Pradeep, Kumar Rajeev, Singh SK, Kumar Anil. Effect of fertility on growth yield and yield attributes of pearl millet (*Pennisetum Glaucum L.*) under rainfed condition. Agriways. 2014;2(2):89-93.

11. Singh V, Ali J, Seema AK, Chauhan TM. Productivity, nutrient uptake and economics of wheat (*Triticum aestivum L.*) under potassium and zinc nutrition. The Indian Society of Soil Science. 2015;60(3):426-430.

12. Patel PC, Kotecha AC. Effect of phosphorus and potassium on growth characters, forage yield, nutrient uptake and quality of lucerne (*Medicago sativa* L.). Indian Journal of Agronomy. 2006;51:242-244.
13. Yadav SS, Tikkoo A, Singh JP. Effect of potassium on pearl millet-wheat cropping system in coarse textured soils of Southern Haryana. Journal of the Indian Society of Soil Science. 2012;60(2):145-149.

© 2023 Sharma et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/99784>