



Effect of Sources of Sulphur and Zinc on Yield and Economics of Sesame (*Sesamum indicum* L.)

Sovan Mondal^{a+++*}, Shikha Singh^{a#}
and Anu Nawhal^{at}

^a Department of Agronomy, SHUATS, Allahabad, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i183333

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/102471>

Original Research Article

Received: 05/05/2023

Accepted: 06/07/2023

Published: 22/07/2023

ABSTRACT

Background: A field experiment was conducted during *kharif* 2022 month of August at Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj (U.P). The experiment was laid out in Randomized Block Design with nine treatments and control which are replicated thrice on the basis of one year experiment. The treatments consisted of 3 levels of source of sulphur (Gypsum 300 kg/ha, Single super phosphate 20 kg/ha, Sulphur 40 kg/ha) and zinc (5, 10, 15 kg/ha) as a basal application and a control. The application of Gypsum 300 kg/ha + Zinc 15 kg/ha recorded significantly maximum number of Capsules/plant (51.40), Seeds/capsules (55.47), Test weight (2.70 g), Seed yield (1.07 t/ha), stover yield (2.59 t/ha), Harvest index (29.30), Gross returns (89,889.00 INR/ha), Net returns (60,180.00INR/ha), B:C ratio (2.03 INR/ha) was recorded with the treatment of Gypsum 300 kg/ha + Zinc 15 kg/ha.

^{**}M.Sc Scholar;

[#] Assistant Professor;

[†] P.hD Scholar;

*Corresponding author: E-mail: sovanmondal36703@gmail.com;

Keywords: Sesame; gypsum; single super phosphate; sulphur; zinc; yield and economics.

1. INTRODUCTION

The experiment was conducted to know the Influence of sources of sulphur and zinc on yield of Sesame (*Sesamum indicum* L.). "Sesame ranks first for having oil content of 46-64% and 6355 k cal/kg dietary energy in seeds" (Sanjay kumar & Goel, 1994). "Seeds of sesame is also rich source of protein (20 - 28%), sugar (14 - 16%) and minerals (5-7%). This oil has 85% unsaturated fatty acid viz., highly stable and has washing effect on cholesterol & prevents coronary heart disease. Sesame as a valuated oil seed appears to have numerous industrial applications. The forecasted results showed for area, production and productivity of Sesame crop for the year 2020-21 to be 51.66 thousand hectare, 17.64 thousand tonnes and 323.43 in kg/hectare respectively" [1]. "However, improved varieties and agro production technologies capable of increasing the productivity levels of sesame are now developed for different agro ecological situations in the country. A well-managed crop of sesame can yield 1200 - 1500 kg/ha under irrigated and 800 - 1000 kg/ha under rainfed conditions. The crop is grown in almost all parts of the country. More than 85% production of sesame comes from West Bengal, Madhya Pradesh, Rajasthan, Uttar Pradesh, Gujarat, Andhra Pradesh and Telangana" [1].

Single Super Phosphate is an excellent source of three plant nutrient (P_2O_5 16%, Calcium 20%, Sulphur 12%). The P component reacts in soil similarly to other soluble fertilizers. The presence of both P and sulfur (S) in SSP can be an agronomic advantage where both of these nutrients are deficient. In agronomic studies where SSP is demonstrated to be superior to other P fertilizers, it is usually due to the S and/or Ca that it contains. When locally available, SSP has found wide-spread use for fertilizing pastures where both P and S are needed. As a source of P alone, SSP often costs more than other more concentrated fertilizers, therefore it has declined in popularity [2].

"Zinc is an indispensable micronutrient for crop growth, an important component of carbonic anhydrase and a stimulator of aldolase, which are involved in carbon metabolism" [3]. "Zn is also an integral component of several biomolecules such as lipids, proteins and co-factor of auxins, and, therefore, it plays an

important role in plant nucleic acid metabolism. Zn application has been proved beneficial in improving crop yield and quality, while its deficiency reduces yield and deteriorates crop quality". Chattha et al. [4], moreover, higher Zn contents have toxic effects on plants and lead to the suppression of cell division and elongation.

2. MATERIALS AND METHODS

The experiment was conducted to know the Influence of sources of sulphur and zinc on yield of Sesame (*Sesamum indicum* L.) was carried out at Crop Research Farm of Sam Higginbottom University, Prayagraj, Uttar Pradesh during 2022. The soil of experimental plot was sandy loam intexture, nearly neutral in soil reaction (pH 7.1), low in organic carbon (0.36 %), available N (171.48 kg/ha), available P (15.2 kg/ha) and available K (232.5 kg/ha). The experiment was laid out in a Randomized Block Design consisting of Ten treatments including Control with 3 replications, with the treatment combinations (T₁) Gypsum 300 kg/ha + Zinc 5 kg/ha, (T₂) Gypsum 300 kg/ha + Zinc 10 kg/ha, (T₃) Gypsum 300 kg/ha + Zinc 15 kg/ha, (T₄) Single super phosphate 20 kg/ha + Zinc 5 kg/ha, (T₅) Single super phosphate 20 kg/ha + Zinc 10 kg/ha, (T₆) Single super phosphate 20 kg/ha + Zinc 15 kg/ha, (T₇) Sulphur 40 kg/ha + Zinc 5 kg/ha, (T₈) Sulphur 40 kg/ha + Zinc 10 kg/ha, (T₉) Sulphur 40 kg/ha + Zinc 15 kg/ha, (T₁₀) Control with spacing of row to row 30 cm and plant to plant 10 cm. The sesame crop was harvested treatment wise at harvesting maturity stage. "Yield parameters viz. number of capsules per plant, number of seeds per capsules, test weight, stover yield and seed yield were recorded manually on five randomly selected representative plants from each plot of each replication separately and after harvesting, seeds were separated from each net plot and were dried under sun for three days. Later winnowed, cleaned and seed yield per ha was computed and expressed in tonnes per hectare. After complete drying under sun for 10 days stover yield from each net plot was recorded and expressed in tonnes per hectare" [5]. The data was computed and analysed by following statistical method of Gomez and Gomez [6]. The benefit: Cost ratio was worked out after price value of seed with stover and total cost included in crop cultivation.

3. RESULTS

3.1 Yield

The perusal of the data of Number of Capsules/plants was recorded at harvest, is presented in Table 1. The data reveals that there was significant effect among different treatments on Number of Capsules/plants. Significantly Maximum Number of Capsules/plants (51.40) was recorded with the treatment of application Gypsum 300 kg/ha + Zinc 15 kg/ha over all the treatments and minimum was recorded in (37.40) control. However, treatment Gypsum 300 kg/ha + Zinc 10 kg/ha (50.67) was found to be statistically at par with Gypsum 300 kg/ha + Zinc 15 kg/ha.

Significantly Maximum test weight (2.70 g) was recorded with the treatment of application Gypsum 300 kg/ha + Zinc 15 kg/ha over all the treatments and minimum was recorded (2.27 g) in control. However, treatment Gypsum 300 kg/ha + Zinc 10 kg/ha (2.67 g) was found to be statistically at par with Gypsum 300 kg/ha + Zinc 15 kg/ha.

Significantly Maximum Number of seeds per capsules (55.47) was recorded with the treatment of application Gypsum 300 kg/ha + Zinc 15 kg/ha over all the treatments and minimum was recorded (45.20) in control. However, treatment Gypsum 300 kg/ha + Zinc 10 kg/ha (54.40) was found to be statistically at par with Gypsum 300 kg/ha + Zinc 15 kg/ha.

Significantly Maximum seed yield (1.07 t/ha) and stover yield (2.59 t/ha) was recorded with the treatment of application Gypsum 300 kg/ha + Zinc 15 kg/ha over all the treatments and minimum was recorded (0.88 t/ha) in control. However, treatment Gypsum 300 kg/ha + Zinc 10 kg/ha (1.06 t/ha) was found to be statistically at par with Gypsum 300 kg/ha + Zinc 15 kg/ha.

Significantly maximum harvest index (29.30 %) was recorded with the treatment of application of Gypsum 300 kg/ha + Zinc 15 kg/ha over all the treatments and minimum was recorded in Control (40:60:40 kg/ha) (26.40 %) and there was significant difference in between the treatments.

3.2 Economics

Higher Gross returns have been recorded with the Gypsum 300 kg/ha + Zinc 15 kg/ha (89,889)

over rest of the treatments followed by Gypsum 300 kg/ha + Zinc 15 kg/ha (57,731).

Higher net returns have been recorded with the Gypsum 300 kg/ha + Zinc 15 kg/ha (60,189) over rest of the treatments followed by Gypsum 300 kg/ha + Zinc 15 kg/ha (58,531).

Higher Benefit ratio have been recorded with the Gypsum 300 kg/ha + Zinc 15 kg/ha (2.03) over rest of the treatments followed by Gypsum 300 kg/ha + Zinc 15 kg/ha (2.00) whereas minimum Benefit ratio was recorded with Control (1.77).

4. DISCUSSION

“The observed improvement in plant height due to zinc might be due to biosynthesis of IAA growth hormones, cell enlargement, cell division and multiplication which ultimately led to better plant height of sesame and boosted plant growth”. Haritha et al [5] Similar findings were also reported by Sharma and Jain [7] in mustard, Choudhary et al. [8] in sunflower and Murthy [9] in sesame. Sulphur through gypsum might have promoted the uptake and translocation of food assimilates from source to sink effectively, resulting in higher yield attributes vi, no of capsules plant, weight of capsules plant leading to higher seed yield. These results are in agreement with the findings of Duboon et al. [10] and Ramakrishna [11] in sesame. “The increase in yield might be attributed to easy availability of sulphate (SO) sulphur present in gypsum compared to sulphide form in elemental sulphur, which essentially requires its oxidation to be converted in to SOS prior to its absorption by the plants. Among sources of sulphur gypsum proved significantly superior to other sources for seed yield have been reported” by Chaurasia et al. [12] and Ramakrishna [11]. “Zinc plays as an activator of several enzymes in plants and it is directly involved in the biosynthesis of growth substances such as auxin thereby producing more plant cells and enhanced dry matter” [13].

“The increase in yield attributes and yield due to the application of Zn might be due to fact that Zn influences on the water economy and crop growth through its effect on water uptake, root growth, maintenance of turgor, transpiration and stomatal behavior, overcomes the adverse effect of water stress and improving the drought tolerance”. Haritha et al [5] Similar results were also reported by Paraye et al. [14], Deosarkar et al. [15] in soybean.

Table 1. Effect of sulphur and zinc on yield parameters of sesame

S. no	Treatments	No. of Capsules/ plant	No. of seed/capsules	Test weight (g)	Seed yield(t/ha)	Stover yield (t/ha)
1.	Gypsum 300 kg/ha + Zinc 5 kg/ha	46.33	51.60	2.57	1.01	2.54
2.	Gypsum 300 kg/ha + Zinc 10 kg/ha	50.67	54.40	2.67	1.06	2.58
3.	Gypsum 300 kg/ha + Zinc 15 kg/ha	51.40	55.47	2.70	1.07	2.59
4.	Single super phosphate 20 kg/ha + Zinc 5 kg/ha	39.40	46.60	2.40	0.91	2.46
5.	Single super phosphate 20 kg/ha + Zinc 10 kg/ha	41.20	47.53	2.43	0.93	2.49
6.	Single super phosphate 20 kg/ha + Zinc 15 kg/ha	43.07	49.47	2.50	0.98	2.52
7.	Sulphur 40 kg/ha + Zinc 5 kg/ha	42.53	47.93	2.47	0.96	2.51
8.	Sulphur 40 kg/ha + Zinc 10 kg/ha	44.87	51.33	2.53	0.99	2.53
9.	Sulphur 40 kg/ha + Zinc 15 kg/ha	48.27	53.27	2.60	1.04	2.56
10.	Control (RDF 40:60:40 NPK kg/ha)	37.40	45.20	2.27	0.88	2.45
	F – Test	S	S	NS	S	S
	SEm (±)	0.61	0.51	0.08	0.01	0.01
	CD (p=0.05)	1.82	1.52	-	0.03	0.03

Table 2. Influence of sulphur and zinc on economics of sesame

S.no	Treatments	Cost of cultivation (INR/ha)	Gross returns (INR/ha)	Net returns (INR/ha)	B:C ratio
1.	Gypsum 300 kg/ha + Zinc 5 kg/ha	28700	83581	54881	1.91
2.	Gypsum 300 kg/ha + Zinc 10 kg/ha	29200	87731	58531	2.00
3.	Gypsum 300 kg/ha + Zinc 15 kg/ha	29700	89889	60189	2.03
4.	Single super phosphate 20 kg/ha + Zinc 5 kg/ha	27100	75779	48679	1.80
5.	Single super phosphate 20 kg/ha + Zinc 10 kg/ha	27600	76941	49341	1.79
6.	Single super phosphate 20 kg/ha + Zinc 15 kg/ha	28100	81589	53489	1.90
7.	Sulphur 40 kg/ha + Zinc 5 kg/ha	30500	79680	49180	1.61
8.	Sulphur 40 kg/ha + Zinc 10 kg/ha	31000	81921	50921	1.64
9.	Sulphur 40 kg/ha + Zinc 15 kg/ha	31500	86320	54820	1.74
10.	Control (RDF 40:60:40 NPK kg/ha)	26400	73040	46640	1.77

5. CONCLUSION

It was concluded that for obtaining higher yield components with better quality of Sesame application of Gypsum 300 kg/ha + Zinc 15 kg/ha was recorded significantly higher number of Capsules/plant, Seeds/capsules, seed yield, stover yield, harvest index and benefit cost ratio as compared to other treatments. Since, the finding based on the research done in one season.

ACKNOWLEDGEMENT

I express my gratitude to the Department of Agronomy & all higher authorities of university for providing us with all facilities for the study. The Authors are thankful to the staff of Agronomy department and staff of Post graduate seed Testing laboratory for their cooperation during the study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Priyanka Evangilin N, Ramana Murthy B, Mohan Naidu G, Aparna B. Statistical model for forecasting area, production and productivity of sesame crop (*Sesamum indicum* L.) in Andhra Pradesh, India. Int. J. Curr. Microbiol. App. Sci. 2020;9(07): 1156-1166.
2. Nutrient Source Specifics (No. 21), International Plant Nutrition Institute.
3. Tsonev T, Lidon FJC. Zinc in plants—An overview. Emir. J. Food Agric. 2012;24: 322–333.
4. Chattha MU, Hassan MU, Khan I, Chattha MB, Mahmood A, Chattha MU, Nawaz M, Subhani MN, Kharal M, Khan S. Biofortification of wheat cultivars to combat zinc deficiency. Front. Plant Sci. 2017; 8:281.
5. Haritha G, Singh R, Indu T. Studies on growth and yield of sesame (*Sesamum indicum* L.) as influenced by spacing and foliar application of zinc. The Pharma Innovation Journal. 2022; 11(3):2366-9.
6. Gomez KA, Gomez AA. Statistical Procedures for Agricultural Research. Edition 2. Wiley Inter Science Publications and John Wiley and Sons, New York; 1984.
7. Sharma PP, Jain NK. Effect of foliar sprays of agrochemicals on growth and yield of Indian mustard (*Brassica juncea* L.). Indian Journal of Agricultural Sciences. 2003; 73:381-383.
8. Choudhary AR, Prabhakara S, Nagarathna TK. Growth and yield of sunflower (*Helianthus annuus* L.) as influenced by micronutrients application in Alfisols. Karnataka Journal of Agricultural Sciences. 2010;23:495-496.
9. Murthy IYLN. Zinc response to oilseed crops. Indian Journal of Fertilisers. 2011;7(10):104-117.
10. Duboon SS, Deshmukh MR, Jyotishi A, Jam HC. Effect of sources and levels of sulphur on seed and oil yield of sesame (*Sesamum indicum* L.) under different agro-climatic situation of India Journal of Oilseed Research. 2005;21(1):199-200.
11. Ramakrishna B. Growth and yield of sesame (*Sesamum indicum* L.) as influenced by sources and levels of sulphur (Master thesis) university of Agriculture Science Dharmad Karnataka (India) lit pknshikoch granth ac in handle 1 \$6993; 2013. (Access on March 21. 2021).
12. Chaurasia SK, Jain N, Jain N. Effects of integrated use of fertilizer, organic manure and micronutrients on productivity of sesame (*Secconum indicum* L.) Annale Agriculture Research New Series. 2009; 30(384):91-94.
13. Christopher A, Oluwagbenga D, Aruna O, Adekiya CC, Khadijat O, Suleiman FO, Okunlola CO, Aremu IO, Owolabi C, Temidayo AJ. Zinc sulphate and boron-based foliar fertilizer effect on growth, yield, minerals, and heavy metal composition of groundnut (*Arachis hypogaea* L) grown on an alfisol. International J. Agronomy. 2019.3:123-127.
14. Paraye PM, Kashyap TL, Chaubey AK. Effect of nutrient management practices and rowspacing on growth yield attributes and yield of Indian mustard (*Brassica juncea* L.). Environment and Ecology. 2009;27(2):589–591.

15. Deosarkar DB, Patinge SP, Bhosale AM, Deshmukh SB. Effect of micronutrient on seed yield of soybean. Annals of Plant Physiology. 2001;15:163-166.

© 2023 Mondal et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/102471>