



# Effect of Phosphorus Application on Groundnut (*Arachis hypogaea* L.): A Review

Kamal<sup>a+++\*</sup>, Ekta Kamboj<sup>a#</sup>, Amit Sharma<sup>a++</sup>, Ravi<sup>a++</sup>,  
B. K. Dhaka<sup>b†</sup> and Preeti<sup>a++</sup>

<sup>a</sup> Department of Agronomy, College of Agriculture, Chaudhary Charan Singh Haryana Agricultural University, Hisar, Haryana, India.

<sup>b</sup> Department of Agronomy, Lovely Professional University, Punjab, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJPSS/2023/v35i183423

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/103501>

Review Article

Received: 20/05/2023

Accepted: 22/07/2023

Published: 03/08/2023

## ABSTRACT

The phosphorus in legumes plays an essential role as part of the membrane system of the cell, chloroplast and the mitochondria. Groundnut, being a legume oilseed crop, requires a significant amount of phosphorus. Plants use it for energy transfer in metabolic processes such as photosynthesis (in the form of ATP and ADP), starch and sugar transformation and nutrient movement. In addition to promoting cell division and the development of good root systems phosphorus plays an important role for ensuring timely and uniform ripening of crops, promoting carbohydrate breakdown for energy release, and hastening maturity, it is an essential component for ensuring the growth and maturity of plants. Phosphorus is one of the principal nutrients, which boosts root growth, thus facilitating water and nutrient uptake from deeper layers of the soil. The

<sup>++</sup> Ph.D. Scholar;

<sup>#</sup> Assistant Scientist;

<sup>†</sup> M.Sc. Scholar;

\*Corresponding author: E-mail: kamalkhroad@gmail.com;

groundnut crop calls for sensible supply of phosphorus for its regular growth. Phosphorus has been found to improve the crop's effective use of soil nutrients and increase biological N fixation by increasing nitrogenase activity. Various research in this paper clearly indicate that application of phosphorus significantly affects the growth, phenology, yield attributes, yield, quality, nutrient uptake and economics of groundnut.

**Keywords:** Growth; phenology; phosphorus; uptake; yield; economics.

## 1. INTRODUCTION

Groundnut (*Arachis hypogaea* L.) is one of the best-known oilseed crops which belongs to the family Leguminosae and sub-family Papilionaceae. It is believed that it has been originated in South America. [1]. The groundnut crop calls for sensible supply of phosphorus for its regular growth. In India, Groundnut is considered the "king of oilseeds". This oil seed crop is primarily found in tropical and subtropical regions, and is also known as peanut, earthnut, monkey nut and poor men's cashew nut. The global area, production and productivity of groundnut is 29.7 million ha, 50.8 million tonnes and 17.1 quintal ha<sup>-1</sup>, respectively. China is the world's largest producer of groundnut with 4.6 million ha area, 17 million tonnes production and productivity of 37.0 quintal ha<sup>-1</sup> followed by India. Groundnut covers an area of 4.8 million ha with production of 9.9 million tonnes and productivity of 20.6 quintal ha<sup>-1</sup> in India during 2019-20 [2]. Groundnut account for 31.7% of India total oilseed production and about 28.3 % of the cultivated area of total oilseeds [3]. The major groundnut producing states in India are Gujrat, Rajasthan and Tamil Nadu. Phosphorus is considered a crucial mineral fertilizer for the flourishing production of the crop [4]. Both organic and inorganic forms of phosphorus are found in soils, and the relative amounts of each form vary substantially amongst soils. Because it is strongly bound to soil colloids, organic P is typically not available for plant uptake unless it is broken down and released during the mineralization process [5]. Plants can access phosphorus in its inorganic forms (H<sub>2</sub>PO<sub>4</sub><sup>-</sup> and HPO<sub>4</sub><sup>-</sup>), but very little of it is present in soil solution [6]. Moisture and temperature have a major impact on the chemistry of P cycling in soils, which is quite complex. To meet the nutrient requirements of plants at each stage of their growth cycle, Labile P one of many potential forms of phosphorus must be supplemented regularly. By adding inorganic P fertilisers, the concentration of phosphorus in the soil solution is increased [7]. For the synthesis of chlorophyll, phosphorus is an essential component. For cell division and the growth of meristematic tissue,

phosphorus is considered an essential component of the cell nucleus [8]. By fixing atmospheric nitrogen within its root nodules, the groundnut crop improves soil fertility. Application of phosphorus determines plant reproductive efficiency and promotes plant growth, development and yield [9]. Available literature on the effect of phosphorus in groundnut is presented under following sub-heads.

## 2. EFFECT OF PHOSPHORUS ON GROWTH PARAMETERS

Jat et al. [10] conducted a study at MPUAT, Udaipur and purported that application of 80 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> significantly increased chlorophyll content (1.535 mg/g) in groundnut, which was significantly higher over 60 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>, 40 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>, 20 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> and 0 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>. Hinduja et al. [11] avouched that application of 70 kg/ha Phosphorus + 40 kg/ha Sulphur in groundnut significantly increase plant height (60.3 cm), number of nodules (79.88 plant<sup>-1</sup>) and dry matter accumulation (32.02 g plant<sup>-1</sup>) in groundnut which was significantly higher over 40 kg/ha phosphorus + 20 kg/ha sulphur by 3.8, 16.1 and 14.8 per cent, respectively. Tekulu et al. [12] conducted study in Ethiopia and observed that application of 30 kg N and 69 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> was highly correlated and produced highest plant height (56.1 cm), which was significantly higher over other treatment combinations in groundnut. Lro et al. [13] conducted a field experiment in Nigeria and revealed that application of 60 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> significantly improved plant height of groundnut (34.6 cm) over 40 kg P<sub>2</sub>O<sub>5</sub>, 20 kg P<sub>2</sub>O<sub>5</sub> and 0 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> by 19.0, 33.3, 62.9 per cent, respectively.

Sibhatu et al. [14] conducted research at Ethiopia and professed that application of 30 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> in groundnut produced highest plant (23.5 cm), which was statistically at par with application of 40 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> and significantly higher over 20 kg, 10 kg and 0 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> by 19.8, 22.1 and 46.1 per cent, respectively. Jeetarwal et al. [15] conducted a field experiment at S.K.N. College of Agriculture, Jobner (Rajasthan) and averred that

application of 60 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> in groundnut produce higher plant height (22.70 cm), dry matter (450.6 g/m row length), number of branches (10.15 plant<sup>-1</sup>), chlorophyll content (0.67 mg/g) and number of nodules (58.10 plant<sup>-1</sup>), which were statistically at par with 40 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> and significantly higher over 20 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> and 0 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>.

Singh et al. [16] conducted an experiment on groundnut at Morena (MP) and reviewed that application of 20 kg N + 40 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> produce higher plant height (66.5 cm), number of primary branch (9.2 plant<sup>-1</sup>), number of secondary branch (21.5 plant<sup>-1</sup>), number of root nodules (174 plant<sup>-1</sup>), dry weight of root nodules (549 mg plant<sup>-1</sup>), tap root length (14.1 cm) and dry weight of root (6.68 g plant<sup>-1</sup>), which were significantly higher over control by 34.1, 33.3, 35.2, 43.8, 44.0, 74.0 and 54.2 per cent, respectively.

Kabir et al. [17] conducted a study at Bangladesh Agricultural University, Mymensingh and stated that application of 50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> in groundnut produce higher plant height (59.6 cm), number of branches (7.5 plant<sup>-1</sup>), dry weight (32.0 plant<sup>-1</sup>), crop growth rate (21.0 g/m<sup>2</sup>/day) between 70-100 DAS and leaf area index (1.7), which were significantly higher over control by 26.4, 20.9, 15.6, 15.5 and 17.3 per cent, respectively. Singh et al. [18] opined that application of 20 kg N + 60 kg P<sub>2</sub>O<sub>5</sub> + 40 kg K<sub>2</sub>O/ha progressively increased root length, shoot length and seedling length of groundnut, which were significantly higher over control by 43.1, 48.7 and 42.7 per cent, respectively. Kamara et al. [19] adduced that in groundnut application of 40 kg P<sub>2</sub>O<sub>5</sub>/ha produced highest plant height (42.3 cm), which was significantly higher over control by 14.0 percent. Karunakaran et al. [20] conducted a field experiment at Tamil Nadu in groundnut and testified that application of 125 % RDF (17 kg N + 34 kg P<sub>2</sub>O<sub>5</sub> + 54 kg K<sub>2</sub>O/ha) increase height of plant (51.8 cm) as compare to RDF (48.3 cm). Salve et al. [21] conducted a field experiment at MPKV, Rahuri and averred that application of 75 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> in groundnut produce highest plant height (34.34 cm) and number of branches (48.45 plant<sup>-1</sup>), which were statistically at par with 50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> and significantly higher over 25 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>. Shiyam [22] concluded that increasing the levels of phosphorus up to 50 kg ha<sup>-1</sup> has a significant effect on plant height, and the highest plant height (56 cm) was found with the application of 50 kg P ha<sup>-1</sup> in groundnut.

Rao and Shaktawat [23] carried out an experiment at Udaipur, Rajasthan and averred

that application of 60 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> in groundnut produced higher number of branches plant<sup>-1</sup> (4.70), leaf area index (4.41), root dry weight plant<sup>-1</sup> (0.72 g), which were significantly higher over 40 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> and 20 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> by 2.2, 2.5, 1.3 and 9.3, 8.6, 9.8 per cent, respectively.

### 3. PHENOLOGY

Tekulu et al. [12] conducted study in Ethiopia and purported that individual N and P fertilizer rates, as well as combinations of N and P fertilizer rates, had no effect on groundnut phenological parameters such as flowering time and maturity. This could be because biological N fixation provided enough nitrogen for the groundnut plant to begin flowering and maturation.

Lro et al. [13] did a field experiment in Nigeria on groundnut and professed that non significant variation in germination percentage was recorded between phosphorus fertilizer application rates. However, the control has a greater germination percentage (68.9 %) than the other treatments (20, 40, and 60 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>) and also concluded that phosphorus rates have a significant effect on days to 50% flowering. The application rate of 60 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> resulted minimum number of days for groundnut to reach 50% flowering (27.7 days), 40 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> and 20 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> reached 50% flowering at the same time (29.5 days), and the control (0 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>) reached 50% flowering at 30.5 days.

Melese and Dechassa [24] purported that applying phosphorus and manure to groundnut genotypes Werer 962 and Oldhale had no influence on flowering and maturity. Sibhatu et al. [14] averred that the highest days to 90% physiological maturity of groundnut (99.0) was obtained with application of 30 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>, while significantly lower days to 90 % physiological maturity (91.3) were obtained in no application of phosphorus So, the crop fertilized with higher application of phosphorus took long time to mature while less phosphorus fertilizer shortened maturity date.

Bala et al. [25] investigated that days to 50% flowering were increased by the application of 30 kg N + 39 kg P<sub>2</sub>O<sub>5</sub> + 39 K<sub>2</sub>O ha<sup>-1</sup> as compared to 10 kg N + 13 kg P<sub>2</sub>O<sub>5</sub> + 13 K<sub>2</sub>O ha<sup>-1</sup> in groundnut. Kamara et al. [19] conducted research and inferred that in groundnut the application of 40 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> resulted in the highest percent field emergence (85.33), while the application of 0 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> resulted in the shortest days to 50%

flowering (25) and the shortest days to maturity (89). Khan et al. [26] reviewed that the application of 27 kg N ha<sup>-1</sup>+ 69 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> in groundnut significantly increased the emergence of plants (11.71 plants m<sup>-2</sup>), days to 50% flowering (33.33) and days to maturity (183.28) in comparison with the control treatment.

#### 4. YIELD ATTRIBUTES, YIELD AND QUALITY

Jat et al. [10] conducted a study at MPUAT, Udaipur and purported that application of 80 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> significantly increased kernel yield (1179 kg ha<sup>-1</sup>) and biological yield (4894 kg ha<sup>-1</sup>) in groundnut, which were statistically at par with application of 60 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> and significantly higher over 40 kg, 20 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> and control. Kumar et al. [27] conducted a field experiment at Odisha in groundnut and averred that application of 100 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> produced higher number of higher number of pods/plant (15.55), pod yield (2112 kg/ha), kernel yield (1370 kg/ha), oil content (47.92 %) and oil yield (998.99 kg/ha), which were significantly higher over control by 76.7, 43.8, 34.7, 12.6 and 51.0 per cent, respectively.

Choudhary and Yadav [28] conducted a field experiment at Anand, Gujarat in groundnut and anticipated that application of 50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> produced higher pod yield (2657 kg ha<sup>-1</sup>) and haulm yield (4467 kg ha<sup>-1</sup>) which were significantly higher over control by 20.3 and 18.9 per cent, respectively. Ikenganyia et al. [29] purported that application of 75 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> produced higher pod yield of groundnut compared to 50 kg P<sub>2</sub>O<sub>5</sub>, 25 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> and control by 15, 32 and 54 per cent, respectively. Sagvekar et al. [30] observed that NP fertilizer ratio 0.40 significantly increased number of pods (30.6 plant<sup>-1</sup>), pod yield (3500 kg ha<sup>-1</sup>) and kernel yield (2630 kg ha<sup>-1</sup>) in groundnut over rest of NP ratio.

Sibhatu et al. [14] conducted research at Ethiopia in groundnut and concluded that increasing the levels of phosphorus up to 20 kg ha<sup>-1</sup> produced higher number of pod (22.6 plant<sup>-1</sup>), seed (3.5 pod<sup>-1</sup>), 100-kernel weight (42.2 g), shelled seed yield (1269 kg ha<sup>-1</sup>) and dry matter yield (3765 kg ha<sup>-1</sup>), which were significantly higher over 10 kg and 0 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> by 31.2, 7.53, 1.5, 7.0, 2.0 and 33.9, 24.8, 3.9, 34.0, 16.6 per cent, respectively.

Kumar et al. [31] reviewed that use of N@7 kg/ha by urea at 60 DAS and NP fertilizer ratio

will be 0.57 significantly increased total pods per plant (29.53), filled pods per plant (24.27), 100-kernel weight (41.16 g), 100-pod weight (135.33 g) and pod yield (4361 kg ha<sup>-1</sup>) in groundnut over other NP fertilizer ratios. Yadav et al. [32] asserted that increasing the level of phosphorus up to 27 kg ha<sup>-1</sup> in groundnut progressively increased the number of pods (70.8 plant<sup>-1</sup>), number of kernels (2.0 pod<sup>-1</sup>), pod yield (2.09 t ha<sup>-1</sup>) and haulm yield (3.58 t ha<sup>-1</sup>), which were significantly higher over control by 60.9, 17.6, 38.4 and 39.2 per cent, respectively.

Jeetarwal et al. [15] conducted a field experiment at S.K.N. College of Agriculture, Jobner (Rajasthan) averred that application of 60 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> in groundnut produce higher number of pods/plant (20.4), number of kernels/pod (1.95), seed index (59.2 g), pod yield (2.83 t/ha), haulm yield (6.85 t/ha), kernels yield (1.98 t/ha) and shelling (69.96 %), which were statistically at par with 40 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> and significantly higher over 20 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> and 0 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>. Singh et al. [16] conducted an experiment at Morena (MP) in groundnut and reviewed that application of 20 kg N + 40 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> produced higher number of pod (27.7 plant<sup>-1</sup>), pod yield (2535 kg ha<sup>-1</sup>), haulm yield (3103 kg ha<sup>-1</sup>), oil content (43.8 %) and protein content (27.04 %), which were significantly higher over control by 33.8, 48.2, 33.9, 3.9 and 11.8 per cent, respectively.

Kabir et al. [17] conducted a study at Bangladesh Agricultural University, Mymensingh and averred that application of 50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> in groundnut gave higher number of pods (18.9 plant<sup>-1</sup>), 100-pod weight (110.8 g), shelling (64.2 %), pod yield (2.8 t ha<sup>-1</sup>), biological yield (10.8 t ha<sup>-1</sup>), straw yield (8.02 t ha<sup>-1</sup>) which was significantly higher over 25 kg P<sub>2</sub>O<sub>5</sub> and 0 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> by 6.4, 0.8, 1.7, 15.3, 6.1, 3.0 and 14.0, 2.4, 10.6, 36.6, 21.5, 17.2 per cent, respectively.

Akbari et al. [33] carried out a field experiment at Gujarat and reckoned that application of 50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> significantly increased haulm yield, pod yield, shelling out percentage and harvest index of groundnut compared to lower doses of phosphorus. Kamara et al. [19] propounded that increasing the levels of phosphorus up to 40 kg ha<sup>-1</sup> significantly increased pod yield in groundnut compared to other lower doses. Nwokwu [34] observed a significant increase in kernel yield of groundnut when 60 kg of P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> was used, which was significantly higher than control by 39.0 percent.

John [35] averred that application of 40 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> significantly increased pod yield, haulm yield and kernel yield of groundnut over control. Salve et al. [21] conducted a field experiment at MPKV, Rahuri and averred that application of 75 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> in groundnut produce highest matured pods/plant (18.34), dry pod yield (34.74 q/ha), haulm yield (43.75 q/ha), oil content (51.50 %), oil yield (13.18 q/ha), protein yield (6.65 q/ha), which were statistically at par with 50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> and significantly higher over 25 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>. Naab et al [36] observed that the application of 30 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> resulted in a significant increase in pod and kernel yield of groundnut as compared to 0 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> but it was statistically equivalent to 90 kg P<sub>2</sub>O<sub>5</sub> and 60 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>.

Kumar et al. [37] averred that application of 60 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> significantly increased pod yield of groundnut compared to 0, 20 and 40 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>. Rajanikant et al. [38] conducted a field experiment at Hyderabad and purported that application of 40 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> significantly increased pod yield and shelling out percentage in groundnut as compare to control. Basu et al. [39] conducted a field experiment at Kharagpur and found that RDF (20 kg N + 40 kg P<sub>2</sub>O<sub>5</sub> + 40 kg K<sub>2</sub>O ha<sup>-1</sup>) significantly increased the pod yield (1465 kg/ha) in groundnut over other treatment combinations. Kausale et al. [40] investigated that increasing the level of phosphorus up to 30 kg ha<sup>-1</sup> in groundnut significantly increased pod yield (2754 kg ha<sup>-1</sup>) over other lower doses of phosphorus. Ranjit et al. [41] conducted a field experiment and professed that use of 112.5 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> in groundnut produced higher haulm and pod yield of groundnut 4600 kg ha<sup>-1</sup> and 3500 kg ha<sup>-1</sup>, respectively.

Gobarah et al. [42] avouched that increasing the levels of phosphorus from 30 to 60 kg/ha significantly increased number of pods plant<sup>-1</sup>, 100-kernel weight, kernel yield, protein and oil content in groundnut. Karmakar et al. [43] conducted an experiment at West Bengal in groundnut and asserted that 30 kg N + 60 kg P<sub>2</sub>O<sub>5</sub> + 20 kg K<sub>2</sub>O ha<sup>-1</sup> produced higher haulm yield (4643 kg/ha) and pod yield (2715 kg/ha) compared to other treatment combinations. Panwar and Singh [44] reviewed that by application of 60 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> significantly increased pod yield (28.40 q ha<sup>-1</sup>) and harvest index (36.5 %) of groundnut as compared to control. Kachot et al. [45] examined that application of 25 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> significantly increase shelling out percentage, oil content and protein content in groundnut over control.

Kumaran [46] reviewed that application of RDF (34 kg N + 17 kg P<sub>2</sub>O<sub>5</sub> + 54 kg K<sub>2</sub>O/ha) + FYM as basal @ 12.5 t/ha + 17 kg P<sub>2</sub>O<sub>5</sub>/ha at 30 DAS in groundnut produced higher number of pods (14.95 plant<sup>-1</sup>) and seed index (30.23 g) which was significantly higher over application of only RDF by 22.8 and 9.5 per cent, respectively. Majumdar et al. [47] conducted a field experiment and stated that application of 75 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> significantly increased haulm (4670 kg ha<sup>-1</sup>) and pod yield (3000 kg ha<sup>-1</sup>) of groundnut whereas, oil and protein content was significantly higher over control by 8.3 and 9.7 per cent, respectively. Rao and Shaktawat [23] carried out an experiment at Udaipur, Rajasthan in groundnut and opined that application of 60 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> produced higher pod yield (18.54 q ha<sup>-1</sup>), biological yield (50.93 q ha<sup>-1</sup>), oil content (49.2 %), protein content (24.3 %), which were significantly higher over 40 kg and 20 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> by 3.6, 3.9, 2.5, 3.0 and 13.6, 9.3, 5.3, 5.1 per cent, respectively.

## 5. NUTRIENT CONTENT AND UPTAKE

Kamal et al. [9] conducted a field experiment at CCS HAU, Hisar, Haryana, and inferred that application of 60 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> progressively increased N uptake (180.5 kg ha<sup>-1</sup>), P uptake (29.4 kg ha<sup>-1</sup>), K uptake (96.4 kg ha<sup>-1</sup>) and S uptake (23.3 kg ha<sup>-1</sup>) in groundnut, which were significantly higher over 50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>, 40 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> and control. Musa et al. [48] conducted a field experiment at Nigeria in groundnut and reproduced that increasing the levels of phosphorus up to 24 kg ha<sup>-1</sup> progressively increased nitrogen (12.5 kg ha<sup>-1</sup>) and phosphorus (4.6 kg ha<sup>-1</sup>) uptake in haulm, which were significantly higher over control by 50.4 and 38.4 per cent, respectively.

Choudhary and Yadav [28] conducted a field experiment at Anand, Gujarat in groundnut and averred that application of 50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> progressively increased phosphorus content in kernel (0.48 %) and phosphorus uptake by kernel (13.25 kg ha<sup>-1</sup>), which were significantly higher over control by 14.3 and 16.1 per cent, respectively. Chirwa et al. [49] conducted a field experiment at Tanzania and concluded that application of 20 kg N + 30 kg P ha<sup>-1</sup> in groundnut significantly increased uptake of N, P and Ca compared to other treatment combination and control. Yadav et al. [32] averred that increasing the level of phosphorus up to 27 kg ha<sup>-1</sup> in groundnut progressively increased uptake of N (120.5 kg ha<sup>-1</sup>), P<sub>2</sub>O<sub>5</sub> (17.7 kg ha<sup>-1</sup>) and K

(71.5 kg ha<sup>-1</sup>), which were significantly higher over control by 43.1, 48.6 and 42.7 per cent, respectively.

Singh et al. [16] conducted an experiment at Morena (MP) and avouched that application of 20 kg N + 40 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> progressively increased phosphorus uptake in kernel (9.23 kg ha<sup>-1</sup>) and stover (8.62 kg ha<sup>-1</sup>), which were significantly higher over control by 100.2 and 119.3 per cent, respectively. Salve et al. [21] conducted a field experiment at MPKV, Rahuri and inferred that application of 75 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> progressively increased N uptake (202.21 kg ha<sup>-1</sup>), P uptake (28.53 kg ha<sup>-1</sup>) and K uptake (92.51 kg ha<sup>-1</sup>) in groundnut, which were significantly higher over 50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> and 25 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>. Ranjit et al. [41] found that applying 112.5 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> in groundnut significantly increased uptake of N, P, K, and Ca compared to lesser phosphorus doses. Mirvat et al. [50] averred that increasing the phosphorus dose from 30 to 60 kg P<sub>2</sub>O<sub>5</sub> faddan<sup>-1</sup> significantly increase protein and NPK content in kernel of groundnut considerably over control. Deka et al. [51] adduced that applying 50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> significantly increased N, P and K uptake in groundnut over control, while 75 and 100 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> had similar results.

## 6. ECONOMICS

Hinduja et al. [11] adduced that maximum net return (₹1,38,787/ha) and benefit cost ratio (1.82) in groundnut were obtained with phosphorus dose 70 kg/ha + sulphur dose 40 kg/ha, which were significantly higher than rest of the treatments. Vali et al. [52] found that application of 50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> + 30 kg Zn ha<sup>-1</sup> fetched highest net returns (₹1,39,103/ha) and benefit cost ratio (1.8) in groundnut, which were significantly higher over 30 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> + 0 kg Zn ha<sup>-1</sup> by 37.1 and 29.5 per cent, respectively.

Kumar et al. [27] conducted a field experiment at Odisha in groundnut and averred that application of 100 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> fetched highest gross return (₹97,152/ha), net return (₹58,858/ha) and benefit cost ratio (1.53), which were significantly higher over control by 43.8, 94.5 and 88.8 per cent, respectively. Sagvekar et al. [30] testified that using 30 kg N ha<sup>-1</sup> + 75 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> resulted in a greater net return (₹58,826 ha<sup>-1</sup>) and benefit cost ratio (1.78), which were significantly higher over other treatment combinations in groundnut. Yadav et al. [32] reported that increasing the level of phosphorus up to 27 kg ha<sup>-1</sup>

progressively increased net returns (₹65,450/ha) and benefit cost ratio (2.54) in groundnut, which were significantly higher over control by 14.1 and 11.9 per cent, respectively.

Singh et al. [16] examined that application of 20 kg N + 40 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> fetched highest net returns (₹30,123/ha) and benefit cost ratio (2.99) in groundnut, which were significantly higher over control by 73.9 and 20.0 per cent, respectively. Vishwakarma et al. [53] reported that highest gross return (₹ 29,000 ha<sup>-1</sup>) and benefit cost ratio (2.23) obtained with application of 30 kg N ha<sup>-1</sup> + 50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> + 40 kg K<sub>2</sub>O ha<sup>-1</sup>, while lowest gross return (₹ 12,060 ha<sup>-1</sup>) and benefit cost ratio (1.21) obtained with control in groundnut.

Patra and Bandyopadhyay [54] found that 150 percent phosphorus through phosphocompost grade-I had the highest net return (₹ 28590.47), while 100 percent phosphorus through phosphocompost grade-I had the highest B:C ratio (1.68) and they were far more cost-effective than sole application of chemical fertilizer and also combined sources of P application in groundnut.

## 7. CONCLUSION

It is concluded that phosphorus plays an important role to increase the yield of groundnut. The optimum dose of phosphorus significantly enhanced oil content, protein content, pod yield, haulm yield and prolonged crop maturity besides significant improvement in the performance of various growth parameters and dry matter production in groundnut.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Hussainy SAH, Brindavathy R, Vaidyanathan R. Influence of Irrigation Regimes on the Performance of Groundnut (*Arachis hypogaea*) under Intercropping Situation. Legume Research- An International Journal. 2023;46(4):496-501.
2. Anonymous. Department of economics and statistics, Ministry of agriculture cooperation and farmers welfare, Government of India; 2020.
3. Ali MA, Pal AK, Baidya A, Gunri SK. Variation in dry matter production,

- partitioning, yield and its correlation in groundnut (*Arachis hypogaea* L.) genotypes. Legume Research- An International Journal. 2021;44(6):706-711.
4. Hasan M, Uddin Md K, Mohamed MTM, Zuan ATK, Motmainna M, Haque ANA. Effect of nitrogen and phosphorus fertilizers on growth, yield, nodulation and nutritional composition of bambara groundnut [*Vigna subterranea* (L.) Verdc.]. Legume Research- An International Journal. 2021;44(12):1437-1442.
  5. Filippelli GM. The global phosphorus cycle. Review on Mineral Geochemistry. 2002;48: 391-425.
  6. Weil RR, Brady NC. Soil phosphorus and potassium. The Nature and Properties of Soils. 2017:643-695.
  7. Sikka R, Kaur S, Gupta RK. Effect of phosphorus application on yield and its uptake by soybean (*Glycine max* L.) in different cropping systems. Indian Journal of Agricultural Research. 2022;56(3):308-312.
  8. Kumar R, Rathore DK, Singh M, Kumar P, Khippal A. Effect of phosphorus and zinc nutrition on growth and yield of fodder cowpea. Legume Research- An International Journal. 2016;39(2):262-267.
  9. Kamal, Dhaka AK, Prakash R, Sharma A, Dhaka BK. Effect of phosphorus and sulphur levels on nutrient content and uptake of groundnut (*Arachis hypogaea* L.). Biological Forum– An International Journal. 2023;15(2):1023-1026.
  10. Jat H, Kaushik MK, Choudhary JL, Meena VK, Meena SK. Weed management in groundnut (*Arachis hypogaea* L.) with phosphorus in Southern Rajasthan. Legume Research- An International Journal. 2023;46(4):513-518.
  11. Hinduja N, Singh S, Tiwari D, Mahapatra A, Mahanta BS, Kumar S. Effect of phosphorus and sulphur on growth and yield of groundnut (*Arachis hypogaea* L.). The Bioscan. 2020;15(4):459-462.
  12. Tekulu K, Taye G, Dereje A. Effect of starter nitrogen and phosphorus fertilizer rates on yield and yield components, grain protein content of groundnut (*Arachis hypogaea* L.) and residual soil nitrogen content in a semiarid north Ethiopia. Heliyon. 2020;6:e05101.
  13. Lro LL, Jameela A, Ninani KN. Growth and yield components of groundnut (*Arachis hypogaea* L.) as affected by phosphorus fertilizer application on the Jos Plateau. Asian Journal of Research in Agriculture and Forestry. 2019;3(3):1-8.
  14. Sibhatu B, Tekle G, Harfe M. Response of groundnut (*Arachis hypogaea* L.) to different rates of phosphorus fertilizer at Tanqua Abergelle District, Northern Ethiopia. Basic Research Journal of Agricultural Science and Review. 2016;5: 24-29.
  15. Jeetarwal RL, Jat NL, Dhaka MS, Jat ML, Naga SD. Performance of groundnut (*Arachis hypogaea* L.) as influenced by phosphorus and zinc fertilization. Annals of agricultural research. 2014;35(4):411-415.
  16. Singh YP, Singh S, Dubey SK, Tomar R. Organic, inorganic sources of phosphorus and method of application on performance of groundnut (*Arachis hypogaea* L.) under rainfed condition. Indian Journal of Soil Conservation. 2014;42(2):204-208.
  17. Kabir R, Yeasmin S, Islam AKMM, Sarkar MAR. Effect of phosphorus, calcium and boron on the growth and yield of groundnut (*Arachis hypogaea* L.). International Journal of Bio-Science and Bio-Technology. 2013; 5(3):51-60.
  18. Singh GP, Singh PL, Panwar AS. Seed yield, quality and nutrient uptake of groundnut (*Arachis hypogaea*) as affected by integrated nutrient management in mid hill altitude of Meghalaya, India. Legume Research- An International Journal. 2013;36(2):147-152.
  19. Kamara EG, Olympio NS, Asibuo JY. Effect of calcium and phosphorus fertilizer on the growth and yield of groundnut (*Arachis hypogaea* L.). International Research Journal of Agricultural Science and Soil Science. 2011;1(8):326-31.
  20. Karunakaran V, Rammohan J, Chellamuthu V, Poonghuzhalan R. Effect of integrated nutrient management on the growth and yield of groundnut (*Arachis hypogaea*) in coastal region of Karaikal. Indian Journal of Agronomy. 2010;55(2): 128-132.
  21. Salve YV, Jadhav AS, Lambade BM. Response of summer groundnut (*Arachis hypogaea* L.) to varying levels of phosphorus and potassium. Journal of Maharashtra Agricultural Universities. 2010;35(2):178-183.
  22. Shiyam JO. Growth and yield response of groundnut (*Arachis hypogaea* L.) to plant densities and phosphorus on an Ultisol in Southeastern Nigeria. Libyan Agriculture

- Research Center Journal International. 2010;1:211-14.
23. Rao SS, Shaktawat MS. Effect of organic manure, phosphorus and growth, yield and quality of groundnut. Indian Journal of Plant Physiology. 2001;6(3):306-311.
  24. Melese B, Dechassa N. Seed yield of groundnut (*Arachis hypogaea* L.) as influenced by phosphorus and manure application at Babile, eastern Ethiopia. International Journal of Advanced Biological and Biomedical Research. 2017;6(1):399–404.
  25. Bala HMB, Ogunlela VB, Tanimu B, Kuchinda NC. Response of two groundnut (*Arachis hypogaea* L.) varieties to sowing date and NPK fertilizer rate in a semi-arid environment: growth and growth attributes. Asian Journal of Crop Science. 2011;3(3): 141-50.
  26. Khan N, Faridullah, Uddin MI. Agronomic characters of groundnut (*Arachis hypogaea* L.) genotypes as affected by nitrogen and phosphorus fertilization under rainfed condition. Electronic Journal of Environmental, Agricultural and Food Chemistry. 2009;8:61-68.
  27. Kumar DP, Mandal TK, Zaman A, Pal A. Studies on the effect of phosphorus levels on yield attributes and yield of groundnut in South Odisha condition. International Journal of Agriculture, Environment and Biotechnology. 2019;12(4):339-344.
  28. Choudhary RR, Yadav HL. Effect of conjunctive use of phosphorus and bio-organics on phosphorus content and uptake in summer groundnut (*Arachis hypogaea* L.). International Journal of Current Microbiology and Applied Sciences. 2017;6(8):1618-1621.
  29. Ikenganyia EE, Anikwe MAN, Ngwu OE. Influence of rhizobacteria inoculant application methods and phosphate fertilizer rates on dry matter accumulation, yield of bambara groundnut [*Vigna subterranea* (L.) Verdc] and soil total nitrogen content in a degraded ultisol in Southeast Nigeria. Agrotechnol. 2017;6:1-7.
  30. Sagvekar VV, Waghmode BD, Kamble AS. Effect of nitrogen and phosphorus management on productivity and profitability of groundnut (*Arachis hypogaea* L.). Indian Journal of Agronomy. 2017;62:338-40.
  31. Kumar BT, Malligawad LH, Halikatti SI, Hiremath SM. Effect of different ratios and levels of nitrogen and phosphorus fertilizers, and top dressing of nitrogen fertilizers on growth and yield of groundnut (*Arachis hypogaea* L.). Karnataka Journal of Agricultural Sciences. 2015;28(1):8-11.
  32. Yadav GS, Datta M, Babu S, Saha P, Singh R. Effect of sources and levels of phosphorus on productivity, economics, nutrient acquisition and phosphorus-use efficiency of groundnut (*Arachis hypogaea*) under hilly ecosystems of North-East India. Indian Journal of Agronomy. 2015;60(2): 307-311.
  33. Akbari KN, Ramdevputra MV, Sutaria GS, Vora VD, Padmani DR. Effect of organics and bio and inorganic fertilizer on groundnut yield and its residue effect on succeeding wheat crop. Legume Research- An International Journal. 2011; 34(1):45-47.
  34. Nwokwu GN. Influence of phosphorus and plant spacing on the growth and yield of groundnut (*Arachis hypogaea* L.). International Journal of Science and Research. 2011;3:97-103.
  35. John OS. Growth and yield response of groundnut (*Arachis hypogaea* L.) to plant densities and phosphorus on an ultisol in southeastern Nigeria. Libyan Agriculture Research Center Journal International. 2010;1:211-14.
  36. Naab JB, Prasad PVV, Boote KJ, Jones JW. Response of peanut to fungicide and phosphorus in on-station and on-farm tests in Ghana. Peanut Science. 2009;36:157-64.
  37. Kumar A, et al. Effect of phosphorus and sulphur on yield and nutrient uptake by groundnut in inceptisols. Asian Journal of Soil Science. 2008;3:139-41.
  38. Rajanikanth E, Subrahmanyam MVR, Rao JV. Effect of integrated nutrient management practices on growth and yield of rainfed groundnut (*Arachis hypogaea* L.) intercropped with guava (*Psidium guajava*). Journal of Oilseeds Research. 2008;25:157-60.
  39. Basu M, Bhadoria PBS, Mahapatra SC. Comparative effectiveness of different organic and industrial wastes on peanut: Plant growth, yield, oil content, protein content, mineral composition and hydration co-efficient of kernels. Archives of Agronomy and Soil Science. 2007;53(6): 645-658.
  40. Kausale SP, Shinde SB, Patel LK, Borse NS. Effect of integrated nutrient

- management on nodulation, dry matter accumulation and yield of summer groundnut at south Gujarat conditions. Legume Research- An International Journal. 2009;32(3):227-29.
41. Ranjit R, Dasog GS, Patil PL. Effect of lime and phosphorus levels on nutrient uptake by groundnut genotypes in acid soils of coastal agro ecosystem of Karnataka. Karnataka Journal of Agricultural Sciences. 2007;20:631-33.
  42. Gobarah ME, Mohamed MH, Tawfik MM. Effect of phosphorus fertilizer and foliar spraying with zinc on growth, yield and quality of groundnut under reclaimed sandy soils. Journal of Applied Sciences Research; 2006;2:491-96.
  43. Karmakar A, Karmakar S, Mukherjee S. Effect of lime and phosphorus levels on the pod, haulm and oil yield of the two groundnut cultivar. Karnataka Journal of Agricultural Sciences. 2005;20(3):627-630.
  44. Panwar AS, Singh NP. Effect of conjunctive use of phosphorus and bio-organics on growth and yield of groundnut (*Arachis hypogaea* L.) Indian Journal of Agronomy. 2003;48(3):214-216.
  45. Kachot NA, Malavia DD, Solanki RM, Sagarka BK. Integrated nutrient management in rainy season groundnut (*Arachis hypogaea* L.). Indian Journal of Agronomy. 2001;46(3):516-22.
  46. Kumaran S. Response of groundnut to organic manure, fertilizer levels, split application of phosphorus and gypsum application under irrigated condition. Research on crops. 2001;2(2):156-158.
  47. Majumdar B, Venkatesh MS, Lal B, Kumar K, Singh CS. Effect of phosphorus and zinc nutrition on groundnut in an acid hapludalf of Meghalaya. Annals of Agricultural Research. 2001;22:354-59.
  48. Musa AM, Singh L, Tame VT, Bubarai ML. Nitrogen, phosphorus and potassium uptake by some varieties of groundnut (*Arachis hypogaea* L.) as influenced by phosphorus application in Yola and Mubi, Adamawa State, Nigeria. IOSR Journal of Agriculture and Veterinary Science. 2017; 10(7):40-45.
  49. Chirwa M, Mrema JP, Mtakwa PW, Kaaya A, Lungu OI. Yield response of groundnut (*Arachis hypogaea* L.) to boron, calcium, nitrogen, phosphorus and potassium fertilizer application. International Journal of Soil Science. 2017;12(1):18-24.
  50. Mirvat EG, Magda HM, Tawfik MM. Effect of phosphorus fertilizer and foliar spraying with zinc on growth, yield and quality of groundnut under reclaimed sandy soils. Journal of Applied Sciences Research. 2006;2:491-96.
  51. Deka NC, Gogi PK, Pinku B, Dutta R. Effect of levels of lime and phosphorus on nutrient content and uptake of groundnut. Annals of Agricultural Sciences. 2001;22: 503-07.
  52. Vali GM, Singh S, Sruthi DSV, Hinduja N, Talasila V, Tiwari D. Effect of phosphorus and zinc on growth and yield of summer groundnut (*Arachis hypogaea* L.). The Bioscan. 2020;5(4):535-540.
  53. Vishwakarma AK, Brajendra, Pathak KA, Ramakrishna Y. Effect of different sources of nutrient application on productivity, nutrient uptake and economics of groundnut (*Arachis hypogaea* L.) in Kolasib district of Mizoram. Indian Journal of Soil Conservation. 2012;40(2):152-157.
  54. Patra PS, Bandyopadhyay S. Effect of phosphocompost on yield, phosphorus uptake pattern and net income of groundnut (*Arachis hypogaea* L.) in acid soil under terai region of west Bengal. Agricultural Science Digest. 2010;30(2): 120- 124.

© 2023 Kamal et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:  
The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/103501>