



Influence of Sowing Time on Varied Duration Redgram Genotypes in YSR Kadapa District

E. Aruna ^{a*} and K. Sunil Kumar ^b

^a Department of Agronomy, Agricultural Research Station, Utukur, YSR Kadapa District, ANGRAU, Andhra Pradesh, 516003, India.

^b Department of Entomology, Agricultural Research Station, Utukur, YSR Kadapa District, ANGRAU, Andhra Pradesh, 516003, India.

Authors' contributions

This work was carried out in collaboration between both authors. Author EA conducted the trial and took the data and did statistical analysis and wrote the paper. Author KSK helped in identifying the pests and guided chemicals for plant protection. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i183483

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/103660>

Original Research Article

Received: 26/05/2023
Accepted: 31/07/2023
Published: 10/08/2023

ABSTRACT

It is well known that redgram is a photo sensitive long duration crop of 180 days and taking up redgram will occupy the land for the year. But now because of the availability of photo insensitive short (120 days) and medium duration redgram genotypes (150 days), farmers can able to go for double cropping after harvesting of redgram with additional benefits. So there is a pressing need to evaluate the performance of these photo insensitive varieties of redgram compared to popularly grown long and medium duration varieties. Hence the experiment was conducted during kharif seasons of 2018 and 2019 at Agricultural Research Station, Utukur, YSR kadapa district of Andhra Pradesh to evaluate the performance of varied duration redgram genotypes under changing climate .The experiment was laid out in split plot design by taking three dates of sowing i.e. July I F.N, July II F.N and august I F.N in main plots and three varied duration i.e. long (180 days) (ICPL 87119),

*Corresponding author: E-mail: arunaettapu@gmail.com;

medium (LRG-52) (150 days) and short duration genotypes (120 days) (ICPL 20338) in sub-plots in three replications. The seeds were sown by dibbling in furrows at a depth of 5 cm at a spacing of 120 cm x 20 cm for long and medium duration genotypes and 30 cm x 10 cm for short duration genotypes. Plot wise data on growth parameters like plant height, number of branches per plant, number of pods per plant and yield parameters like pod length, number of seeds per pod, and 100 seed weight was taken and analysed by using SPSS software. The results indicated that Sowing of long or medium duration redgram varieties in July first or second fortnight depending on the onset of the monsoon is recommended for higher yield in redgram in YSR kadapa district of Andhra Pradesh whereas the performance of short duration varieties of redgram was poor in both the years making them unsuitable under rainfed conditions during kharif as rainfall affect the flowering which finally reduced the yield.

Keywords: Growth; time of sowing; varied duration redgram genotypes; yield.

1. INTRODUCTION

Redgram (*Cajanus cajan* L.) commonly known as pigeon pea, arhar or tur is a major pulse crop with its deep roots and drought resistance ability cultivated in around 3.86 M ha in India during *kharif* season under rainfed conditions. India ranks first in area and production of pigeon pea contributing to 70-80% of its world's production. Redgram with its virtue of having 22% protein in the seed, meeting the major share of protein needs of the vegetarians. Being a legume crop, Pigeon pea also enhances soil fertility through leaf litter and biological nitrogen fixation [1]. Though benefits of redgram to contribute to food security, nutrition, forage and income generation are indisputable, erratic climate patterns, long duration together with photo and thermo sensitivity of the crop restricted its expansion to wider latitudes and altitudes. So efforts are being continuously made for reducing the duration for redgram crop and as a result medium (155-170 days) and short duration (120 days) varieties were released in to the market. It was further reported that these short duration varieties due to photo and thermo insensitive nature very well fit in to the intensive cropping areas in all the seasons. Further, the reasons for low yields of pulses in India were mainly attributed to growing in poor/marginal lands, unawareness of recommended package of practices especially optimum sowing time, suitable variety, improper plant population, non-fertilization and improper plant protection measures due to improper diagnosis of the pest and diseases. Among all these, time of sowing largely determines the growth and yield of any crop including pulses. Hence there is a great need to determine the performance of varied duration genotypes of redgram under changing climate scenario during kharif in YSR kadapa district of Andhra Pradesh.

2. MATERIALS AND METHODS

The experiment was conducted during kharif seasons of 2018 and 2019 at Agricultural Research Station, Utukur, YSR kadapa district of Andhra Pradesh to evaluate the performance of varied duration redgram genotypes under changing climate. The experimental site was sandy loam and it was neutral in reaction with a P^H of 7.38, EC of 0.20 dS m^{-1} , low in organic carbon and available nitrogen, medium in available phosphorus and high in available potassium. The experiment was laid out in split plot design by taking three dates of sowing i.e. July I F.N, July II F.N and August I F.N in main plots and three varied duration i.e. long (180 days) (ICPL 87119), medium (LRG-52) (150 days) and short duration genotypes (120 days) (ICPL 20338) in sub-plots in three replications. By using tractor drawn cultivator the field was ploughed twice and levelled by using blade harrow. Healthy seeds of redgram varieties with good germination per cent (95%) were used for sowing purpose. The seeds were sown by dibbling in furrows at a depth of 5 cm at a spacing of 120 cm x 20 cm for long and medium duration genotypes and 30 cm x 10 cm for short duration genotypes and which were covered and compacted immediately for better germination after receipt of sufficient rainfall. Thinning was done at 15 days after sowing by retaining one healthy seedling $hill^{-1}$ and gap filling was done wherever necessary. The recommended dose of 20 and 50 kg N and P_2O_5 ha^{-1} was applied through urea and single super phosphate respectively at the time of sowing. Inter cultivation was done in inter rows and hand hoe weeding was done between the plants. Other crop husbandry practices like plant protection were followed as per ANGRAU recommendation for successful raising of the crop. Plot wise data on growth parameters like plant height, number

of branches per plant, number of pods per plant taken by selecting five plants randomly in each plot at the time of harvest and yield parameters like pod length, number of seeds per pod by collecting 10 pods from each plot and 100 seed weight was taken by taking seed from the yield obtained from each plot. The data used in the study are the mean values of replicated observations, which was tabulated and subjected to statistical analysis using analysis of variance (ANOVA) as applicable to split plot design by using SPSS software. A total rainfall of 147 and 766 mm was received in 13 and 48 days during 2018 and 2019 respectively against the average rainfall of 750 mm for the region. The weather data during the crop period is presented in Table 3.

3. RESULTS AND DISCUSSION

3.1 Effect of Time of Sowing

Plant height was not significantly influenced by the time of sowing in both the years. The number of branches per plant were also not significantly influenced by the time of sowing during 2018 where as during 2019, August I fortnight sown crop recorded lower number of branches compared to July sown crop. The time periods of days to 50% flowering and number of days to maturity were not significantly influenced by the time of sowing in both the years. During both the years, the number of pods per plant were more in early sown crop and with delay in sowing time the number of pods were decreased. Rain fall also significantly influenced the length of pods, the number of seeds per pod, 100 seed weight with lower values during 2018 which was drought year compared to 2019. But all these yield attributing characters like length of pods, number of seeds per pod, and 100 seed weight were not significantly influenced by the time of sowing. Consequently, there is significant impact of rainfall on the productivity of the redgram with lower yields during drought year of 2018 compared to 2019. Higher seed yield of redgram was recorded in July I fortnight sown crop which was on par with July II fortnight crop but was significantly superior to August I fortnight sown crop. The beneficial effect of early sowing on yield of redgram was also confirmed by Nagaraju et al. [2] and Kithan et al. [3]. The higher seed yield from first fortnight of July may be due to cumulative favourable effect of various growth and yield attributes. As early sowing get favourable environmental condition for proper growth and development which resulted in higher leaf area development and high biomass

accumulation which ultimately lead to significant improvement in seed yield. The crop sown on first fortnight of July produced more number of primary and secondary branches which ultimately resulted in production of more number of pods per plant which contributed towards higher seed yield. With delay in sowing, flowering was induced earlier resulting in less vegetative growth and earliness in maturity resulting in low seed yield. These results are in close conformity with those of [4] and Nene and Sheila, [5]. Pigeonpea suffers more when sowing is delayed [6]. Reddy et al. [7] reported that significant reduction occurs in branches/plant and dry weight/plant at harvest with delayed sowing than normal sowing. Ram et al. [8] revealed that the time to achieve harvest maturity was decreased in case of late sowing as compared to early sowing.

3.2 Effect of Genotype

Taller plants with a greater number of branches per plant was recorded in long and medium duration genotypes compared to short duration genotype. This might be due to genetic makeup of the plant. The long duration variety ICPL 87119 recorded higher number of pods per plant which was closely followed by medium duration variety LRG-52. Whereas the short duration variety (ICPL 20338) recorded significantly lower number of pods per plant. The pod length and number of seeds per pod were not significantly influenced by the varied duration genotypes. Higher 100 seed weight was reported with long and medium duration genotypes which was on par with each other but significantly lower seed weight was recorded with short duration genotype i.e ICPL 20338. Long duration genotype ICPL 87119 recorded higher seed yield followed by medium duration genotype (LRG- 52) but both are on par with each other. Lower seed yield was recorded with short duration genotype in both the years. Sahajadeva et al. [9] also reported higher yield and BC ratio with long duration varieties like LRG 41 but suggested to go for short duration varieties for late sown conditions under rain fed situation which can escape terminal moisture stress and under intensive cropping situations. The difference in grain yield of pigeonpea varieties was also reported by Prashanthi et al., [10]. Cultivars have a profound impact on the crop performance. Long duration genotypes produce maximum yield than early maturing genotypes, but they take more time to mature which may delay the sowing of succeeding crop such as wheat in a cropping system [11,12].

Table 1. Growth of varied duration redgram genotypes as influenced by time of sowing

Treatments	Plant height (cm)		Number of branches per plant		Days to 50% flowering		Days to maturity	
	2018	2019	2018	2019	2018	2019	2018	2019
Time of sowing								
July I F.N	145	139	13.4	12.43	135	140	180	186
July II F.N	140	134	13.0	12.42	130	136	175	178
August I FN	141	145	13.6	10.88	128	130	170	175
Sig	NS	NS	NS	*	NS	NS	NS	NS
P value	0.72	0.07	0.67	0.015	0.08	0.07	0.09	0.07
Varieties								
ICPL-87119	146	152	19.2	12.36	146	149	180	185
LRG-52	142	139	21.0	13.13	130	134	170	175
ICPL-20338	102	107	3.00	3.24	50	55	110	115
Sig	**	**	**	**	**	**	**	**
P value	0.00	0.00	0.00	0.000	0.00	0.00	0.00	0.00

Table 2. Yield attributes and yield of varied duration redgram genotypes as influenced by time of sowing

Treatments	No. of Pods/ plant		Pod length (cm)		Seeds/pod		100 seed wt (g)		Seed yield (kg ha ⁻¹)	
	2018	2019	2018	2019	2018	2019	2018	2019	2018	2019
Time of sowing										
July I F.N	86.08	109.75	3.03	4.79	2.11	3.67	7.33	10.05	452	832
July II F.N	81.99	93.77	3.13	4.99	2.16	3.87	7.22	10.22	406	814
August I FN	64.99	76.93	3.20	4.90	2.12	3.87	7.22	10.55	311	788
Sig	*	*	NS	NS	NS	NS	NS	NS	NS	**
P value	0.04	0.64	0.14	0.36	0.74	0.11	0.78	0.08	0.16	0.07
Varieties										
ICPL-87119	100.6	113.84	4.37	4.82	3.68	3.82	11.0	10.44	652	1010
LRG-52	94.5	107.91	4.98	5.01	3.79	3.78	10.7	10.83	516	930
ICPL-20338	20.3	25.04	3.98	4.00	3.60	3.80	7.38	7.55	405	495
Sig	**	**	NS	NS	NS	NS	**	**	**	**
P value	0.000	0.000	0.35	0.33	0.63	0.94	0.00	0.00	0.007	0.008

Table 3. Month wise Rainfall distribution during the crop period

Month	2018		2019		2018		2019	
	Rainfall (mm)	Rainy days	Rainfall (mm)	Rainy days	Maximum Temperature (°C)	Minimum Temperature (°C)	Maximum Temperature (°C)	Minimum Temperature (°C)
July	0	0	125.2	6	35.70	27.0	36.20	27.25
August	51.6	5	168.8	8	34.30	26.1	33.95	26.83
September	19.8	2	126.6	9	35.15	25.3	33.43	25.35
October	37.0	2	181.2	10	35.04	23.0	31.97	24.29
November	14.8	2	50.6	5	32.25	21.8	31.42	21.96
December	0	0	23.4	2	31.14	20.8	29.35	19.87
January	0	0	9.0	2	31.43	17.5	31.92	19.30
February	0	0	0	0	35.71	21.4	33.91	20.02

4. CONCLUSION

Delay in sowing significantly reduced the number of primary and secondary branches per plant, pods per plant, seeds per pod, test weight and seed yield as compared to early sown crop and found highest in crop sown on first fortnight of July followed by second fortnight of July and first fortnight of August. Sowing of long or medium duration redgram varieties in July first or second fortnight is recommended for higher yield in redgram in YSR kadapa district of Andhra Pradesh. The performance of short duration varieties in this region is not lucrative to farmers.

ACKNOWLEDGEMENTS

The authors are highly thankful to the Acharya. N.G. Ranga Agricultural University, Lam. Guntur for the financial help provided for conducting the research.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Udhaya ND, Vimalendren L, Letha KR, Sangamithra S, Kalaiyarasan V. A Review on biological advantage of Pigeonpea intercropping influenced by different cropping geometries. *Int. J. of Agric. Sciences Research*. 2015;5:103-112.
2. Nagaraju D, Kharbade SP, Sthool VA, Shaik AA, Jadhav JD, Balasubramaniyan R. Impact of yield attributes and yields of pigeon pea (*Cajanus cajan* L.) varieties under different sowing windows. *Int. J. of Environment and Climate Change*. 2022; 12(10):753–762.
3. Kithan L, Malini BS, Longchar A. Effect of planting dates on the performance of promising pigeonpea genotypes under NEZ. *Int. J. of Economic Plants*. 2020;7(1): 006-008.
4. Reddy GK, Reddy PM, Kumari PL, Krishna TG. Response of Pigeonpea varieties to time of sowing during rabi season. *Journal of Agriculture and Veterinary Sciences*. 2015;8(2):12-15.
5. Nene YL, Sheila VK. Pigeonpea: Geography and Importance. In: *The Pigeonpea*, Nene YL, Hall SH, Sheila VK. Eds. CAB International, Wellingford, U.K. 1990;1-14.
6. Padhi AK. Effect of sowing date and planting geometry on yield of redgram (*Cajanus cajan*) genotypes. *Indian Journal of Agronomy*. 1995;40(1):72-76.
7. Reddy MM, Padmaja B, Malathi S. Evaluation of pigeonpea genotypes for delayed Sowing in Telanganaregion of Andhra Pradesh under rainfed conditions. *Indian Journal of Dryl and Agricultural Research and Development*. 2012;27(2): 59-62.
8. Ram H, Singh G, Sekhon HS, Khanna V. Effect of sowing time on the performance of pigeonpea genotypes. *Journal of Food Legumes*. 2011;24:207-10.
9. Sahajadeva D, varaprasadarao ch, vinayalakshmi P, Prasada Rao GMV. Short duration varieties of pigeon pea perform better under late sown conditions in rainfed areas. *J. of Krishi Vignan*. 2021;9(2):225-228.
10. Prashanthi L, Vasanthi RP, Muneendra BABU. Evaluation of pigeonpea genotypes for rainfed conditions in the Southern Zone of Andhra Pradesh, India. *Int. Chickpea and Pigeonpea News I*. 2001;50-52.
11. Singh RS Performance of late duration pigeonpea varieties under delayed planting. *Indian Journal of Pulses Research*. 2016;19: 255-256.
12. Kumar A, Dhanoji MM, Meena MK. Phenology and productive performance of pigeonpea as influenced by date of sowing. *Journal of Pharmacognosy and Phytochemistry*. 2018;7(5):266-268.

© 2023 Aruna and Kumar; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/103660>