



# Effect of Nutrients and Mulches on Vegetative Growth and Yield Characters in Strawberry cv. Chandler

Charu Shekhawat <sup>a\*</sup>, Sanjay Kumar <sup>a</sup>, Pooja Meena <sup>a</sup>, Dharmendra Kumar <sup>a</sup>, Kehar Raj <sup>a</sup> and Deeksha Gautam <sup>b</sup>

<sup>a</sup> Department of Horticulture, School of Agricultural Sciences and Technology, Babasaheb Bhimrao Ambedkar (A Central University) Vidya Vihar, Rae Bareilly Road, Lucknow – 226025, U.P., India.

<sup>b</sup> Department of Fruit Science and Horticulture Technology, Odisha University of Agriculture and Technology, Bhubaneswar-751003, Odisha, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJPSS/2023/v35i203861

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/104101>

Original Research Article

Received: 03/06/2023

Accepted: 05/08/2023

Published: 02/10/2023

## ABSTRACT

A field experiment was conducted at the Horticulture Research Farm-1, of the Department of Horticulture, School of Agricultural Sciences and technology, Babasaheb Bhimrao Ambedkar University (A Central University), Vidya- Vihar, Rae Bareilly Road, Lucknow to evaluate the significance of mulching and nutrients on the performances of Strawberry (*Fragaria x ananassa*) with respect to the vegetative and yield character's during 2019-20. The experiment was laid out in RBD having three replications. The result revealed significantly highest plant height (24.82 cm), number of leaves (20.72cm), length of leaf (9.85cm), length of petiole (8.98 cm), no. of pickings

\*Corresponding author: E-mail: Charushekhawat95@gmail.com;

(22.86), no. of fruits per plant (24.58), length of fruit (42.88 mm), width of fruit (27.98 mm), weight of fruit (21.87g) found in treatment T-14 (Black polythene vermicompost 50% + NPK 50%) followed by the treatment T-13 (Black polythene FYM 50% + NPK 50%) at 90 Days after planting. However, treatment T<sub>1</sub> (control) showed poor performance with respect to all vegetative and yield parameters. Hence, the result led to the conclusion that adoption of Black polythene with vermicompost 50% + NPK 50% in Strawberry enhances all growth and yield attributing parameters and black polythene with FYM 50% + NPK 50% showed second-best result.

**Keywords:** Strawberry; nutrients; mulches; growth and yield.

## 1. INTRODUCTION

“The Strawberry is a most important man made horticultural fruit crop. The modern cultivated octaploid Strawberry (*Fragaria X ananassa* Duch.) originated in France. It belongs to the family Rosaceae. It is one of the delicious and nutritious among soft fruit with wider adoptability. This crop is chiefly grown in the temperate world. However, it can be grown under sub-tropical climate and even at high altitude of tropical climate. Strawberry can be grown wide range of soil from heavy clays to light sand and gravels. However, strawberry plant grows well in sandy loam soil with pH of 5.5- 6.5 [1]. “The strawberry requires an optimum day temperature of 22°C & night temperature of 7°C to 13°C for maximum growth and development. In cold climate frost damage as well as winter injury greatly reduced yield of strawberry. Frost may damage center of open flower causing the characteristics ‘Black eye’ disease. Frost free site make strawberry cultivation comparatively easy, it is also grown under the green house and covered plastic. Flower before opening and after set are slightly more resistant to frost damage. Proper nutrition is very important for its profitable cultivation. ‘Albinism’ is a physiological disorder probably caused by certain climatic condition and extremes in nutrition. Being a quick growing crop, it is also suitable for the kitchen garden. The fruit of strawberry is a complete fruit with 98% edible portion. The first plant of the cultivated strawberry (*Fragaria x ananassa*) has originated in France and the two American diploids *F.chiloensis* and *F.virginiana* considered in progenitors” [2].

The practice of applying mulch is as old agriculture itself[3]. Black polyethylene, sawdust, wheat straw, paddy straw, grasses, sugarcane trash, litters, crop residues etc. are common mulching materials used for covering plant basin. “Both organic and inorganic mulches are used in strawberry cultivation. Mulching protects fruits from soil contact, reduces the number of dirty fruits and prevents the occurrence of fruit rot.

Mulching reduces the water evaporation by interfering the radiation falling on the soil surface ensuring better moisture regime and more availability of moisture” [4,5].

## 2. MATERIALS AND METHODS

The experiment was conducted at Horticulture Research Farm – I of Department of Horticulture, School of Agricultural Sciences and Technology, Babasaheb Bhimrao Ambedkar University, Lucknow, India; during the academic year of 2019-20 it is geographically situated in the subtropical tract of Central U.P. at 80°55’ East longitude and 26°56’ North latitude. The Horticulture Research Farm is situated at 10 Km away from the Lucknow Railway Station towards South-East in Lucknow Rae Bareli Road, Vidya-Vihar, near South City, Lucknow. “This region received an average annual rainfall of 650-750 mm which is distributed over a period of more than 100 days with peak period during July-August. It is also receiving scattered show during winter months. In general, the temperature ranges from 3.5°C to 45°C. The coldest month is January while, the maximum temperature is observed during May and June. The relative humidity (RH) is 50-77% in different season of the year. Lucknow is characterized by sub-tropical climate with hot, dry summer and cold winter. The soil of experimental farm was saline with soil pH less than 8.2, Electrical conductivity more than 4.0 and sodium exchangeable percentage less than 15.0”[2].

### 2.1 Experimental Materials

The runners of Chandler variety of strawberry were brought from the Central Institute of Temperate Horticulture, Kashmir (J&K) in the month of October, 2019. The runners were kept for two days in shade for hardening before transplanting in well-prepared beds under open field condition plots which were distributed randomly in three replications. Standard cultural practices were followed during the period of the experiment for maintaining the runners.

## 2.2 Planting System and Crop Management

An area of 18 m x 7.80 m size was divided into 42 plots having the size of 1.8 m x 1.2 m and arranged in the three replications of 14 plots. The experiment was laid out in R. B. D. under 14 treatments with spacing of 45 cm row to row and 30 cm plant to plant. The field was prepared with basal dose of N.P.K. fertilizer as indicated earlier. Sixteen plants, in each bed were transplanted and watered uniformly. Water was applied in channels between two beds. Mulching was done two days after transplanting black polythene (200 gauge) was used for covering the soil surface in the bed and also used white polythene (200 gauge) to cover the plots.

**Table 1. The details of treatments used in the experiments are given below**

T <sub>1</sub>	White Polythene
T <sub>2</sub>	White Polythene + FYM 100%
T <sub>3</sub>	White Polythene + vermi-compost 100%
T <sub>4</sub>	White Polythene + NPK 100%
T <sub>5</sub>	White polythene FYM 50%+ Vermi-compost 50%
T <sub>6</sub>	White Polythene FYM 50% + NPK 50%
T <sub>7</sub>	White polythene Vermi-compost 50% + NPK 50%
T <sub>8</sub>	Black Polythene
T <sub>9</sub>	Black polythene + FYM 100%
T <sub>10</sub>	Black Polythene + Vermi-compost 100%
T <sub>11</sub>	Black polythene +NPK 100%
T <sub>12</sub>	Black Polythene FYM 50%+ Vermi-compost 50%
T <sub>13</sub>	Black Polythene FYM 50%+ NPK 50%
T <sub>14</sub>	Black Polythene Vermi-compost 50%+ NPK 50%

## 3. RESULTS AND DISCUSSION

Among these mulching treatments, black polythene mulch showed significant superiority in reducing weed population over rest of the treatments. This effect may be due to smothering effect and causing physical barrier to photosynthetic activity imparted by polythene mulches due to which plant performance may have improved by better nutrient uptake by the plants. Which ultimately enhances the vegetative growth of the strawberry In case of nutrient vermin-compost and FYM enhances the growth and yield possibly due to the modification in the soil environment viz., moderating soil temperature during early Growth of the crop which coincides with hot dry month that

preserves soil moisture also contributed to increase the production of the more vegetative Growth and yield.

### 3.1 Growth Character

The plant height of various treatment combination has been represented in Table-2, which was varied from (15.19 cm) to (24.82 cm). In which, the maximum plant height was recorded in treatment T-14 (Black polythene + vermi-compost 50% + NPK 50%) showed value (24.82 cm), followed by the treatment T-13 (Black polythene + FYM 50% + NPK 50%) showed value (22.68 cm). However, the minimum plant height was recorded in treatment T<sub>1</sub> (control) showed value (15.19 cm). Christopher Lourduraj et al. 1996 [6] found that plastic mulch resulted in significantly higher plant height as compared to organic mulches in tomato. Mulching with black polythene enhanced growth of strawberry as compared with unmatched one [7].

Findings of number of leaves per plant as influenced by different treatment combinations and the relevant data are presented in Table-2, which recorded from (14.45 cm) to (20.72 cm) at the 90 days after planting. Minimum number of leaves was found in treatment T<sub>1</sub> control (14.45 cm). The maximum number of leaves was recorded in treatment T-14 (20.72 cm) (Black polythene + vermi-compost 50% + NPK 50%) followed by treatment T-13 Black polythene + FYM 50% +NPK 50% (19.56cm). Chattopadhyay and Sarad, Gurung[8] reported that mulching with black polythene resulted in best vegetative growth, pseudo stem height, girth, leaf, number and sucker production in banana cv. Giant Cavendish.

The data regarding length of leaf under different treatment combination presented in table number 2, the maximum length of leaf was assessed in the treatment T-14 (Black polythene + vermi-compost 50% + NPK 50%) 9.85 cm followed by the length of leaf occurs in treatment T-13 (Black polythene + FYM 50% +NPK 50%) 9.65 cm. The minimum length of leaf found in the treatment T<sub>1</sub> Control (7.45) cm. Gaiwatet al. [9] reported that grass mulch increased vegetative growth in term of number of shoot/ branch and leaf area as compared to polythene mulch in Nagpur Mandarin.

It is clear data presented in the table number 2. Length of petiole varies from 3.98 cm to 8.98 cm. Maximum length of petiole observed in the

**Table 2. Variation in growth parameters by application of different treatments.**

Treatments	Plant height (cm)			No. of Leaves			Petiole length			Length of leaf		
	30	60	90	30	60	90	30	60	90	30	60	90
T <sub>1</sub>	3.53	6.65	15.19	3.25	10.48	14.45	2.63	3.03	3.98	3.52	4.35	7.45
T <sub>2</sub>	3.73	6.88	15.69	3.72	10.84	14.66	3.29	4.26	5.15	3.68	4.55	7.67
T <sub>3</sub>	3.98	7.68	16.86	3.92	10.93	14.82	3.35	4.36	5.38	3.76	4.65	7.72
T <sub>4</sub>	4.15	7.82	17.23	4.22	11.38	15.68	3.41	4.45	5.41	3.96	4.75	7.85
T <sub>5</sub>	4.54	7.98	17.73	4.29	11.74	15.74	3.58	4.52	5.77	4.25	5.32	8.92
T <sub>6</sub>	4.73	8.36	18.34	4.41	11.85	16.49	3.92	5.22	6.26	4.42	5.48	8.63
T <sub>7</sub>	4.94	8.59	18.69	4.82	12.25	16.73	4.12	5.37	6.38	4.68	5.58	8.75
T <sub>8</sub>	5.13	8.85	18.89	5.18	12.42	17.25	4.19	5.42	6.43	4.75	5.66	8.97
T <sub>9</sub>	5.45	8.92	19.46	5.56	12.77	17.54	4.28	5.52	6.54	4.82	5.75	8.52
T <sub>10</sub>	5.66	9.18	19.82	5.92	13.52	18.35	4.35	5.63	6.68	4.96	5.95	8.75
T <sub>11</sub>	6.42	9.55	20.48	6.35	13.67	18.75	5.15	5.76	7.06	5.28	6.42	9.22
T <sub>12</sub>	6.58	9.92	20.85	6.68	14.68	19.25	5.39	6.05	7.21	5.34	6.58	9.48
T <sub>13</sub>	7.14	10.78	22.68	6.87	14.86	19.56	5.55	6.32	8.78	5.76	6.72	9.65
T <sub>14</sub>	7.29	10.82	24.82	7.18	15.75	20.72	5.78	6.53	8.98	5.96	6.92	9.85
S.E m(±)	0.20	0.35	0.55	0.17	0.38	0.51	0.14	0.2	0.26	0.15	0.20	0.31
C.D. (P=0.05)	0.60	1.05	1.62	0.50	1.12	1.49	0.49	0.59	0.78	0.45	0.60	0.90

**Table 3. Variation in yield parameters by application of different treatments**

Treatments	No. of pickings	Length of fruit (mm)	Width of fruit (mm)	Yield/plant (g)	Weight of fruit (g)	No. of fruit per plant
T <sub>1</sub>	13.22	28.93	21.48	233.47	15.24	15.32
T <sub>2</sub>	13.54	30.92	21.52	244.72	15.45	15.84
T <sub>3</sub>	14.22	32.62	21.82	267.76	16.14	16.59
T <sub>4</sub>	14.82	33.15	22.67	277.70	16.55	16.78
T <sub>5</sub>	15.55	33.38	22.96	285.26	16.80	16.98
T <sub>6</sub>	16.78	34.25	23.18	300.05	17.62	17.65
T <sub>7</sub>	16.98	35.44	23.28	328.56	17.77	18.49
T <sub>8</sub>	17.68	35.84	23.85	341.31	18.04	18.92
T <sub>9</sub>	18.24	36.42	24.55	355.71	18.45	19.28
T <sub>10</sub>	18.88	36.65	24.73	372.94	18.76	19.88
T <sub>11</sub>	19.65	37.59	25.43	399.74	19.33	20.68
T <sub>12</sub>	20.54	38.15	25.84	422.14	20.15	20.95
T <sub>13</sub>	20.78	38.32	26.88	471.83	20.85	22.63
T <sub>14</sub>	22.86	42.88	27.98	537.56	21.87	24.58
S.E m(±)	0.68	1.27	0.77	10.84	0.54	0.77
C.D. (P=0.05)	1.99	3.72	2.26	31.70	1.60	2.27

treatment T-14 (Black polythene + vermi-compost 50% + NPK 50%) 8.98 cm followed by length of petiole treatment T-13 (8.78) cm. Lowest length of petiole regarded in treatment T<sub>1</sub> Control (3.98) cm. Byun et al. [10] reported the black polythene mulch increased tree height, leaf area, leaf petiole length and number of branches in young jujube tree.

### 3.2 Yield Character

The data presented in Table-3 indicates that maximum No. of pickings (22.86) were noted in treatment T-14 at 90days after planting. Where treatment T-13 which was significantly followed by the black polythene + FYM 50% + NPK 50% application observed data calculated (20.78) at 90 days after planting . However, the minimum harvesting data calculated (13.22) under controlled plants treatment at 90 days after planting. Plastic mulch increased yield of strawberry and keep fruit clean [11].

The mean values presented in Table-3 clearly indicate that the maximum length of fruit was recorded under (Black polythene + vermi-compost 50% + NPK 50%) (42.88 mm) at treatment T-14 followed by treatment T-13 (38.32 mm) with effect of (Black polythene + FYM 50% +NPK 50%) However, the minimum fruit length (28.93 mm) was observed under controlled T-1.

Findings of width of fruit clearly indicate that the maximum width of fruits were recorded under Black polythene + vermi-compost 50% + NPK 50% (27.98 mm) at treatment T-14. Followed by T-13 (26.88 mm) with (Black polythene + FYM 50% +NPK 50%) However, the minimum fruit width (21.48 mm) was observed under controlled treatment T<sub>1</sub>.

Data regarding yield per plant presented under different treatment combination in Table-3 indicates that the maximum yield per plant were recorded under Black polythene + vermi-compost 50% + NPK 50% (537.56 g) treatment T-14 followed by T-13 (471.83) with Black polythene + FYM 50% +NPK 50% However, the minimum yield per plant (233.47) was observed under controlled treatment T<sub>1</sub>. Ali and Gaur [12] reported the mulching significantly increased no of fruits, fruit weight and fruit yield. Maximum number of fruits (13.75), Fruit weight (7.9g) & fruit yield (109.68/plant) were recorded in black polythene mulch plants, while minimum in control.

The mean values presented in Table-3 clearly indicate that the maximum weight of fruit was

recorded under Black polythene + vermi-compost 50% + NPK 50% (21.87g) at T-14. Followed by T-13 (20.85 g) with effect of Black polythene + FYM 50% +NPK 50% However, the minimum weight of fruit (15.24g) was observed under controlled T<sub>1</sub>. Kasperbauer [13] observed that most commonly used plastic mulch color is black in strawberry and also reported that the yield per plant and size per berry were greater over the red plastic mulch.

Findings of no. of fruit clearly indicate that the maximum no. of fruits were recorded under Black polythene + vermi-compost 50% + NPK 50% (24.58) at treatment T-14. Followed by T-13 (22.63) with (Black polythene + FYM 50% +NPK 50%) However, the minimum no. of fruit (15.32) was observed under controlled treatment T<sub>1</sub>.

### 4. CONCLUSION

From the investigation with nutrient and mulches on strawberry cv. Chandler it can safely be concluded that T-14 Black polythene + vermi-compost 50% + NPK 50% significantly increase the growth and yield characters i.e. plant height, number of leaves per plant, length of leaves, length of petioles and total number of fruits, content of fruits, whereas, T-14 produced maximum weight of fruits. It can be recommended to the strawberry growers for obtaining a better quality of winter season strawberry fruits in Lucknow condition.

### COMPETING INTERESTS

Authors have declared that no competing interests exist.

### REFERENCES

1. Anonymous. Wealth of India, CSIR, New Delhi, India. 1956;IV:57-60.
2. Shukla KK, Verma RS, Pandey L, Shukla JK, Singh SN, Prakash S. Effect of different types of mulches on vegetative growth characters in strawberry (*Fragaria x ananassa*Duch.) cv. Chandler. The Pharma Innovation Journal. 2021;10(12):2641-4.
3. Jacks GV, Brind WD, Smith, R. Mulching. Tech Commun. Bur Soil Sci. Rathamsted. 1955;49.
4. Gupta JP. Evaporation from a sandy soil under mulches. Ann. Arid Zone. 1978; 17(3):287-290.

5. Gupta JA. Effect of mulch on hydro-thermal environment of soil and crop production in arid zone of western Rajasthan. Ann. Arid Zone. 1985;24(2):131-140.
6. Cristopher, Lourduraj A, Shreenarayanan VV, Rajendran R, Ravi V, Padmini K, Pandurajan T. Effect of plastic mulching on tomato yield and economics. South Indian Hort. 1996;44(5-6):134-142.
7. Srivastava RP, Agrawal NC. A note on the mulching of strawberry. Indian J. Hort. 1965;22:211-213.
8. Chattopadyay PK, Gurung, Sarad. Effect of soil covers on growth flowering and yield of banana cv. GaintGoverner. Prog. Hort. 1993;25(3/4):181-183.
9. Gaiwat SC, Ingle HV, Panchbhai DM. A note on the effect of different type of mulches on growth, yield and quality of Nagpur Mandarin. The Orisa J. Hort. 2004;30(3):137-138.
10. ByunJK, Do JH, Chang KN. The effect of black polythene and rice straw mulches on growth of young jujube (*Z. jujube* Miller) plants. Korean Soc. Hort. Sci.1989;7(1):138-139.
11. Martin NLF, Westphalen SL. Effect of plastic mulches on yield of two strawberry (*Fragaria x ananassa*Duch.) cultivars and on soil temperature. Agronomia Sulriograndense. 1983;19(2):67-88.
12. Ali A, Gaur GS. Effect of various mulches on yield and quality of strawberry (*Fragaria x ananassa*Duch.) cv. Sweet Charlie. International Seminar on Rec. Trend Hi-Tech. Horticulture & PHT, Kanpur. February2004;4-6:121-122.
13. Kasperbauer MJ. Strawberry yield over red verses black plastic mulch. Crop Sci. 2000;40(1):171-174.

© 2023 Shekhawat et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/104101>