



Unveiling the Proximate Composition of Pigmented and Non-Pigmented Rice Bran

Gosangi Avinash^{a*} and Neerja Sharma^b

^a Department of Biochemistry, Punjab Agricultural University, Ludhiana, Punjab, India.

^b Department of Plant Breeding and Genetics, Punjab Agricultural University, Ludhiana, Punjab, India.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i214020

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/100929>

Original Research Article

Received: 02/04/2023

Accepted: 08/06/2023

Published: 27/10/2023

ABSTRACT

This present study aimed to evaluate the proximate composition of pigmented and non-pigmented rice. We have evaluated the proximate components of 16 rice bran genotypes, 9 pigmented rice bran, and 7 NP rice bran, grown and cultivated in the research fields. The proximate composition (moisture, ash, crude fat, crude proteins, crude fiber) were determined using proximate analysis methods. Raising awareness regarding rice's bran nutritional quality and health benefits is important. The results shown that black rice genotypes exhibited high range of almost all the parameters. Crude protein in black rice genotypes, ranged from 18.02-21.01%, crude fat content in black rice genotypes, ranged from 19.83-21.73% and ash content in the black rice genotypes, ranged from 14.08-16.80%, ash in black rice genotypes, ranged from 11.28-12.35%, crude fibre. The result exhibited a significant difference in the content of proximate composition. The study concludes that the significant distinctiveness and difference in nutritional and is use full in rice bran products. The results suggest that the black rice genotypes have better nutrition as compared to NP rice genotypes. The results showed great variations in each parameter among different rice bran varieties.

*Corresponding author: E-mail: avinashgosangi@gmail.com;

Keywords: Pigmented rice; NP-non pigmented rice; crude protein; ash.

1. INTRODUCTION

Rice (*Oryza sativa* L.) is a significant and staple food among all the major cereal crops consumed by over 50 % of the total world's population. It is fundamental to Asian societies not just as a food source, but also in an array of different social, financial and religious activities [1] (Hedge et al 2013). Different layers of rice have known to have varying amounts of nutrients [2]. In contrast to the aleurone layer, which is deficient in vital nutrients, the bran layer is rich in dietary fibre, minerals, and vitamin B complex [3,4]. When compared to germ and bran, the rice endosperm consists of low concentrations of proteins, lipids, and fibre but an increased amount of carbohydrates [5,6]. Most of the bioactive compounds are present in bran and germ fractions. Tocopherols (vitamin E), phenolics, carotenoids (lutein, β -cryptoxanthin, zeaxanthin, and β -carotene), as well as compounds containing nitrogen and organo-sulphur are among the phytochemicals found in whole grains of rice [7,8]. Most of the phytochemicals are highly concentrated in the outer layer (pericarp) and testa or the bran part of the rice kernel, which results in rice with different colours *i.e.*, purple, black and red.

Rice is naturally brown after harvesting, but once the nutrient-rich outer layer of bran is removed, it is white in color. Red, black, and purple rice, all feature unique pigmentation in the bran.

Rice bran contains 12-20 % protein of excellent quality which can be a source of dietary protein in human nutrition [9]. Among cereals, rice bran protein has the highest nutritional value due to its high content of lysine and threonine (limiting amino acids), which are generally low in other cereals. Rice bran protein (RBP) contains approximately 3-4% lysine, an essential amino acid, which is highest among other cereal proteins [10]. Protein digestibility of rice bran protein is greater than 90% and the protein efficiency ratio (PER) values of rice bran concentrate is relatively high, around 2.0 to 2.5, whereas casein is 2.5 [11]. The RBP can be used as a source of protein in infant food formula and restricted food for children with allergies, due to its hypoallergenicity. After hydrolysis of RBP with protease enzyme, small peptides with higher antioxidative activity were obtained by Adebisi et al. [12].

2. MATERIALS AND METHODS

2.1 Collection of Samples

Rice crop was harvested at maturity. After drying the paddy to moisture content of 14%, cleaned it thoroughly to get rid from impurities. Cleaned paddy was stored in cloth bags at ambient temperature for two months. Paddy was dehulled on Satake Rice Machine (Kitashiba electronics Co.ltd., Japan) to obtain brown rice and then milled on Satake Test Mill (Kitashiba electronics Co.ltd., Japan) to obtain rice bran. The bran which obtained after milling was sieved through Standard Test Sieve (425 MICs) and analyzed for various biochemical parameters by standard procedures reported in literature.

2.2 Moisture Content (AACC 2000)

Weighed rice fractions (10 g) were dried in a hot air oven at $130 \pm 1^\circ\text{C}$ for one hour and the content of moisture was calculated in percent from the loss in weight of the sample

2.3 Crude Protein (Micro Kjeldahl Method AOAC 2000)

Reagents

- i. Digestion mixture ($\text{CuSO}_4: \text{K}_2\text{SO}_4, 1:9$)
- ii. Concentrated sulphuric acid (H_2SO_4)
- iii. 1 M sodium hydroxide (NaOH)
- iv. 0.01N hydrochloric acid (HCl)
- v. 4% Boric acid (H_3BO_3)
- vi. Indicator: 0.5 g bromocresol green and 0.1g methyl red dissolve in 100 ml of ethanol (95%)

Procedure

Took 100 mg rice powder in digestion tubes and added 10 ml concentrated H_2SO_4 along with 5 g digestion mixture. Digestion was done till a clear solution was obtained. Digestion tubes were allowed to cool to room temperature and the content was transferred to 100 ml volumetric flask, quantitatively and made to volume. An aliquot of 20 ml was distilled using Kjeldahl distillation unit (FOSS Kjeltac 2100) after the addition of 50 ml 1 N NaOH. Distillate was collected in 25 ml 4% H_3BO_3 and titration was done against 0.01 N HCl till grey end point. Percent nitrogen content (%N) was calculated as follows:

Percent nitrogen (% N) = $\frac{(X - \text{Blank}) \times 0.00014}{Y} \times 100$

Crude protein content = Percent nitrogen (% N) x 5.95

Where X= Titre value (ml)
Y = Sample weight (g)

2.4 Crude Fat (AACC 2000)

Reagent

Petroleum ether.

Procedure

SOCS plus apparatus (Pelican Equipment's, Chennai, India) was used for the extraction of crude fat in the samples. Thimble containing 5 g rice powder sample was placed in the pre-weighed extraction beaker containing 70 ml petroleum ether. Beakers were loaded in the chambers of the apparatus. Extraction was carried out for 1 hour 30 minutes after which the extraction beakers were allowed to cool. The extraction beakers were oven dried (60- 65°C). The solvent was recovered. The crude fat was calculated as percentage from the increase in weight of the extraction beakers.

2.5 Ash Content (AACC 2000)

Procedure

Rice powder (5g) was taken in pre- weighed crucible and charred completely on a hot plate. The crucibles were placed in the muffle furnace for 5 hours at 550°C. The crucibles were allowed to cool to room temperature and were weighed accurately. Content of ash was calculated from the increase in the weight of the crucibles and expressed in percentage.

2.6 Estimation of Crude Fibre (AACC 2000)

Reagents

- i. 0.12 N H₂SO₄
- ii. 1 N NaOH

Procedure

The sample (2 g) was boiled with 250 ml H₂SO₄ solution for 30 min. Filtered it through sintered

glass crucibles and washed thoroughly with hot water. The residue was boiled again with 250 ml 1 N NaOH for 30 min followed by filtration and washing with hot water until a clear filtrate was obtained. The residue was dried and ashed in muffle furnace at 600° C. The crude fibre content was calculated from the loss in weight of samples and expressed as a percentage.

3. RESULTS AND DISCUSSION

3.1 Moisture Content

Moisture content in bran ranged similar in all the three-color groups. In bran fraction, in black rice genotypes, the moisture content ranged from 7.41-9.59%, in red rice genotypes, it ranged from 7.35-10.49%, and in NP rice genotypes it ranged from 6.87-8.71%. The moisture content of all three colour groups were lower than 14%, Champagne et al. [13] stated that the 14% moisture content was considered safe for storing grains to avoid damage and deterioration of seed quality. The difference in the moisture content might be due to climatic conditions where the crop is cultivated and the difference in the genetic makeup of an individual genotype. Ebuehi et al. [14] reported that the moisture content plays a significant role in determining the shelf life of foods. Xheng and Lan [15] reported that moisture influences the milling characteristics and the taste of cooked rice (Table 1).

3.2 Crude Protein

Crude protein content in bran fraction, in black rice genotypes, ranged from 18.02-21.01%, in red rice genotypes, it ranged from 16.21-20.45%, and in NP rice genotypes it ranged from 15.78-21.06% (Table 1). The average crude protein content was found to be higher in pigmented rice groups compared to NP rices. Simonelli et al. [16] stated that proteins play a crucial role in cooked rice texture because they form a complex with starch that restricts the starch granule swelling. Suwannaporn et al. [17] reported that high protein rice is less sticky and has a harder texture of cooked rice. Kennedy et al. [5] reported that the second major component next to starch are proteins which influence the eating quality and the nutritional quality of rice. The dietary supply of rice per person per day is 207.9 g, which provides about 24.1% of the required dietary protein.

3.3 Crude Fat

Crude fat refers to the crude mixture of fat-soluble material present in a sample and is also known as the ether extract or the free lipid content. The lipid materials may include free fatty acids, fat-soluble vitamins, monoglycerides, diglycerides, triglycerides, phospholipids, steroids, carotene pigments, chlorophylls, etc. Juliano et al. (1986) and Tanaka et al. (1978) Starch lipids present in rice are composed of monoacyl lipids (fatty acids and lysophosphatides) complexed with amylose [18]. In the present study crude fat content in bran

fraction, in black rice genotypes, ranged from 19.83-21.73%, in red rice genotypes, it ranged from 15.25-20.97%, and in NP rice genotypes it ranged from 17.38-22.39%. Among three colours, the mean crude fat content was found to be the highest in black rices while red rices and NP rice genotypes exhibited overlapping content. It has been reported that the lipids or fats in rice are mainly confined to the rice bran (20%, dry basis) as lipid bodies in the aleurone layer and bran. Priya et al. [19] reported that the amount of crude fat present in pigmented rices was high in red rices.

Table 1. Proximate composition of the bran of pigmented and non-pigmented rice genotypes

Genotype	Moisture (%)	Ash (%)	Crude Protein (%)	Crude Fat (%)	Crude Fibre (%)
Black rices					
FCP	9.59 ± 0.12 ^a	15.41 ± 0.29 ^b	20.1 ± 0.46 ^a	19.96 ± 0.23 ^b	12.35 ± 0.04 ^a
FCAP	8.36 ± 0.03 ^b	14.08 ± 0.26 ^c	18.02 ± 0.4 ^b	19.83 ± 0.11 ^b	11.28 ± 0.02 ^c
CP	7.41 ± 0.06 ^c	16.80 ± 0.16 ^a	21.01 ± 0.34 ^a	21.73 ± 0.14 ^a	11.75 ± 0.01 ^b
Mean	8.45 ^A	15.43 ^A	19.71 ^A	20.51 ^A	11.79 ^A
Range	7.41-9.59	14.08-16.80	18.02-21.01	19.83-21.73	11.28-12.35
C.V.	1.65	2.73	3.54	1.43	0.38
P-Value	<0.001	<0.001	<0.001	<0.001	<0.001
Red rices					
GSL	6.82 ± 0.13 ^e	12.93 ± 0.06 ^b	19.23 ± 0.07 ^{bc}	20.97 ± 0.16 ^a	8.33 ± 0.02 ^c
DMI	8.46 ± 0.01 ^c	14.75 ± 0.01 ^a	20.45 ± 0.47 ^a	17.62 ± 0.12 ^b	9.04 ± 0.02 ^b
KMA	9.65 ± 0 ^b	10.72 ± 0.04 ^c	19.87 ± 0.1 ^{ab}	15.25 ± 0.08 ^d	7.0 ± 0.04 ^d
BKG	10.49 ± 0.11 ^a	13.06 ± 0.25 ^b	16.21 ± 0.28 ^d	16.76 ± 0.07 ^c	10.72 ± 0.06 ^a
MSTY	9.65 ± 0.21 ^b	12.87 ± 0.29 ^b	18.47 ± 0.11 ^c	15.49 ± 0.06 ^d	6.94 ± 0.02 ^d
YKM	7.35 ± 0.07 ^d	14.35 ± 0.07 ^a	20.36 ± 0.47 ^a	17.63 ± 0.04 ^b	5.93 ± 0.08 ^e
Mean	8.73 ^A	13.11 ^B	19.10 ^A	17.29 ^B	7.99 ^B
Range	6.82-10.49	10.72-14.75	16.21-20.45	15.25-20.97	5.93-10.72
C.V.	2.25	2.13	2.72	0.97	0.97
P-Value	<0.001	<0.001	<0.001	<0.001	<0.001
Non-pigmented rices					
PR 114	8.46 ± 0.06 ^b	10.45 ± 0 ^e	18.29 ± 0.34 ^c	18.61 ± 0.08 ^{cd}	6.4 ± 0.03 ^e
PR 121	6.92 ± 0.15 ^d	12.35 ± 0.01 ^c	21.06 ± 0.37 ^a	22.39 ± 0.14 ^a	5.91 ± 0.02 ^f
PR 126	8.91 ± 0.04 ^a	10.69 ± 0.11 ^e	16.27 ± 0.32 ^{de}	19.84 ± 0.18 ^b	7.66 ± 0.03 ^d
BAS 370	7.82 ± 0.02 ^c	13.76 ± 0.07 ^b	19.36 ± 0.19 ^b	18.72 ± 0 ^{cd}	7.91 ± 0 ^c
BAS 386	8.71 ± 0.11 ^a	15.4 ± 0.06 ^a	15.78 ± 0.03 ^e	19.02 ± 0 ^c	8.17 ± 0.16 ^b
Pb Bas7	8.02 ± 0.08 ^c	11.27 ± 0.19 ^d	17.02 ± 0.39 ^d	18.28 ± 0.18 ^d	8.33 ± 0.05 ^b
PB 1121	6.87 ± 0.05 ^d	10.49 ± 0.04 ^e	16.02 ± 0.19 ^e	17.38 ± 0.3 ^e	10.24 ± 0.1 ^a
Mean	7.96 ^B	12.06 ^B	17.69 ^B	19.18 ^{AB}	7.80 ^B
Range	6.87-8.91	10.45-15.40	15.78-21.06	17.38-22.39	5.91-10.24
C.V.	1.78	1.31	2.83	1.45	1.69
P-Value	<0.001	<0.001	<0.001	<0.001	<0.001
Average (Overall)	8.34	13.09	18.60	18.72	8.62
Range (Overall)	6.82-10.49	10.45-16.80	15.78-21.06	15.25-22.39	5.91-12.35

Each value represents the mean ± SD of three independent replications

3.4 Ash Content

The ash content in bran fraction of the black rice genotypes, ranged from 14.08-16.80%, in red rice genotypes, it ranged from 10.72-14.75% and in NP rice genotypes it ranged from 10.45-15.40%. In the bran fraction of the black rice genotype CP, red rice genotype YKM, and NP rice genotype Bas 386 displayed higher amounts of ash that is; 16.80, 14.35, and 15.4%. Black rice possessed a high amount of ash followed by NP rice and red rice genotypes. Kalpanadevi et al. [20] reported that the bran from the pigmented rice (Jyothi) possessed marginally higher ash content (10.2–14.2%). Deepa et al [21] reported that most minerals are abundant in the pigmented varieties. Ao et al. [22] stated that the ash content of different cultivars of rice may be due to genetic factors or the mineral content of the soil. Ramaiah and Rao [23] reported that the zinc and iron content of red rice is two to three times higher than that of white rice. The most common minerals found in rice includes calcium, magnesium, iron, and zinc. Resurreccion et al. [24] reported that the outer layers (bran) of the rice grain had the highest concentration of ash content and it decreases towards the centre of the grain.

3.5 Crude Fibre

Crude fibre is the organic residue that remains after the food sample has been treated under the standardized condition with boiling dilute sulphuric acid, boiling dilute sodium hydroxide solution, and alcohol. The fibre in the diet increases the bulk of faeces, which has a laxative effect on the gut [25]. In the present study crude fibre content in bran fraction, in black rice genotypes, ranged from 11.28-12.35%, in red rice genotypes, it ranged from 5.93-10.72%, and in NP rice genotypes it ranged from 5.91-10.24% (Table 1). In bran fraction, black rice genotype FCP, red rice genotype BKG and NP rice PB1121 possessed the highest crude fibre content of 12.35, 10.72 and 10.24 respectively. Red rice is known to be rich in iron and zinc, while black and purple rice is especially high in protein and crude fibre. Priya et al. [19] reported that among pigmented rice varieties, Chak-hao amubi (Manipur black rice) has a significantly lower content of crude fibre. In the current study, we found high crude fibre content in black rice genotypes in all three fractions followed by red and NP rice genotypes. In all the nutritional components determined in the three colours, the black rice genotypes possessed the highest

amount of ash, crude protein, crude fat, and crude fibre content. We didn't find distinct differentiation in the nutritional components among the red and NP rice groups [26-28].

The results are presented as percentage, and values in each column with different capital letters are significantly different ($P < 0.001$) means of the three coloured groups, while the different small letters in each column are significantly different ($P < 0.001$) proximate composition like moisture (%), ash (%), crude protein (%), Crude fat (%) and crude fiber (%) of bran in each coloured group

4. CONCLUSION

The primary interest of this study is to unveil the proximate composition of various pigmented rice which in turn serves as an alternative food for the malnourished. Therefore, bran which by-product of rice milling that was observed appeared to have an excellent nutrition, among the three colour groups black rice bran has identified best.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Sathya A. Are the Indian rice landraces a heritage of biodiversity to reminisce their past or to reinvent for future. *Asian Agrihist.* 2013;17:221-32.
2. Sharif MK, Butt MS, Anjum FM, Khan SH. Rice bran: a novel functional ingredient. *Crit Rev Food Sci Nutr.* 2014;54:807-16.
3. Park HY, Lee KW, Choi HD. Rice bran constituents: Immunomodulatory and therapeutic activities. *Food Funct.* 2017; 8:935-43.
4. Devi R, Veliveli VL, Devi SS. Nutritional composition of rice bran and its potentials in the development of nutraceuticals rich products. *J Pharmacogn Phytochem.* 2021;10:470-73.
5. Kennedy G, Burlingame B, Nguyen N. Nutrient impact assessment of rice in major rice-consuming countries: NewsI, Food and Agriculture Organization, Rome. 2002;51:33-41.
6. Ma ZH, Wang YB, Cheng HT, Zhang GC, Lyu WY. Biochemical composition

- distribution in different grain layers is associated with the edible quality of rice cultivars. *Food chem.* 2020;311:125896.
7. Usman H, Abdulrahman FI, Kaita HA, Khan IZ, Tijjani MA. Flavonoids: The bioactive phytochemical agent—A review. *Chem Res J.* 2017;2:59-72.
 8. Ravichanthiran K, Ma ZF, Zhang H, Cao Y, Wang CW, Muhammad S, Aglago EK, Zhang Y, Jin Y, Pan B. Phytochemical profile of brown rice and its nutrigenomic implications. *Antioxidants.* 2018;7:71.
 9. Adebiji AP, Adebiji AO, Ogawa T, Muramoto K. Preparation and characterization of high-quality rice bran proteins. *J Sci Food Agric.* 2007;87:1219-27.
 10. Shih FF, Champagne ET, Daigle K, Zarins Z. Use of enzymes in the processing of protein products from rice bran and rice flour. *Nahrung.* 1999 ;43:14–18.
 11. Wang M, Hettiarachchy NS, Qi M, Burks W, Siebenmorgen T. Preparation and functional properties of rice bran protein isolate. *J Agric Food Chem.* 1999;47:411–16.
 12. Adebiji AO, Adebiji AP, Ogawa T, Muramoto K. Purification and characterization of antioxidative peptides derived from rice bran protein hydrolysates. *Int J Food Sci Tech.* 2009b;43:35-43.
 13. Champagne ET, Thompson JF, Bett-Garber KL, Muttters R, Miller JA, Tan E. Impact of storage of freshly harvested paddy rice on milled white rice flavor. *Cereal Chem.* 2004b;81:444-49.
 14. Ebuehi OAT, Oyewole AC. Effect of cooking and soaking on physical characteristics, nutrient composition and sensory evaluation of indigenous and foreign rice varieties in Nigeria. *Afr J Biotechnol.* 2007;6:1016-20.
 15. Xheng X, Lan Y. Effects of drying temperature and moisture content on rice taste quality. In *Proc 5th Asia-Pacific Drying Confr.* Hong-Kong University of Science and Technology, China. 2007;2:1112-1117.
 16. Simonelli C, Abbiati A, Cormegna M. Physicochemical characterization of some Italian rice varieties. *J Food Sci Nutr.* 2016;45:9-23.
 17. Suwannaporn P, Pitiphunpong S, Champangern S. Classification of rice amylose content by discriminant analysis of physicochemical properties. *Starch.* 2007;59:171-77.
 18. Choudhury NH, Juliano BO. Effect of amylose content on the lipids of mature rice grain. *Phytochem.* 1980;19:1385-89.
 19. Priya TS, Eliazer Nelson ARL, Ravichandran K, Antony U. Nutritional and functional properties of coloured rice varieties of South India: a review. *J Ethn Foods.* 2019;6:1-11.
 20. Kalpanadevi C, Singh V, Subramanian R. Influence of milling on the nutritional composition of bran from different rice varieties. *J Food Sci Technol.* 2018; 55:2259-69.
 21. Deepa G, Singh V, Naidu KA. Nutrient composition and physicochemical properties of Indian medicinal rice – Njavara. *Food Chem.* 2008;106:165-71.
 22. Ao O, Be U, Aa E, ND. Comparative analysis of the chemical nutrient composition of selected local and newly introduced rice varieties grown in Ebonyi State of Nigeria. *Int J Agric For.* 2012;2:16-23.
 23. Ramaiah K, Rao MVB. Rice breeding and genetics. Pp.19. New Indian Council of Agricultural Research, New Delhi, India; 1953.
 24. Resurreccion AP, Tanaka Y. Nutrient content and distribution in milling fractions of rice grain. *J Sci Food Agric.* 1979;30:5-481.
 25. Oko A, Okpani and Chijioke S, Ugwu D, Onyedikachi, Kalu N, Onyia C, Ayanniran F, Omoniyi A, Okorie J, Michael, Azubuikokechukwu S. Proximate, mineral compositions and gelatinization temperatures of pap (Ogi) from fermented maize, sorghum and millet. *Interdiscip Sci Rev.* 2019;14:106-13.
 26. Hegde S, Yenagi NB, Kasturiba B. Indigenous knowledge of the traditional and qualified ayurveda practitioners on the nutritional significance and use of red rice in medications. *Indian J Tradit Knowl.* 2013;12:506-11.

27. Juliano BO (ed). Polysaccharides, proteins, and lipids of rice. pp. 59-174. American Association of Cereal Chemists, Minnesota; 1985.
28. Tanaka K, Sugimoto T, Ogawa M, Kasai Z. Isolation and characterization of two types of protein bodies in rice endosperm. *Agric Biol Chem.* 1980;44:1633–39.

© 2023 Avinash and Sharma; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/100929>