



# Field Efficacy of Insecticides and Botanicals for Management of Spotted Stem Borer (*Chilo partellus*) on Maize (*Zea mays* L.)

Ashok Sakharam Chandar <sup>a</sup> and Anoorag R. Tayde <sup>a\*</sup>

<sup>a</sup> Department of Entomology, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj – 211007, Uttar Pradesh, India.

## Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJPSS/2023/v35i234232

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/110371>

Original Research Article

Received: 11/10/2023  
Accepted: 16/12/2023  
Published: 18/12/2023

## ABSTRACT

The field investigation was conducted during the *Kharif* 2021 and 2022 at the Central Research Farm (CRF) of Naini Agriculture Institute, Sam Higginbottom University of Agriculture Technology and Sciences, Prayagraj, Uttar Pradesh, India. The experiment was organized in a Randomized Block Design, comprising twelve treatments, each replicated thrice. The treatments included Chlorantraniliprole 18.5 SC (0.5 ml/l), Lambda cyhalothrin 2.5% EC (10 ml/l), Thiamethoxam 25 WG (10 gm/kg), Profenophos 50 EC (3ml/l), Cypermethrin 10 EC (4 ml/l), Neem oil 3% (30 ml/l), *Verticillium lecani* 1.15 WP (15 ml/l), Spinosad 45 SC (0.3 ml/l), Emamectin benzoate 5 SG (0.40 gm/kg), Indoxacarb 15.8 EC (1.5 ml/l), NSKE 5% (50 ml/l), and control plot. The results showed that the lowest population of stem borer was identified with Emamectin benzoate 5 SG (2.24), (0.82) and (3.61), (2.33) followed by Chlorantraniliprole 18.5 SC (2.41), (1.01) and (3.82), (2.46) during the *Kharif* seasons of both 2021 and 2022. Neem oil 3%, NSKE 5% were found to be least effective over the other treatments and is significantly superior over the control.

\*Corresponding author: E-mail: anurag.tayde@shiats.edu.in;

**Keywords:** *Chilo partellus*; efficacy; insecticides; maize.

## 1. INTRODUCTION

“Maize, also known as corn, is a widely cultivated cereal grain and is an important staple food in many parts of the world. It is a highly versatile crop that is used for a variety of purposes, including human consumption, animal feed, and industrial products. Maize is grown in diverse agroecological zones, ranging from tropical to temperate regions, and is produced in large quantities globally. According to the Food and Agriculture Organization (FAO), maize is the third most important cereal crop in the world, after wheat and rice, and is grown on an estimated 191 million hectares globally, with a total production of 1.14 billion tonnes in 2021. Maize is an important source of carbohydrates, protein, fiber, and other essential nutrients, and plays a critical role in food security and livelihoods for millions of people worldwide” [1].

“The nutritious cereal grain known as maize, or corn, is a great source of vitamins, minerals, fibre, protein, and carbs. One cup of cooked yellow maize contains 123 calories, 26 grammes of carbohydrates, 3 grammes of protein, 2.5 grammes of fibre, and a number of vital vitamins and minerals, including vitamin C, vitamin B6, potassium, magnesium, and phosphorus, according to the United States Department of Agriculture” (USDA, 2021). Maize has also high antioxidants and has been associated with a number of health advantages, including better digestion, a decreased risk of chronic illnesses, and better heart health [2].

“The average production of maize per hectare in the world increased to 5.9 metric tonnes in 2019 from 4.4 metric tonnes in 2000. These increases in maize productivity have supported economic growth and rural development in many nations while also helping to fulfil the rising demand for food, feed, and industrial purposes. (Source: retrieved March 31, 2023 from FAOSTAT, "Maize - Production, Quantity"). With an output of 28.98 million tonnes of maize in 2020–21, India is one of the top producers in the world” [3]. “Most Indian states cultivate maize, with Karnataka, Andhra Pradesh, Telangana, Bihar, Uttar Pradesh, and Rajasthan being the top producers” [4]. With an average yield of 2.93 tonnes per hectare in 2020-21 compared to 2.80 tonnes per hectare in 2019–20, maize productivity in India

has been continuously rising over time [3]. Use of more advanced technology, including as hybrid cultivars, high-density planting, and enhanced crop management techniques, are responsible for this rise in production [4].

“In many regions of Africa and Asia, the Spotted Stem Borer (*Chilo partellus*), a serious pest of maize crops, reduces yields significantly depending on stage of larvae. Early on in the larvae's development, the plant's whorl and leaves sustain the majority of the damage, which stunts the plant's growth and reduces photosynthesis. But, when the larvae grow older, they begin to dig into the stem, severely harming the vascular tissue, which causes wilting and ultimately plant death. According to studies, the yield losses due by the stem borer range from 10% to 70%” [5,6]. To lessen the harm due to stem borer, efficient control methods are required, such as the adoption of resistant maize cultivars and integrated pest management techniques.

## 2. MATERIALS AND METHODS

The study was carried out during the *Kharif* seasons of 2021 and 2022 at the Central Research Farm (CRF) of SHUATS, Prayagraj, Uttar Pradesh, India. The experimental design employed was a Randomized Block Design with twelve treatments, each replicated three times. The plot size was (2m×2m) with a spacing of (60×20cm). The recommended package of practices, excluding plant protection, was followed. The treatments included Chlorantraniliprole 18.5 SC (0.5 ml/l), Lambda cyhalothrin 2.5% EC (10 ml/l), Thiamethoxam 25 WG (10 gm/kg), Profenophos 50 EC (3ml/l), Cypermethrin 10 EC (4 ml/l), Neem oil 3% (30 ml/l), *Verticillium lecani* 1.15 WP (15 ml/l), Spinosad 45 SC (0.3 ml/l), Emamectin benzoate 5 SG (0.40 gm/kg), Indoxacarb 15.8 EC (1.5 ml/l), NSKE 5% (50 ml/l), and a control.

Two sprays were carried out at interval of 15 days during the experimentation to assess the effectiveness of insecticides on the *chilo partellus* larval population. Five plants were randomly selected in each treatment and observations were taken one day before spraying application and three, seven and fourteen-days after spraying. Foliar and granular insecticides were

applied at their recommended doses at economic threshold level (ETL=10% infestation) [7].

### 3. RESULTS AND DISCUSSION

#### 3.1 Efficacy of *Chilo partellus* Kharif 2021 (First Spray)

**Three days after spraying:** The data on the larval population of spotted stem borer on three days after spray revealed that all treatments were significantly superior over control. Among the treatments lowest larval population of spotted stem borer was recorded in Emamectin benzoate 5 SG (1.53) followed by Chlorantraniliprole 18.5 SC (1.66), Indoxacarb 15.8 EC (1.93), Thiamethoxam 25 WG (2.20), Lambda cyhalothrin 2.5% EC (2.26) and Spinosad 45 SC (2.33). In this Profenophos 50 EC (2.53), Cypermethrin 10 EC (2.80), *Verticillium lecani* 1.15 WP (2.93), Neem oil 3% (3.13) and NSKE 5% (3.33) is found to be least effective than all the treatments and is significantly superior over the control (3.60).

**Seven days after spraying:** The data on the larval population of spotted stem borer on seven days after spray revealed that all treatments were significantly superior over control. Among the treatments lowest larval population of spotted stem borer was recorded in Emamectin benzoate 5 SG (0.40) followed by Chlorantraniliprole 18.5 SC (0.46), Indoxacarb 15.8 EC (0.60), Thiamethoxam 25 WG (0.66), Lambda cyhalothrin 2.5% EC (0.80) and Spinosad 45 SC (1.06). In this Profenophos 50 EC (1.26), Cypermethrin 10 EC (1.33), *Verticillium lecani* 1.15 WP (1.73), Neem oil 3% (1.86) and NSKE 5% (2.13) is found to be least effective than all the treatments and is significantly superior over the control (3.86).

**Fourteen days after spraying:** The data on the larval population of spotted stem borer on fourteen days after spray revealed that all treatments were significantly superior over control. Among the treatments lowest larval population of spotted stem borer was recorded in Emamectin benzoate 5 SG (1.06) followed by Chlorantraniliprole 18.5 SC (1.20), Indoxacarb 15.8 EC (1.33), Thiamethoxam 25 WG (1.46), Lambda cyhalothrin 2.5% EC (1.66) and Spinosad 45 SC (1.73). In this Profenophos 50 EC (1.80), Cypermethrin 10 EC (2.26),

*Verticillium lecani* 1.15 WP (2.53), Neem oil 3% (2.66), NSKE 5% (2.93) is found to be least effective than all the treatments and is significantly superior over the control (4.20).

**Overall mean of first spray:** The data on the larval population of spotted stem borer on mean (3<sup>rd</sup>, 7<sup>th</sup> and 14<sup>th</sup> DAS) days after spray revealed that all treatments were significantly superior over control. Among the treatments lowest larval population of spotted stem borer was recorded in Emamectin benzoate 5 SG (0.99) followed by Chlorantraniliprole 18.5 SC (1.10), Indoxacarb 15.8 EC (1.28), Thiamethoxam 25 WG (1.44), Lambda cyhalothrin 2.5% EC (1.57) and Spinosad 45 SC (1.70). In this Profenophos 50 EC (1.86), Cypermethrin 10 EC (2.13), *Verticillium lecani* 1.15 WP (2.39), Neem oil 3% (2.55) and NSKE 5% (2.79) is found to be least effective than all the treatments and is significantly superior over the control (3.88).

#### 3.2 Efficacy of *Chilo partellus* Kharif 2021 (Second Spray)

**Three days after spraying:** The data on the larval population of spotted stem borer on three days after spray revealed that all treatments were significantly superior over control. Among the treatments lowest larval population of spotted stem borer was recorded in Emamectin benzoate 5 SG (0.60) followed by Chlorantraniliprole 18.5 SC (0.80), Indoxacarb 15.8 EC (0.86), Thiamethoxam 25 WG (1.06), Lambda cyhalothrin 2.5% EC (1.20) and Spinosad 45 SC (1.33). In this Profenophos 50 EC (1.60), Cypermethrin 10 EC (1.80), *Verticillium lecani* 1.15 WP (2.13), Neem oil 3% (2.20) and NSKE 5% (2.40) is found to be least effective than all the treatments and is significantly superior over the control (4.46).

**Seven days after spraying:** The data on the larval population of spotted stem borer on seven days after spray revealed that all treatments were significantly superior over control. Among the treatments lowest larval population of spotted stem borer was recorded in Emamectin benzoate 5 SG (0.20) followed by Chlorantraniliprole 18.5 SC (0.26), Indoxacarb 15.8 EC (0.40), Thiamethoxam 25 WG (0.60), Lambda cyhalothrin 2.5% EC (0.73) and Spinosad 45 SC (0.93). In this T<sub>4</sub> Profenophos 50 EC (1.20), Cypermethrin 10 EC (1.33), *Verticillium lecani*

1.15 WP (1.46), Neem oil 3% (1.73) and NSKE 5% (2.20) is found to be least effective than all the treatments and is significantly superior over the control (5.00).

**Fourteen days after spraying:** The data on the larval population of spotted stem borer on fourteen days after spray revealed that all treatments were significantly superior over control. Among the treatments lowest larval population of spotted stem borer was recorded in Emamectin benzoate 5 SG (0.40) followed by Chlorantraniliprole 18.5 SC (0.53), Indoxacarb 15.8 EC (0.60), Thiamethoxam 25 WG (0.80), Lambda cyhalothrin 2.5% EC (0.93) and Spinosad 45 SC (1.06). In this Profenophos 50 EC (1.40), Cypermethrin 10 EC (1.53), *Verticillium lecani* 1.15 WP (1.80), Neem oil 3% (2.00) and NSKE 5% (2.46) is found to be least effective than all the treatments and is significantly superior over the T<sub>0</sub> control (5.33).

**Overall mean of second spray:** The data on the larval population of spotted stem borer on mean (3<sup>rd</sup>, 7<sup>th</sup> and 14<sup>th</sup> DAS) days after spray revealed that all treatments were significantly superior over control. Among the treatments lowest larval population of spotted stem borer was recorded in Emamectin benzoate 5 SG (0.40) followed by Chlorantraniliprole 18.5 SC (0.53), Indoxacarb 15.8 EC (0.62), Thiamethoxam 25 WG (0.82), Lambda cyhalothrin 2.5% EC (0.95) and Spinosad 45 SC (1.10). In this Profenophos 50 EC (1.40), Cypermethrin 10 EC (1.55), T<sub>7</sub> *Verticillium lecani* 1.15 WP (1.79), Neem oil 3% (1.97) and NSKE 5% (2.35) is found to be least effective than all the treatments and is significantly superior over the control (4.93).

### 3.3 Efficacy of *Chilo partellus* Kharif 2022 (First Spray)

**Three days after spraying:** The data on the larval population of spotted stem borer on three days after spray revealed that all treatments were significantly superior over control. Among the treatments lowest larval population of spotted stem borer was recorded in Emamectin benzoate 5 SG (1.66) followed by Chlorantraniliprole 18.5 SC (1.73), Indoxacarb 15.8 EC (1.80), Thiamethoxam 25 WG (1.86), Lambda cyhalothrin 2.5% EC (1.93) and Spinosad 45 SC (2.00). In this Profenophos 50 EC (2.06), Cypermethrin 10 EC (2.13), *Verticillium lecani*

1.15 WP (2.20), Neem oil 3% (2.26) and NSKE 5% (2.33) is found to be least effective than all the treatments and is significantly superior over the control (3.86).

**Seven days after spraying:** The data on the larval population of spotted stem borer on seven days after spray revealed that all treatments were significantly superior over control. Among the treatments lowest larval population of spotted stem borer was recorded in Emamectin benzoate 5 SG (0.40) followed by Chlorantraniliprole 18.5 SC (0.46), Indoxacarb 15.8 EC (0.53), Thiamethoxam 25 WG (0.60), Lambda cyhalothrin 2.5% EC (0.66) and Spinosad 45 SC (0.73). In this Profenophos 50 EC (0.86), Cypermethrin 10 EC (0.93), *Verticillium lecani* 1.15 WP (1.00), Neem oil 3% (1.06) and NSKE 5% (1.20) is found to be least effective than all the treatments and is significantly superior over the control (4.00).

**Fourteen days after spraying:** The data on the larval population of spotted stem borer on fourteen days after spray revealed that all treatments were significantly superior over control. Among the treatments lowest larval population of spotted stem borer was recorded in Emamectin benzoate 5 SG (0.80) followed by Chlorantraniliprole 18.5 SC (0.86), Indoxacarb 15.8 EC (0.93), Thiamethoxam 25 WG (1.00), Lambda cyhalothrin 2.5% EC (1.06) and Spinosad 45 SC (1.13). In this Profenophos 50 EC (1.26), Cypermethrin 10 EC (1.33), *Verticillium lecani* 1.15 WP (1.40), Neem oil 3% (1.46) and NSKE 5% (1.60) is found to be least effective than all the treatments and is significantly superior over the control (4.13).

**Overall mean of first spray:** The data on the larval population of spotted stem borer on mean (3<sup>rd</sup>, 7<sup>th</sup> and 14<sup>th</sup> DAS) days after spray revealed that all treatments were significantly superior over control. Among the treatments lowest larval population of spotted stem borer was recorded in Emamectin benzoate 5 SG (0.95) followed by Chlorantraniliprole 18.5 SC (1.01), Indoxacarb 15.8 EC (1.08), Thiamethoxam 25 WG (1.15), Lambda cyhalothrin 2.5% EC (1.21) and Spinosad 45 SC (1.28). In this Profenophos 50 EC (1.39), Cypermethrin 10 EC (1.46), *Verticillium lecani* 1.15 WP (1.53), Neem oil 3% (1.59) and NSKE 5% (1.71) is found to be least effective than all the treatments and is significantly superior over the control (3.99).

Table 1. Efficacy against *Chilo partellus* Kharif-2021

Treatments	Larval population on number basis/5 plants										Overall Mean
	One day before spray	After First Spray				After Second Spray					
		3 DAS	7 DAS	14 DAS	Mean	3 DAS	7 DAS	14 DAS	Mean		
T <sub>1</sub>	Chlorantraniprole 18.5 % SC	3.20	1.66 <sup>ij</sup>	0.46 <sup>f</sup>	1.20 <sup>f</sup>	1.10 <sup>hi</sup>	0.80 <sup>hi</sup>	0.26 <sup>i</sup>	0.53 <sup>gh</sup>	0.53 <sup>hi</sup>	0.81 <sup>g</sup>
T <sub>2</sub>	Lambda Cyhalothrin 5% EC	3.06	2.26 <sup>fg</sup>	0.80 <sup>ef</sup>	1.66 <sup>de</sup>	1.57 <sup>efg</sup>	1.20 <sup>fg</sup>	0.73 <sup>fg</sup>	0.93 <sup>f</sup>	0.95 <sup>g</sup>	1.26 <sup>efg</sup>
T <sub>3</sub>	Thiamethoxam 25% WG	3.46	2.20 <sup>gh</sup>	0.66 <sup>ef</sup>	1.46 <sup>def</sup>	1.44 <sup>fgh</sup>	1.06 <sup>fgh</sup>	0.60 <sup>gh</sup>	0.80 <sup>fg</sup>	0.82 <sup>gh</sup>	1.13 <sup>efg</sup>
T <sub>4</sub>	Profenophos 50% EC	3.53	2.53 <sup>ef</sup>	1.26 <sup>d</sup>	1.80 <sup>d</sup>	1.86 <sup>de</sup>	1.60 <sup>de</sup>	1.20 <sup>de</sup>	1.40 <sup>e</sup>	1.40 <sup>ef</sup>	1.63 <sup>cdef</sup>
T <sub>5</sub>	Cypermethrin 10% EC	3.46	2.80 <sup>de</sup>	1.33 <sup>cd</sup>	2.26 <sup>c</sup>	2.13 <sup>cd</sup>	1.80 <sup>cd</sup>	1.33 <sup>d</sup>	1.53 <sup>de</sup>	1.55 <sup>de</sup>	1.84 <sup>bcde</sup>
T <sub>6</sub>	Neem oil 3%	3.53	3.13 <sup>bc</sup>	1.86 <sup>b</sup>	2.66 <sup>bc</sup>	2.55 <sup>b</sup>	2.20 <sup>b</sup>	1.73 <sup>c</sup>	2.00 <sup>c</sup>	1.97 <sup>c</sup>	2.26 <sup>bc</sup>
T <sub>7</sub>	<i>Verticillium lecani</i> 1.15 % WP	3.46	2.93 <sup>cd</sup>	1.73 <sup>bc</sup>	2.53 <sup>bc</sup>	2.39 <sup>bc</sup>	2.13 <sup>bc</sup>	1.46 <sup>cd</sup>	1.80 <sup>cd</sup>	1.79 <sup>cd</sup>	2.09 <sup>bcd</sup>
T <sub>8</sub>	Spinosad 45% SC	3.06	2.33 <sup>fg</sup>	1.06 <sup>de</sup>	1.73 <sup>de</sup>	1.70 <sup>ef</sup>	1.33 <sup>ef</sup>	0.93 <sup>ef</sup>	1.06 <sup>f</sup>	1.10 <sup>fg</sup>	1.40 <sup>defg</sup>
T <sub>9</sub>	Emamectin benzoate 5 % SG	3.26	1.53 <sup>j</sup>	0.40 <sup>f</sup>	1.06 <sup>f</sup>	0.99 <sup>i</sup>	0.60 <sup>i</sup>	0.20 <sup>i</sup>	0.40 <sup>h</sup>	0.40 <sup>i</sup>	0.69 <sup>g</sup>
T <sub>10</sub>	Indoxacarb 15.8% EC	3.53	1.93 <sup>hi</sup>	0.60 <sup>f</sup>	1.33 <sup>ef</sup>	1.28 <sup>ghi</sup>	0.86 <sup>ghi</sup>	0.40 <sup>hi</sup>	0.60 <sup>gh</sup>	0.62 <sup>hi</sup>	0.95 <sup>fg</sup>
T <sub>11</sub>	NSKE 5 %	3.86	3.33 <sup>ab</sup>	2.13 <sup>b</sup>	2.93 <sup>b</sup>	2.79 <sup>b</sup>	2.40 <sup>b</sup>	2.20 <sup>b</sup>	2.46 <sup>b</sup>	2.35 <sup>b</sup>	2.57 <sup>b</sup>
T <sub>0</sub>	Control	3.20	3.60 <sup>a</sup>	3.86 <sup>a</sup>	4.20 <sup>a</sup>	3.88 <sup>a</sup>	4.46 <sup>a</sup>	5.00 <sup>a</sup>	5.33 <sup>a</sup>	4.93 <sup>a</sup>	4.40 <sup>a</sup>
F-test		NS	S	S	S	S	S	S	S	S	S
S.Ed ( $\pm$ )		1.68	0.13	0.20	0.19	0.20	0.78	0.13	0.73	0.15	0.27
C.D. (P=0.05)		-	0.27	0.42	0.40	0.41	0.34	0.28	0.32	0.31	0.73

DBS- Day before Spraying, DAS- Day after Spraying, NS- Non significant, S- Significant

Table 2. Efficacy against *Chilo partellus* Kharif-2022

Treatments	Larval population on number basis/5 plants										Overall Mean	Pooled Mean
	One day before spray	After First Spray				After Second Spray						
		3 DAS	7 DAS	14 DAS	Mean	3 DAS	7 DAS	14 DAS	Mean			
T <sub>1</sub>	Chlorantraniprole 18.5 % SC	3.80	1.73 <sup>fg</sup>	0.46 <sup>bc</sup>	0.86 <sup>gh</sup>	1.01 <sup>g</sup>	0.66 <sup>gh</sup>	0.33 <sup>ij</sup>	0.40 <sup>hi</sup>	0.46 <sup>hi</sup>	0.73 <sup>fg</sup>	0.77 <sup>g</sup>
T <sub>2</sub>	Lambda Cyhalothrin 5% EC	3.93	1.93 <sup>cdefg</sup>	0.66 <sup>efg</sup>	1.06 <sup>efg</sup>	1.21 <sup>defg</sup>	0.86 <sup>efg</sup>	0.53 <sup>fgh</sup>	0.60 <sup>fg</sup>	0.66 <sup>fg</sup>	0.93 <sup>cdefg</sup>	1.09 <sup>efg</sup>
T <sub>3</sub>	Thiamethoxam 25% WG	3.86	1.86 <sup>defg</sup>	0.60 <sup>fgh</sup>	1.00 <sup>fgh</sup>	1.15 <sup>efg</sup>	0.80 <sup>fgh</sup>	0.46 <sup>ghi</sup>	0.53 <sup>fgh</sup>	0.59 <sup>fgh</sup>	0.87 <sup>defg</sup>	1.00 <sup>efg</sup>
T <sub>4</sub>	Profenophos 50% EC	4.00	2.06 <sup>bcdef</sup>	0.86 <sup>cde</sup>	1.26 <sup>cde</sup>	1.39 <sup>bcdef</sup>	1.06 <sup>cde</sup>	0.66 <sup>def</sup>	0.80 <sup>de</sup>	0.84 <sup>de</sup>	1.11 <sup>bcdef</sup>	1.37 <sup>cdef</sup>
T <sub>5</sub>	Cypermethrin 10% EC	4.06	2.13 <sup>bcde</sup>	0.93 <sup>cd</sup>	1.33 <sup>cd</sup>	1.46 <sup>bcde</sup>	1.13 <sup>cd</sup>	0.73 <sup>cde</sup>	0.86 <sup>cd</sup>	0.90 <sup>cd</sup>	1.18 <sup>bcde</sup>	1.51 <sup>bcde</sup>
T <sub>6</sub>	Neem oil 3%	4.20	2.26 <sup>bc</sup>	1.06 <sup>bc</sup>	1.46 <sup>bc</sup>	1.59 <sup>bc</sup>	1.26 <sup>bc</sup>	0.86 <sup>bc</sup>	1.00 <sup>bc</sup>	1.04 <sup>bc</sup>	1.31 <sup>bc</sup>	1.78 <sup>bc</sup>
T <sub>7</sub>	<i>Verticillium lecani</i> 1.15 % WP	4.13	2.20 <sup>bcd</sup>	1.00 <sup>bc</sup>	1.40 <sup>bc</sup>	1.53 <sup>bcd</sup>	1.20 <sup>bc</sup>	0.80 <sup>cd</sup>	0.93 <sup>cd</sup>	0.97 <sup>cd</sup>	1.25 <sup>bcd</sup>	1.67 <sup>bcd</sup>
T <sub>8</sub>	Spinosad 45% SC	4.00	2.00 <sup>bcdefg</sup>	0.73 <sup>def</sup>	1.13 <sup>def</sup>	1.28 <sup>cdefg</sup>	0.93 <sup>def</sup>	0.60 <sup>efg</sup>	0.66 <sup>ef</sup>	0.73 <sup>ef</sup>	1.00 <sup>cdefg</sup>	1.20 <sup>defg</sup>
T <sub>9</sub>	Emamectin benzoate 5 % SG	3.66	1.66 <sup>d</sup>	0.40 <sup>h</sup>	0.80 <sup>h</sup>	0.95 <sup>d</sup>	0.60 <sup>h</sup>	0.26 <sup>i</sup>	0.33 <sup>i</sup>	0.39 <sup>i</sup>	0.67 <sup>d</sup>	0.68 <sup>d</sup>
T <sub>10</sub>	Indoxacarb 15.8% EC	3.86	1.80 <sup>efg</sup>	0.53 <sup>fgh</sup>	0.93 <sup>fgh</sup>	1.08 <sup>fg</sup>	0.73 <sup>fgh</sup>	0.40 <sup>hij</sup>	0.46 <sup>ghi</sup>	0.53 <sup>ghi</sup>	0.80 <sup>efg</sup>	0.87 <sup>fg</sup>
T <sub>11</sub>	NSKE 5 %	4.26	2.33 <sup>b</sup>	1.20 <sup>b</sup>	1.60 <sup>b</sup>	1.71 <sup>b</sup>	1.40 <sup>b</sup>	1.00 <sup>b</sup>	1.13 <sup>b</sup>	1.17 <sup>b</sup>	1.44 <sup>b</sup>	2.00 <sup>b</sup>
T <sub>0</sub>	Control	3.73	3.86 <sup>a</sup>	4.00 <sup>a</sup>	4.13 <sup>a</sup>	3.99 <sup>a</sup>	4.26 <sup>a</sup>	4.40 <sup>a</sup>	4.56 <sup>a</sup>	4.37 <sup>a</sup>	4.18 <sup>a</sup>	4.29 <sup>a</sup>
F-test		NS	S	S	S	S	S	S	S	S	S	S
S.Ed ( $\pm$ )		0.20	0.16	0.10	0.11	0.17	0.10	0.08	0.09	0.06	0.15	0.25
C.D. (P=0.05)		-	0.34	0.20	0.23	0.35	0.21	0.18	0.19	0.14	0.41	0.56

DBS- Day before Spraying, DAS- Day after Spraying, NS- Non significant, S- Significant

### 3.4 Efficacy of *Chilo partellus* Kharif 2022 (Second Spray)

**Three days after spraying:** The data on the larval population of spotted stem borer on three days after spray revealed that all treatments were significantly superior over control. Among the treatments lowest larval population of spotted stem borer was recorded in Emamectin benzoate 5 SG (0.60) followed by Chlorantraniliprole 18.5 SC (0.66), Indoxacarb 15.8 EC (0.73), Thiamethoxam 25 WG (0.80), Lambda cyhalothrin 2.5% EC (0.86) and Spinosad 45 SC (0.93). In this Profenophos 50 EC (1.06), Cypermethrin 10 EC (1.13), *Verticillium lecani* 1.15 WP (1.20), Neem oil 3% (1.26) and NSKE 5% (1.40) is found to be least effective than all the treatments and is significantly superior over the control (4.26).

**Seven days after spraying:** The data on the larval population of spotted stem borer on seven days after spray revealed that all treatments were significantly superior over control. Among the treatments lowest larval population of spotted stem borer was recorded in Emamectin benzoate 5 SG (0.26) followed by Chlorantraniliprole 18.5 SC (0.33), Indoxacarb 15.8 EC (0.40), Thiamethoxam 25 WG (0.46), Lambda cyhalothrin 2.5% EC (0.53) and Spinosad 45 SC (0.60). In this Profenophos 50 EC (0.66), Cypermethrin 10 EC (0.73), *Verticillium lecani* 1.15 WP (0.80), Neem oil 3% (0.86) and NSKE 5% (1.00) is found to be least effective than all the treatments and is significantly superior over the control (4.40).

**Fourteen days after spraying:** The data on the larval population of spotted stem borer on fourteen days after spray revealed that all treatments were significantly superior over control. Among the treatments lowest larval population of spotted stem borer was recorded in Emamectin benzoate 5 SG (0.33) followed by Chlorantraniliprole 18.5 SC (0.40), Indoxacarb 15.8 EC (0.46), Thiamethoxam 25 WG (0.53), Lambda cyhalothrin 2.5% EC (0.60) and Spinosad 45 SC (0.66). In this Profenophos 50 EC (0.80), Cypermethrin 10 EC (0.86), *Verticillium lecani* 1.15 WP (0.93), Neem oil 3% (1.00) and NSKE 5% (1.13) is found to be least effective than all the treatments and is significantly superior over the control (4.56).

**Overall mean of second spray:** The data on the larval population of spotted stem borer on mean (3<sup>rd</sup>, 7<sup>th</sup> and 14<sup>th</sup> DAS) days after spray revealed that all treatments were significantly superior over control. Among the treatments lowest larval population of spotted stem borer was recorded in Emamectin benzoate 5 SG (0.39) followed by Chlorantraniliprole 18.5 SC (0.46), Indoxacarb 15.8 EC (0.53), Thiamethoxam 25 WG (0.59), Lambda cyhalothrin 2.5% EC (0.66) and Spinosad 45 SC (0.73). In this Profenophos 50 EC (0.84), Cypermethrin 10 EC (0.90), *Verticillium lecani* 1.15 WP (0.97), Neem oil 3% (1.04) and NSKE 5% (1.17) is found to be least effective than all the treatments and is significantly superior over the control (4.37).

**Pooled mean:** The data on the larval population of spotted stem borer on pooled mean revealed that all treatments were significantly superior over control. Among the treatments lowest larval population of spotted stem borer was recorded in Emamectin benzoate 5 SG (0.68) followed by Chlorantraniliprole 18.5 SC (0.77), Indoxacarb 15.8 EC (0.87), Thiamethoxam 25 WG (1.00), Lambda cyhalothrin 2.5% EC (1.09) and Spinosad 45 SC (1.20). In this Profenophos 50 EC (1.37), Cypermethrin 10 EC (1.51), *Verticillium lecani* 1.15 WP (1.67), Neem oil 3% (1.78) and NSKE 5% (2.00) is found to be least effective than all the treatments and is significantly superior over the control (4.29).

## 4. DISCUSSION

### 4.1 Overall (First and Second Spray) *Chilo partellus* (Kharif 2021)

The data on the larval population of spotted stem borer on overall mean (first and second spray) revealed that all treatments were significantly superior over control. Among the treatments lowest larval population of spotted stem borer was recorded in Emamectin benzoate 5 SG (0.69) followed by Chlorantraniliprole 18.5 SC (0.81), Indoxacarb 15.8 EC (0.95), Thiamethoxam 25 WG (1.13), Lambda cyhalothrin 2.5% EC (1.26) and Spinosad 45 SC (1.40). In this Profenophos 50 EC (1.63), Cypermethrin 10 EC (1.84), *Verticillium lecani* 1.15 WP (2.09), Neem oil 3% (2.26) and NSKE 5% (2.57) is found to be least effective than all the treatments and is significantly superior over the control (4.40).

Among all the treatments, Emamectin benzoate 5 % SG was found to be most effective in managing the larval population of spotted stem borer. The values obtained in the first and second spray were (0.99) and (0.40). These results are supported by Rathod et al. [8]. Chlorantraniliprole 18.5% SC was also found to be very effective in reducing the larval population of spotted stem borer. The values obtained in the first and second spray were (1.10) and (0.53). The same results were observed by Arunkumara et al. [9]; Devananda et al. [10]. The application of Indoxacarb 15.8% EC reduced the larval population of spotted stem borer. Where the observations of first and second sprays obtained were (1.28) and (0.62). These results are supported by Prakash et al. [11]; Rathod et al. [8]. The efficacy of Thiamethoxam 25% WG on spotted stem borer in first and second spray were (1.44) and (0.82). These results are as per the findings of Joshi et al. [12].

The next best treatment was found to be Lambda Cyhalothrin 5% EC in which the efficacy values of first and second spray were (1.57) and (0.95) respectively to the similar findings of Prakash et al. [11]. The next best treatment was found to be Spinosad 45% SC in which the efficacy values of first and second spray were (1.70) and (1.10) respectively; these results were supported by Joshi et al. [12] and Malar *et al.*, (2018). The next effective treatment was found to be Profenophos 50% EC in which efficacy values of first and second sprays were (1.86) and (1.40) respectively; these results were supported by Devananda et al. [10]. The efficacy of Cypermethrin 10% EC on spotted stem borer in first and second spray were (2.13) and (1.55). These results are as per the findings of Ahmad et al. [13]; Bhat and Baba [14]; Dinesh et al. [15].

#### **4.2 Overall (First and Second Spray) *Chilo partellus* (Kharif 2022)**

The data on the larval population of spotted stem borer on overall mean (first and second spray) revealed that all treatments were significantly superior over control. Among the treatments lowest larval population of spotted stem borer was recorded in Emamectin benzoate 5 SG (0.67) followed by Chlorantraniliprole 18.5 SC (0.73), Indoxacarb 15.8 EC (0.80), Thiamethoxam 25 WG (0.87), Lambda cyhalothrin 2.5% EC (0.93) and Spinosad 45 SC (1.00). In this Profenophos 50 EC (1.11),

Cypermethrin 10 EC (1.18), *Verticillium lecani* 1.15 WP (1.25), Neem oil 3% (1.31) and NSKE 5% (1.44) is found to be least effective than all the treatments and is significantly superior over the control (4.18).

Among all the treatments, Emamectin benzoate 5 % SG was found to be most effective in managing the larval population of spotted stem borer. The values obtained in the first and second spray were (0.95) and (0.39). These results are supported by Rathod et al. [8]. Chlorantraniliprole 18.5% SC was also found to be very effective in reducing the larval population of spotted stem borer. The values obtained in the first and second spray were (1.01) and (0.46). The same results were observed by Kumar et al. [16]; Kumar and Jindal [17]. The application of Indoxacarb 15.8% EC reduced the larval population of spotted stem borer. Where the observations of first and second sprays obtained were (1.08) and (0.53). These results are supported by Devananda et al. [10]; Prakash et al. [11]. The efficacy of Thiamethoxam 25% WG on spotted stem borer in first and second spray were (1.15) and (0.59). These results are as per the findings of Joshi et al. [12].

The next best treatment was found to be Lambda Cyhalothrin 5% EC in which the efficacy values of first and second spray were (1.21) and (0.66) respectively to the similar findings of Prakash et al. [11]. The next best treatment was found to be Spinosad 45% SC in which the efficacy values of first and second spray were (1.28) and (0.73) respectively; these results were supported by Devananda *et al.*, [10]; Kumar et al. [16]. The next effective treatment was found to be Profenophos 50% EC in which efficacy values of first and second sprays were (1.39) and (0.84) respectively; these results were supported Justin and Preetha [18]. The efficacy of Cypermethrin 10% EC on spotted stem borer in first and second spray were (1.46) and (0.90). These results are as per the findings of Babu and Kumar [19]; Reddy and Kumar [20]; Prakash et al. [11]; [21].

#### **5. CONCLUSION**

From the current investigation, the findings indicate that Emamectin benzoate 5% SG succeeded by Chlorantraniliprole 18.5% SC and Indoxacarb 15.8% EC, emerge as the most efficacious treatments against the maize spotted

stem borer. These treatments not only resulted in the highest yield but also demonstrated the most favourable Cost-Benefit ratio when compared to alternative treatments. Thiamethoxam 25% WG, Lambda Cyhalothrin 5% EC, Spinosad 45% SC, and Profenophos 50% EC exhibited moderate efficacy in the management of the spotted stem borer. Conversely, Cypermethrin 10% EC, *Verticillium lecani* 1.15 % WP, Neem oil 3%, and NSKE 5% were identified as the least effective in controlling all maize pests. Based on these findings, it is strongly recommended to integrate the use of effective insecticides into existing Integrated Pest Management programs. Emphasis should be placed on preventing issues such as insecticide resistance and pest resurgence. The inclusion of botanical substances is crucial in Integrated Pest Management as it helps mitigate the indiscriminate use of pesticides, minimizing environmental harm, and preserving the impact on beneficial insects.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. FAOSTAT; 2021.
2. Source: United State Department of Agriculture. USDA; 2021.
3. Ministry of agriculture and farmer welfare; 2021.
4. Directorate of Maize Research; 2021.
5. Sisay B, Tefera T, Wakgari M, Ayalew G, Mendesil E. The efficacy of selected synthetic insecticides and botanicals against fall armyworm, *Spodoptera frugiperda*, in maize. *Insects*. 2019;10(45): 2-14.
6. Sharma H, Jaglan MS, Yadav SS. Population dynamics of pink stem borer, *Sesamia inferens* (Walker) on maize as influenced by weather conditions. *Journal of Applied and Natural Science*. 2017;9(4):1975-1980.
7. Iqbal J, Farooq SU, Alqarni AS, Ali H, Zeshan A, Ansari MJ. Management of maize stem borer (*Chilo partellus*) with insecticides of three formulations under field conditions. *Indian Journal of Agricultural Sciences*. 2017;87(12):1720-1724.
8. Rathod SN, Lande GK, Bhalkare SK, Sadawarte AK. Evaluation of Some Newer Insecticides Against Stem Borer, Aphids and Natural Fauna on Maize. 2020; 44(1):17-24. [www.PKVResearchJournal.ac.in](http://www.PKVResearchJournal.ac.in)
9. Arunkumara CG, Bheemanna M, Shaila HM. Field evaluation of insecticides for management of spotted stem borer, *Chilo partellus* (Swinhoe) (Lepidoptera: Crambidae) on maize. *Journal of Entomology and Zoology Studies*. 2017;5 (5):1719-1723.
10. Devananda KM, Khanpara AV, Vaja AM. Bio-efficacy of various insecticides against maize stem borer *Chilo partellus* (Swinhoe) Crambidae: Lepidoptera in Junagadh conditions. *Journal of Pharmacognosy and Phytochemistry*. 2018;7(4):2011-2014.
11. Prakash V, Singh DV, Singh R, Singh G, Kumar S. Efficacy of some novel insecticide against maize stem borer, *Chilo partellus* (Swinhoe) in maize. *Journal of Pharmacognosy and Phytochemistry*. 2017;SP1:481-484.
12. Joshi AR, Patel JR, Chaudhari SJ, Joshi MJ. Field efficacy of newer molecules against stem borer, *Chilo partellus* (Swinhoe) infesting maize. *International Journal of Chemical Studies*. 2019;7 (2):1588-1592.
13. Ahmed S, Saleem MA, Rauf I. Field efficacy of some bioinsecticides against Maize and Jowar stem borer, *Chilo partellus* (Pyralidae: Lepidoptera). *International Journal of Agriculture and Biology*. 2002;4(3):332-334.
14. Bhat ZH, Baba ZA. Efficacy of different insecticides against maize stem borer *Chilo partellus* (Swinhoe) and maize aphid *rhopalosiphum maidis* (Fitch) infesting maize. *Pakistan Entomological*. 2007;29 (2):73-76.
15. Dinesh MV, Simon S, Nagar S. Biorational management of stem borer, *Chilo partellus* in maize. *Journal of Pharmacognosy and Phytochemistry*. 2018;7(4):1142-1145.
16. Kumar GS, Madhumathi T, Kumar DS, Kumar VM, Ahamad M. Bioefficacy of different biorational insecticides for the management of spotted stem borer, *chilo partellus* (swinhoe) in maize (*Zea mays* L.). *Journal of Research ANGRAU*. 2017;45(2):19-30.

17. Kumar R, Jindal J. Economic evaluation of biorational and conventional insecticides for the control of maize stem borer *Chilo partellus* (Swinhoe) in *Zea mays*. Journal of Applied and Natural Science. 2015; 7(2):644-648.
18. Justin CGL, Preetha G. Survey on the occurrence, distribution pattern and management of stem borers on rice in Kanyakumari District, Tamil Nadu. Journal of Entomology and Zoology Studies. 2014;2(6):86-90.
19. Babu JS, Kumar A. Efficacy of certain chemicals and biopesticides against maize stem borer, *Chilo partellus* (Swinhoe). The Pharma Innovation Journal, SP. 2022;11 (7):4156-4159.
20. Reddy GV, Kumar A. Efficacy of various insecticides against maize stem borer *Chilo partellus* (Swinhoe) and their cost-benefit analysis at Prayagraj. Journal of Entomology and Zoology Studies. 2021; 9(6):130-134.
21. Malav SK, Lyall H, Sharma AK, Malav SSVMH. Efficacy of certain chemical insecticides and neem oil against stem borer (*Chilo partellus* Swin) on maize (*Zea mays* L.). International Journal of Advanced Scientific Research and Management. 2008;1:152-154.

© 2023 Chandar and Tayde; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/110371>