



Mapping of Soil Fertility Status in a Coastal Village of Odisha Using Geospatial Technology

Pradipta Majhi ^a, Prasannajit Mishra ^b, Jibanjit Sen ^a,
Dwarika Mohan Das ^a, Amit Phonglosa ^{b*}, Debasis Panda ^a,
Hemanta Kumar Sahoo ^b and Pravat Kumar Roul ^b

^a Krishi Vigyan Kendra, Jagatsinghpur, Odisha University of Agriculture and Technology,
Odisha 754160, India.

^b Directorate of Extension Education, Odisha University of Agriculture and Technology,
Bhubaneswar, Odisha 751003, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i234252

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/110265>

Original Research Article

Received: 10/10/2023
Accepted: 14/12/2023
Published: 21/12/2023

ABSTRACT

The most fundamental decision-making tool for sustainable soil nutrient management is the assessment of soil fertility in a region. During 2021, a soil fertility status inventory was prepared through National Innovations on Climate Resilient Agriculture (NICRA) project in Achyutadaspur village of Jagatsinghpur district, Odisha. A total of 125 surface soil samples were randomly collected from a depth of 0-15 cm with GPS locations and soil parameters like pH, electrical conductivity, organic carbon, available N, P, K, S, Zn, and B were evaluated using standard analytical procedures. GIS maps have been prepared using ArcGIS software based on the analysis report. The majority of the soil was found acidic around 62.4% area of the village, followed by

*Corresponding author: E-mail: soilamit12@rediffmail.com;

neutral soil pH in 33.6% area. The organic carbon status of the soil was found low in 79.2% area and medium in 20.8% area, available N was found low in 96% area and medium in 4% area, available P was found medium in 82.4% area and high in 17.6% area, available K was low in 27.2% area, medium in 64% area, and high in 8.8% area, and available S was found low in 37.6% area and medium in 62.4% area. Among the available micronutrients, Zn was found low in 79.2% area and medium in 20.8% area. Similarly, hot water extractable boron was found low in 92.6% area of the region and medium in 7.4% area. By adopting soil map based fertilizer recommendation with improved practices in the Achyutadaspur village, 17.92% yield was increased over farmers' practices. The benefit cost ratio (B:C ratio) of rice was found to be 1.89 whereas 1.65 in farmers practices.

Keywords: Fertility status; fertility maps; rice; yield; coastal area.

1. INTRODUCTION

Soil is critical factor for agriculture and it is the cradle for all crops and plants. It is the reservoir of nutrients that plays a crucial role for agricultural production [1]. In India, the soil resources accessible for agriculture are decreasing hence, optimized the exploitation of these resources is highly required for agricultural intensification. As a result, it is necessary to assess the fertility status of the soil on a regular basis in order to maintain the soil health [2]. Nutrient mining is an important threat in Indian agriculture since there is a large gap between nutrient addition and nutrient removal. In India, the nutrient loss rate is 10 million tonnes per year [3]. The one of the major reasons for reduced production is indiscriminate application of fertilizers without knowledge of soil fertility status and crop nutrient requirements, which has a negative impact on soil and crop in terms of nutrient toxicity and inadequacy [4]. Assessing the soil fertility status of a region is difficult to access and labour intensive due to presence of high spatial variability and heterogeneity [5]. Jagatsinghpur is coastal district of Odisha situated in the East and South Eastern Coastal Plain Agro Climatic Zone. Primary, secondary, and micronutrient deficiencies have been identified intensively in coastal areas. Farmers have been impelled to apply high amounts of chemical fertilizers due to rising food grain demand. Chemical fertilizer overuse is a severe danger to the sustainability of agricultural production. Soil test-based fertilizer recommendation and management is an excellent method for enhancing soil productivity. Achyutadaspur village is one of the coastal villages of Jagatsinghpur having vulnerability of water logging, soil salinity and flood. So, soil health management in such area is highly essential for sustainable crop production.

Therefore, this village is selected as adopted village under National Innovations on Climate Resilient Agriculture (NICRA) project of KVK Jagatsinghpur. Keeping this in view, GIS based soil nutrient mapping of Achyutadaspur village has been carried out and necessary advisory has been given to the farmers at each cadastral level.

2. MATERIALS AND METHODS

The study was conducted in NICRA adopted village (Achyutadaspur) of Jagatsinghpur district of Odisha. The district is situated in East and South East Coastal Plain Agro Climatic Zone of Odisha between 19°58' to 20°23' North Latitude and 86°03' to 86°45' East Longitude and total geographical area of district is 1668 square km and geographical area of the village is 204 ha, as shown in Fig.1.

Total 125 numbers of representative soil samples have been collected from the farmers field with the help of a soil auger for testing in KVK Jagatsinghpur soil testing laboratory. Surface soil samples within 0-15 cm depth were collected with geo-coordinates of the sampling location. The soil samples were mixed thoroughly and about 500g of composite soil samples were taken for analysis. First the samples were air-dried, passed through 2 mm and 0.5 mm sieve and stored in properly labeled plastic bags for physical and chemical analysis. The prepared samples were analyzed for using standard analytical methods. Soil pH, EC and organic carbon were estimated by Jackson method [6] [7]. Available nitrogen was determined following Subbiah and Asija method [8]. Available phosphorus and potassium were determined by Bray and Kurtz [9]; Olsen *et al.* [10]. Available sulphur was determined by the procedure described by Massoumi and Cornfield [11].




Fig. 1. Location map of study area

Available Zn and B were determined by [12] and [13], respectively. The method involved in analyses of soil samples is depicted in Table 1.

Based on soil test results, a front line demonstration (FLD) was conducted in rice crop during kharif-2022 in the selected village (Achyutadaspur) of Jagatsinghpur district. The farmers' practice was transplanting the rice variety Swarna in local check plot. The complete package of practice demonstrated in farmers' field through FLD including improved variety, recommended seed rate, seed treatment, sowing method, fertilizer dose, weed management

and plant protection which are presented in Table 11.

3. RESULTS AND DISCUSSION

3.1 Soil Reaction (pH)

The soil reaction of the soil samples were identified as acidic to neutral in condition with the value ranged from 5.50 to 6.50 with 6.08 as a mean value. The current findings showed that 62.4% area of the village is acidic reaction whereas 37.6% area is under neutral reaction (Table 2 and Fig.2).The results are being in conformity with the observations made by Majhi et al. [14].

Table 1. Analytical methodologies for soil analyses

Parameter	Methodology	References
pH	1:2.5:: Soil : Water	Jackson (1967) [6]
EC	1:2.5:: Soil : Water	Jackson (1967) [6]
Organic carbon	Wet oxidation method	Jackson (1973) [7]
Available N	Hot alkaline KMnO ₄ Method	Subbiah and Asija (1956) [8]
Available P	0.03 N NH ₄ F + 0.025 N HCL (pH 3.5) and 0.5 M NaHCO ₃ at pH 8.5	Bray and Kurtz (1945) [9] and Olsen et al. (1954) [10]
Available K	Neutral N NH ₄ OAc extraction	Jackson (1973) [7]
Available S	Extraction with 0.15% CaCl ₂	Massoumi and Cornfield, 1963 [11]
Available Zn	DTPA extractant	Lindsay and Norvel, 1978 [12]
Available B	Hot water extraction	Berger and Truog (1939) [13]

Table 2. pH status of the soil samples

Class	Rating	% of area
Acidic	<6.50	62.4
Neutral	6.50-7.50	37.6
Alkaline	>7.50	0




Fig. 2. Soil pH GIS Map

3.2 Electrical Conductivity

The electrical conductivity of the soil samples ranged from 0.86 to 2.68 dSm⁻¹, with a mean value 1.49 dSm⁻¹ (Table 3). The present findings observed that 33.6% of area of the village has normal salinity (suitable for all crop types) whereas 66.4% area of the village has critical EC (Fig. 3). Similar findings were also recorded by Majhi *et al.* [14] and Mishra *et al.*, 2015 [15]. The low EC might be due to free drainage conditions which favored the removal and release of bases by percolating and drainage water [16].

3.3 Organic Carbon

The organic carbon content of the soil samples ranged from 0.25 to 0.75% with a mean value of 0.47%. Based on the limits prescribed by Nanda *et al.*, 2008 [17], it is observed that 79.2% area of the village is under low in organic carbon content

whereas 20.8% area is within the medium carbon range (Table 4 and Fig. 4). It may be due to low application of FYM and crop residue as well as rapid rate of decomposition [18].

Table 3. Status of electrical conductivity in soil of the study area

Class	Rating (dSm ⁻¹)	% of Area
Low	<1.00	33.6
Medium	1.00-2.00	40.8
High	>2.00	25.6

Table 4. Status of organic carbon in soil of the study area

Class	Rating (%)	% of Area
Low	<0.50	79.2
Medium	0.50-0.75	20.8
High	>0.75	0




Fig. 3. Soil electrical conductivity (dSm^{-1}) GIS Map




Fig. 4. Soil organic carbon (%) GIS Map

3.4 Available Nitrogen

The soil sample analysis revealed that the available nitrogen content ranged from 168 to 274 kg ha^{-1} , with a mean value of 229.48 kg ha^{-1}

(Table 5 and Fig. 5). On the basis of prescribed limits by Nanda *et al.*, [17], the result showed that 96% soil of area has low in nitrogen content and 4% soil has medium available nitrogen status. It may be due to the nitrogen lost through various

mechanism like ammonia volatilization, nitrification, chemical and microbial fixation, leaching, runoff and may be due to very low content of organic carbon [19].

Table 5. Status of available nitrogen in soil of the study area

Class	Rating (kg ha ⁻¹)	% of Area
Low	<250	96
Medium	250-500	4
High	>500	0

3.5 Available Phosphorus

The available phosphorus content of the area ranged from 9 to 35 kg ha⁻¹, with a mean value of 17.57 kg ha⁻¹ (Table 6 and Fig.6). Based on the limits prescribed by Nanda *et al.* [17], the present results revealed that 82.4% area of the village has medium range of available phosphorus whereas 17.6% area has high range of available phosphorus. It might be due to slightly acidic pH of soil which is making available phosphorus immobile [20].

Table 6. Status of available phosphorus in soil of the study area

Class	Rating (kg ha ⁻¹)	% of Area
Low	<9.0	0
Medium	9.0-22.0	82.4
High	>22.0	17.6

3.6 Available Potassium

The available soil potassium content varied from between 118 to 280 kg ha⁻¹, with a mean value of 201.36 kg ha⁻¹. Based on the limits prescribed by Nanda *et al.* [17], the present results showed that 27.2% area of the village has low range of potassium whereas 64% and 8.8% area has medium and high range of available phosphorus, respectively (Table 7 and Fig. 7).

Table 7. Status of available potassium in soil of the study area

Class	Rating (kg ha ⁻¹)	% of Area
Low	<118	27.2
Medium	118-280	64
High	>280	8.8

3.7 Available Sulphur

The available sulphur status of the soil samples ranged between 5.18 to 20.00 mg kg⁻¹, with a mean value 10.08 mg kg⁻¹ (Table 8 and Fig. 8). Results revealed that 62.4% area of the village has medium range of available sulphur and 37.6% area of the village has low range of available sulphur content. Deficiency of soil S might be due to the presence of variable properties of different components of organic matter, improper fertilizer and water management practices [21].




Fig. 5. Soil available N (kg ha⁻¹) GIS Map




Fig. 6. Soil available P (kg ha⁻¹) GIS Map




Fig. 7. Soil available K (kg ha⁻¹) GIS Map

3.8 Available Zinc

The available zinc content of the soil samples varied from 0.3 to 1.2 mg kg⁻¹, with a mean value 0.46 mg kg⁻¹. Results revealed that 79.2% area of the village has low range of available Zn whereas 20.8% area has medium range of available Zn (Table 9 and Fig. 9). This revealed

that majority of soil in the village is deficient in Zn.

3.9 Available Boron

The hot water extractable B content of the soil samples varied from 0.23 to 0.74 mg kg⁻¹, with a mean value of 0.40 mg kg⁻¹ (Table 10 and Fig.

10). The present results showed that 92.6% area of the village has low range of available B content whereas 7.4% area of the village has medium range of hot water extractable B content.

Table 8. Status of available sulphur in soil of the study area

Class	Rating (mg kg ⁻¹)	% of Area
Low	<10	37.6
Medium	10-20	62.4
High	>20	0

3.10 Yield Performance and Economic Analysis of Demonstration and Farmers' Practices during Kharif Season of 2022

The yield data indicated that due to frontline demonstration, rice yield was found 48.78 q ha⁻¹ in demonstration plot and 41.34 q ha⁻¹ in farmers' practice plot during *kharif* 2022 (Table 11). The demonstration results also revealed that 17.92% yield was increased over farmers' practice in rice (Table 11). This result clearly indicated the

higher yield in demonstration plots over farmers' practice was due to adoption of improved package of practices following soil test results of the farmer's field. Similar findings in rice and others crops have been reported by Mishra et al. [22], Pandey et al. [23] and Phonglosa et al. [24].

Rice cultivation under improved practices has shown higher net return of Indian Rupee (INR) 46,850 ha⁻¹ as compared to farmers' practices with INR 33,434 ha⁻¹ during *kharif* season of 2022. The benefit cost ratio (B:C ratio) of rice under improved practices was found to be 1.89 as compared to 1.65 under farmers' practices (Table 12). Based on the cost of production and the current input and output prices, the economic feasibility of the improved technology over the farmers' practice was found superior and was proved though higher B:C ratio. The increased cost of demonstration was mostly attributed by the higher costs of pesticides, fertilizer and seeds. These results are also in conformity with the findings of Girish et al. [23] and Phonglosa et al. [24] [27].




Fig. 8. Soil available S (mg kg⁻¹) GIS Map

Table 9. Status of available zinc in soil of the study area

Class	Rating (mg kg ⁻¹)	% of Area
Low	<0.60	79.2
Medium	0.60-1.20	20.8
High	>1.20	0

Table 10. Status available boron in soil of the study area

Class	Rating (mg kg^{-1})	% of Area
Low	<0.50	92.6
Medium	0.50-0.75	7.4
High	>0.75	0




Fig. 9. Soil available Zn (mg kg^{-1}) GIS Map




Fig. 10. Soil available B (mg kg^{-1}) GIS Map

Table 11. Improved practices and farmers practices of rice under frontline demonstrations

Sl. No.	Particulars	Improved practices for demonstration	Existing Farmer practices
1	Variety	Swarna sub-1	Pooja, Swarna
2	Field preparation	Ploughing, Harrowing and puddling	Ploughing, Harrowing and puddling
3	Seed rate	50 kg ha ⁻¹	70 kg ha ⁻¹
4	Seed treatment	Tricyclazole 75WP @ 2g kg ⁻¹ of seed	No seed treatment
5	Nursery management and transplanting	Nursery seeding is done during May to June and transplanting is done during July	Nursery seeding is done during May to June and transplanting is done during July
6	Sowing method	Line transplanting	Random transplanting
7	Fertilizer dose	Soil test based NPK @ 100:40:40 kg ha ⁻¹ + Zn @ 5 kg ha ⁻¹ + B @ 1 kg kg ha ⁻¹ + well decomposed FYM @ 5 t ha ⁻¹	Imbalance use of fertilizer
8	Weed management	Bispyribac Sodium 10% + Clomazone 48% @ 0.25 litre ha ⁻¹	Butachlor @ 2.0 litre ha ⁻¹
9	Plant protection	Need based plant protection measures	Injudicious use of plant protection chemicals

Table 12. Yield performance of demonstration and farmers' practices during *Kharif* season of 2022

Name of crop and variety	Improved yield (q ha ⁻¹)	Farmers practice (q ha ⁻¹)	Yield increase over FP (%)
Rice, Swarna Sub-1	48.75	41.34	17.92

Table 13. Economic analysis of demonstration and farmers' practices during *Kharif* season of 2022

Improved practice				Farmers Practice			
Cost of cultivation (INR ha ⁻¹)	Gross Return (INR ha ⁻¹)	Net Income (INR ha ⁻¹)	B:C ratio	Cost of cultivation (INR ha ⁻¹)	Gross Return (INR ha ⁻¹)	Net Income (INR ha ⁻¹)	B:C ratio
52600	99450	46850	1.89	50900	84334	33434	1.65

INR=Indian Rupee; B:C=Benefit cost ratio

4. CONCLUSION

The collected soil samples on grid basis were tested in KVK soil laboratory. Soil pH, EC, OC, N, P, K, some secondary nutrient like S and micro nutrients like Zn and B were quantified using various standard methodologies in the soil laboratory. Then the gridded data were converted into thematic layers and interpolated using Kriging Technique in ArcGIS software to develop soil nutrient map of the district. Further, the nutrients maps were classified into high, medium and low zones to identify the nutrient deficient and sufficient zones. In this context, soil was found acidic around 62.4% area of the village, followed by neutral soil pH in 33.6% area. The organic carbon status of soil was found low

in 79.2% area to medium in 20.8% area, available N was found low in 88% area and medium in 12% area, available P was found medium in 82.4% area and high in 17.6% area, available K was low in 27.2% area, medium in 64% area and high in 8.8% area and available S was low in 37.6% area and medium in 62.4% area, respectively. Among the micronutrients, available Zn was found low in 79.2% area and medium in 20.8% area of the village. Likewise hot water extractable boron was found low in 92.6% area and medium in 7.4% area. Using the GIS map, cadastral level soil nutrient status was estimated easily using GIS which will be much helpful for farmers for getting proper fertilizer recommendation. Thus, the GIS based soil fertilizer recommendation is now specific to a

particular farmer not to a group of farmers in a patch. It was also observed that production of rice in the Achyutadaspur village has been enhanced by adopting GIS soil map based fertilizer recommendation along with improved package of practices.

5. RECOMMENDATION

The soil spatial mapping of the village in GIS revealed that the majority of the soil in the village is acidic, so farmers are recommended to do various improved management practices like green manuring, lime and other soil amendments application to neutralize the soil pH. Around 25.6% area in the village is now affected by soil salinity, hence ponding of fresh rain water in bonded lands are recommended for leaching the salt from root zone. In addition saline tolerant local rice varieties *i.e.*, Luna Suvarna, Luna Barihal, Lunisri, Luna Ambiki are recommended in high saline fields. Organic carbon is low in most of the area which is associated with low available nitrogen. Hence, farmers are recommended to apply FYM and vermi-compost along with 25% higher dose of nitrogen above the recommended dose. In majority of area available phosphorus and potassium is medium, hence recommended fertilizer dose should be adopted in those patches. The secondary nutrient sulphur is medium in majority of the area and low in around 38% area. In low sulphur area farmers are advised to apply organic manure along with sulphur based fertilizers and amendments. Among the micro-nutrients, the majority of the land is deficient with zinc and boron. Khaira disease in rice due to zinc deficiency is observed in the village and fruit cracking is observed in areca nut, coconut, water melon etc. due to boron deficiency. Therefore, farmers are advised to apply zinc and boron based fertilizer in soil or as foliar spray as per soil test based recommendations.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Sahu DK, Samadhiya VK, Chandrakar T, Netam S, Sahu M, Homeshvari. Macro and micro nutrient status of research farm, college of agriculture and research station, Kurud, district Dhamtari, Chhattisgarh. The Pharma Innovation International Journal. 2021;8(10):32-35.
2. Sashikala G, Naidu MVS, Ramana KV, Nagamadhuri KV, Reddy A, Kumar P, Sudhakar P, Krishna T. Giridhara. Mapping of nutrients status in Tatrakallu village of Anantapuramu district of Andhra Pradesh using geographic information system. Journal of the Indian Society of Soil Science; 2021 69(2):133-141.
3. Ramakrishna PVR, Munawery A. Sustainable soils nutrient management. Journal of the Indian Institute of Science. 2012;92:1-8.
4. Ray PK, Jana AK, Maitra DN, Saha MN, Chaudhury J, Saha S, Saha AR. Fertilizer prescriptions on soil test basis for jute, rice and wheat in Typic Ustochrept. Journal of the Indian Society of Soil Science.2000;48:79-84.
5. Singh RP, Mishra SK. Available macro nutrients (N, P, K and S) in the soils of Chiraigaon block of district Varanasi (U.P.) in relation to soil characteristics. Indian Journal Science Research. 2012;3(1): 97-100.
6. Jackson ML. Soil Chemical Analysis, Prentice Hall of India Pvt. Ltd., 1967 New Delhi. 1967;205.
7. Jackson ML. Soil Chemical Analysis. Prentice Hall Inc. Englewood Cliffs, 1973 New Jersey, USA; 1973.
8. Subbiah BV, Asija CL. A rapid procedure for the determination of available nitrogen in soils. Current Science. 1956;25:259-262.
9. Bray RH, Kurtz LT. Determinations of total, organic and available forms of phosphorus in soils. Soil Science. 1945;59:39-45.
10. Olsen SR, Cole CV, Watanable FS, Dean LA. Estimation of available phosphorus in soils by extraction with sodium bicarbonate USDA Cric. 1954;939.
11. Massoumi A and Cornfield AH. A rapid method for determining sulphate in water extracts of soils. Analyst. 1963;88(1045):321-322.
12. Lindsay WL, Norvell WA. Development of a DTPA soil test for zinc, Iron, Manganese, and Copper. Soil Science Society of America Journal. 1978;42:421-428.
13. Berger KC, Truog E. Boron determination in soils and plants using the quinalizarin

- reaction. Indus. Eng. Chem. 1939;11:540–545.
14. Majhi P, Phonglosa A., Mohanty SK, Roul PK. Soil Fertility Status of Different Blocks of Puri District of Coastal Odisha. International Journal of Agriculture Sciences. 2018;21(10):7445-7447.
 15. Mishra A, Das D, Saren S, and Dey P. GPS, GIS Based Soil Fertility Maps of Puri District of Coastal Odisha. Indian Society of Coastal Agricultural Research. 2015;33(2):7-11.
 16. Seema, Ghosh AK. Characterization and classification of alluvium derived soils under different land uses in Varanasi district of Uttar Pradesh. Journal of the Indian Society of Soil Science. 2019;67(3): 360-364.
 17. Nanda SK, Mishra A, Pradhan NK, Muralidharudu Y. Soil testing and fertilizer recommendation in Orissa, STCR technical bulletin, OUAT, Bhubaneswar; 2008.
 18. Sathish A, Ramachandrappa BK, Devaraja K, Savitha MS, Gowda MN, Prashanth K.M. Assessment of spatial variability in fertility status and nutrient recommendation in Alanatha Cluster Villages, Ramanagara District Karnataka using GIS. Journal of the Indian Society of Soil Science. 2017;66(2):149-157.
 19. Vaisnow B, Sengar SS, Jatav GK, Patel T, Bhagat RK. Soil fertility status of major nutrient in vertisol of Dhamtari block. Journal of Plant Development Sciences. 2014;6(4):587-591.
 20. Krishna Bal, Sahu KK, Tedia K, Singh CB, Kumar R. Evaluation of Soil Fertility Status of Palari Block under Baloda Bazar District of Chhattisgarh. Journal of Pharmacognosy and Phytochemistry. 2018;SP2:243-246.
 21. Sutar RK, Pujar AM, Kumar BNA, Hebsur NS. Sulphur nutrition in maize- A critical review. International Journal of Pure & Applied Bioscience. 2017; 5(6): 1582-1596.
 22. Mishra PK, Singh VP, Singh SN, Kumar P, Pandey MK. Impact of frontline demonstration in adoption extent and horizontal spread of tomato (*Lycopersicon esculentum* Mill.) cultivation in Tarai region of Siddharthnagar district, Uttar Pradesh, India. Journal of pharmacognosy and Phytochemistry. 2019;8(3):4024-4028.
 23. Pandey SK, Tiwari DK, Singh S, Singh P. Promotion of long duration rice variety Swarna sub-1 through frontline demonstration in Chandauli district of Uttar Pradesh, India. International Journal of Current Microbiology and Applied Sciences. 2018;7(5):2870-2874.
 24. Phonglosa A, Sahoo HK, Mishra P, Sangramsingh SP, Patra B. Performance of Submergence Tolerant Shallow Lowland Rice Variety Swarna sub-1 under Frontline Demonstrations in East and South Eastern Coastal Plain Zone of Odisha, India. International Journal of Plant & Soil Science. 2022;34(20):17-23.
 25. Girish R, Bharath Kumar TP, Shruthi HR, Shivakumar L, Praveen KM. Frontline demonstration on paddy variety KPR 1 by KVK in Chikkamagaluru district of Karnataka, India: An impact study. Journal of Pharmacognosy and Phytochemistry. 2020;9(2):303-305.
 26. Priyadarshini PR, Saren S, Mishra A, Acharya BP. Soil fertility status of some villages under North-Eastern Coastal Plain Agro climatic Zone of Odisha. Journal of Indian Society of Coastal Agricultural Research. 2017;35(2):42-47.
 27. Phonglosa A, Dalei BB, Murmu P, Behera B, Sinha B, Saren S, Nayak L, Meena MK. Integrated nutrient management in rainfed maize (*Zea mays* L.) under Eastern Ghat High Land Zone of Odisha. International Journal of Plant & Soil Science. 2022;34(6):102-108.

© 2023 Majhi et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/110265>