



Estimation of Genetic Variability Parameters for Various Yield Attributing Traits in Germplasm Collection of Rice (*Oryza sativa* L.)

Abhinav Sao ^{a*}, Guman Singh ^a, S. K. Nair ^a
and Harish G. D. ^b

^a Department of Genetics and Plant Breeding, Indira Gandhi Krishi Vishwavidyalaya Raipur 492012, Chhattisgarh, India.

^b Germplasm Conservation Division, ICAR-National Bureau of Plant Genetic Resource, New Delhi, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2024/v36i34411

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/112515>

Original Research Article

Received: 24/11/2023

Accepted: 27/01/2024

Published: 07/02/2024

ABSTRACT

Genetic variability parameters were studied in 2879 germplasm accessions of rice for 11 quantitative traits. Moderate value of genotypic coefficient of variation (GCV) and phenotypic coefficient variation (PCV) was observed for the traits, 100 seed weight, seedling height, L:B ratio, plant height, and number of effective tillers, whereas rest of the traits had low GCV and PCV values. High heritability was observed for all the traits except leaf length, leaf width and grain yield per plant. High heritability coupled with high genetic advance as percent of mean was observed for 100 seed weight, seedling height and L:B ratio. High heritability coupled with moderate genetic advance as percent of mean was recorded for days to 50% flowering and plant height. The presence of high heritability and high genetic advance as percent of the mean indicated the presence of additive gene effects for controlling aforesaid traits, hence selection in later generations will be rewarding in a breeding programme.

*Corresponding author: E-mail: saoabhi27@yahoo.co.in;

Keywords: Rice germplasm; heritability; genetic advance; phenotypic coefficient of variation (PCV) and genotypic coefficient of variation (GCV).

1. INTRODUCTION

Rice is the most consumed cereal grain in the world, constituting the dietary staple food for more than half of the planet's human population. India being the second largest producer of rice still lags behind its productivity per unit area. The rising demand, saturation of cultivable field and low gross domestic production of rice are likely to cause a supply shortage of a crop in near future. By the year 2025, about 785 million tons of paddy, which is 70 per cent more than the current production will be needed to meet the growing demand [1]. This ranks first among the three major cereals, followed by wheat and maize. It is one of the oldest cultivated sources of nutrients to flourish under a wide range of environments and it has 22 wild species in the *Oryza* genus. It may be cultivated in a variety of environments, including upland, lowland, and rain-fed. It is one among India's most important crops. Moreover, it is a nutrient-rich cereal crop that provides 20 % of the calories and 15 % of the protein consumed by the world's population.

Crop improvement is heavily dependent on understanding the genetic diversity and population structure of germplasm collections. Because rice is an important crop in the globe, there is a lot of curiosity in the origins and diversity of *Oryza sativa*. Current theories suggest that domesticated rice originated from a polyphyletic source, most likely from either an annual or perennial variety of *O. rufipogon*, through at least two separate domestication processes [2,3,4].

Minerals and fibers are also present in the rice grains. Collection and classification of existing germplasm is critical not only for using appropriate attribute-based donors in breeding programmes, but also for protecting the distinctive rice for the future. In Asia, rice production contributes to almost 90 % of global output [5]. Estimating genetic gain under selection will be more accurate when heritability and evolution are combined [6]. To enhance rice production, many programmes and activities were developed. However, the successes continue to be up to par, which has prompted the country to shift from being a net rice producer to an import in recent years [7,8]. Crop failure in recent years has been mainly

caused by climate change scenarios such as unpredictable rainfall, disease and pest attack, and genetic degradation owing to developmental activities [9].

The characterization and evaluation of conserved genetic resources in gene banks are very essential to document the presence of genetic variability and diversity for different important traits. Keeping in view the importance of variability, the present study has been conducted to estimate the variability parameters in large number of germplasm accessions.

2. MATERIALS AND METHODS

The materials for the study comprised of 2879 germplasm accessions of rice i.e. 2879 germplasm accessions received from ICAR-NBPGR (National Bureau of Plant Genetic Resources), New Delhi including 6 checks namely, Annada, IR 64, Pusa Basmati-1, Swarna, NDR-97 and Jaya of different duration groups. The experimental materials were grown in *Kharif*, 2022 in Augmented Block Design. Twenty one days old seedlings of each genotype were transplanted manually in 2 row of 2-meter length maintaining a spacing of 20 cm between rows and 15 cm between plants in each row. Observations were recorded on 3 randomly selected plants from each genotype for 11 quantitative traits viz., days to 50% flowering, no. of effective tillers, days to maturity, panicle length, plant height, 100 grain weight, seedling height, leaf length, leaf width, grain yield per plant and grain length/breadth ratio. The mean values of these were used for statistical analysis through XL Stat.

3. RESULTS AND DISCUSSION

The experiment was carried out in Augmented Block Design with the objective to assess the variability present in the rice accessions. In almost all the traits evaluated in the experiment, the phenotypic coefficient of variation (PCV) was greater in magnitude than the genotypic coefficient of variation (GCV). This indicated that the apparent variance is attributable not just by genes but also due to environmental effects. The mention of a slight distinction between GCV and PCV was also reported by Mustafa and Isheikh [10], Kole et al. [11] and Mulugeta Syoum et al. [12]. Classifying the overall variation into its

constituent parts could facilitate the preservation and application of genetic resources. Also, it provides the possibility to use appropriate germplasm for crop enhancement of certain plant features [13].

The phenotypic coefficient of variation (PCV) with higher magnitude is presented in Table No.1. Higher phenotypic coefficient of variation (PCV) was recorded for the traits grain yield per plant (37.43), similar to the findings of Lingaiah et al. [14]. Moderate phenotypic coefficient of variation was observed for the traits, seedling height (16.34), L/B ratio (15.65), leaf width (14.12), leaf length (13.18), plant height (12.04), 100 seed weight (19.4) and number of effective tillers (19.39) and rest of the traits had low PCV. The lowest GCV was recorded in days to 50% flowering (9.35), days to maturity (7.3) and panicle length (5.81). The results are in agreement with Bisen et al. [15], Singh et al. [16], Subbaiah et al. [17] and Babu et al. [18]. Moderate GCV was recorded for 100 seed weight (17.93), seedling height (16.34), L/B ratio (14.05), plant height (10.98) and the number of effective tillers (11.27). Prajapati et al. [19] and Ananadarao et al. [20] also reported similar observations in their study.

The presence of higher genotypic coefficient of variation (GCV) suggests that the population has a considerable genetic diversity. The phenotypic coefficient of variation was higher than the GCV in the present research, but the difference was moderate, indicating that environment impact on traits expression was low. Similar findings has also been reported by Sarawgi et al. [21], Ganesan et al. [22] and Sharma and Dubey [23].


To identify a trait's response to phenotypic selection, heritability is crucial. For a breeding programme to be successful, improving quantitative traits in plants requires accurate heritability estimation [24]. The estimates of genetic advance expressed as a percentage of the mean offer greater confidence about the efficacy of selection in enhancing the traits. Genetic progress indicates the new population's genotypic value is higher than that of the original population. Heritability estimates alone are less useful in predicting the gain under selection than heritability estimates combined with significant genetic advancement.

However, it's not always the case that a trait with high heritability also has strong genetic advances [25]. High heritability was recorded for seedling height (100), days to 50 % flowering (100), days to maturity (100), 100 seed weight (85.45), plant height (83.23) and L/B ratio (80.68). Li-Yun et al. [26] and Satyanaryan et al. (2005) also reported similar findings. Genetic advance as percent of mean was higher for the 100 seed weight (33.72) and L/B ratio (26.04). High heritability coupled with higher genetic advance as percent of the mean were observed for 100 seed weight, seedling height and L/B ratio. High heritability coupled with moderate genetic advance as percent of mean was recorded for plant height and days to maturity and rest of the traits are low heritability and genetic advance as percent of mean, the findings were also in agreement with Kujur et al. [27]. The result shows the presence of additive gene effects. Hence, selection of desirable genotype for future breeding programme will be rewarding [28,29].


Table 1. Estimation of genetic variability parameters for quantitative traits

S. No.	Traits	Mean	Range		PCV (%)	GCV (%)	h ² bs (%)	GAM
			Min.	Max.				
1.	Leaf length (cm)	55.04	27.16	2.06	13.18	12.16	85.07	23.13
2.	Leaf width (cm)	1.09	0.6	2.06	14.12	11.28	63.07	18.57
3.	Days to 50% flowering	106.59	65.00	148.00	9.35	9.35	99.79	19.29
4.	Plant height (cm)	150.48	52.00	210.33	12.04	10.98	83.23	20.67
5.	Panicle length (cm)	25.10	9.10	32.63	9.15	5.81	40.29	7.61
6.	No. of effective tillers	6.53	1.33	13.33	19.39	11.27	33.76	13.51
7.	Days to maturity	136.59	95.00	178.00	7.30	7.30	99.79	15.05
8.	L/B ratio	3.23	1.70	5.40	15.65	14.05	80.68	26.04
9.	100 seed weight (g)	2.36	0.77	4.90	19.40	17.93	85.45	34.02
10.	Grain yield per plant (g)	19.86	2.00	116.00	37.43	35.20	82.03	32.03


Where, Min: minimum, Max: maximum, PCV: phenotypic coefficient of variation, GCV: genotypic coefficient of variation, h²bs: heritability (broad sense), GA: genetic advance, GAM: genetic advance as percent of mean




Seedling height




Plant height




Days to 50% flowering




Panicle length




No. of effective tillers



Days to maturity



100 grain weight



Grain L/B ratio




Fig. 1. Graphical representation of variability parameters for different quantitative characters

4. CONCLUSION

The present study aimed at the assessment of the extent of variability present in the germplasm accessions. The PCV was slightly higher than the GCV for most of the characters, indicating a little bit of influence of environment on the expression of traits. High variability was observed for grain yield per plant and moderate for other traits means ample diversity is present in the germplasm accessions. High heritability coupled with high genetic advance was also observed for flowering, maturity duration and yield traits indicating the selection for these traits will also be passed on to the next generation, hence genotypes are useful for future breeding programmes.

ACKNOWLEDGEMENT

The authors are highly thankful to ICAR- National Bureau of Plant Genetic Resources, New Delhi for funding support under the project Consortium Research Platform on Agro-Biodiversity (CRP-AB) and supply of rice Germplasm Accessions for the study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Manomani S, Fazlullah Khan AK. Studies on combining ability and heterosis in rice. Madras Agric. J. 2003;90: 228-231.
2. Second G. Origin of the genetic diversity of cultivated rice (*Oryza spp.*): Study of the polymorphism scored at 40 isozyme loci. Jpn. J. Genet.1982;57:25–57
3. Cheng CY, Motohashi R, Tsuchimoto S, Fukuta Y, Ohtsubo H, Ohtsubo E. Polyphyletic origin of cultivated rice: based on the interspersed pattern of SINEs. Mol. Biol. Evol. 2003;20:67–75.
4. Vitte C, Ishii T, Lamy F, Brar DS, Panaud O. Genomic paleontology provides evidence for two distinct origins of Asian rice (*Oryza sativa* L.). Mol. Genet. Genomics. 2004;272:504–511.
5. Faysal ASM, Ali L, Aza MG, Sarker U, Ercisli S. Genetic variability, character association, and path coefficient analysis in transplant Aman rice genotypes. Plants. 2022;11(21):2952.
6. Demeke B, Dejene T, Abebe D. Genetic variability, heritability, and genetic advance of morphological, yield related and quality traits in upland rice (*Oryza sativa* L.) genotypes at pawe, northwestern Ethiopia. Cogent Food & Agriculture. 2023;9(1): 2157099.
7. Joshi NP, Maharjan KL, Piya L. Production economics of rice in different development regions of Nepal. J. Int. Dev. Coop .17.1. 2011:103-112. Online.
8. Bishwajit G, Sarker S, Kpoghomou MA, Gao H, Jun L, Yin D, Ghosh S. Self-sufficiency in rice and food security: A South Asian perspective. Agri. & Food Secu. 2013;2.1: 10.
9. Ali S, Liu Y, Ishaq M, Shah T, Abdullah, Ilyas A, Din IU. Climate Change and Its Impact on the Yield of Major Food Crops: Evidence from Pakistan. Foods.2017;6 (6):39.
10. Mustafa MA, Elsheikh MAY. Variability, correlation and path coefficient analysis for yield and yield components in rice. Afri. Crop. Sci. J. 2007;15:183-189.
11. Kole PC, Chakraborty NR, Bhatt JS. Analysis of variability, correlation, path coefficients in induced mutants of aromatic non basmati rice. Trop. Agric. Res. Exten. 2008;113: 60-64.
12. Seyoum M, Sentayehu A, Kassahum B. Genetic variability, heritability, correlation coefficient and path analysis for yield and yield related traits in upland rice. J. Plant Sciences. 2012;7(1):1320.
13. Pecetti L, Annicchiaro P, Damania AB. Geographic variation in tetraploid wheat (*Triticum turgidum* spp. *Turgidum convar. Durum*) landraces from two provinces in Ethiopia. Genetic Resources and Crop Evolution. 1996;43:395-407.
14. Lingaiah N, Venkanna V, Cheralu C. Genetic variability analysis in rice (*Oryza sativa* L.). Int. J. Pure App. Biosci. 2014; 2(5):203-204.
15. Bisen R, Sarawgi AK, Verulkar SB. Study of heritability, genetic advance and variability for yield contributing characters in rice. Bangladesh J. Agric. Res. 2009;34(2):175-179.
16. Singh SK, Singh CM, Gil GM. Assessment of genetic variability for yield and its component characters in rice. Research in Plant Biology. 2011;1(4):73-76.

17. Subbaiah PV, Sekhar MR, Reddy KHP, Reddy NPE. Variability and genetic parameters for grain yield and its components and kernel quality attributes in CMS based rice hybrids (*Oryza sativa* L.). International Journal of Applied Biology and Pharmaceutical Technology. 2011;2(3):603-609.
18. Babu VR, Shreya K, Dangi KS, Usharani G, Nagesh P. Genetic variability studies for qualitative and quantitative traits in popular rice (*Oryza sativa* L.) hybrids of India. International Journal of Scientific and Research Publication. 2012;2(6):1-5.
19. Prajapati M, Singh CM, Suresh BG, Lavanya GR, Jadhav P. Genetic parameters for grain yield and its component characters in rice. Ele. J. Pl. Bred. 2011;2(2):235-238.
20. Ananadrao SD, Singh CM, Suresh BG, Lavanya GR. Evaluation of rice hybrids for yield and yield component characters under North East Plain Zone. The Allahabad farmer. 2011;67(1): 63-68.
21. Sarawgi AK; Soni DK, Shrivastava MN. Variability analysis in rice under irrigated and rain-fed situations. Current Research. 1994;23(3/4):33-35.
22. Ganesan K, Manuel WW, Sundram T. Analysis in Rice. Int Rice Res. Newl. 1995; 20:1- 4.
23. Sharma RK, Dubey SD. Variation and association among panicle traits in rice. *Oryza*, 1997;34:8-12.
24. Saidon SA, Kamaruzaman R, Razak MSFA, Ramli A, Sarif HM. Studies on heritability and genetic variability for grain physical properties in Malaysian rice germplasm. Earth Environ. Sci., 2020;482.
25. Johnson HW, Robinson HF, Comstock RE. Estimates of genetic and environmental variability in soybean. Agron. J. 1955; 47:314-318.
26. Li Y, Zhao GL, Zuo YS, Zhang CW. Studies on the combining ability of head rice milled rate under different environmental condition in three-line *Indica* hybrid rice. Southwest China J. Agric. Sciences. 2003;16(2):22-25.
27. Kujur, Vivek Kumar, Sao, Abhinav, Singh MK, Tiwari, Ajay. Genetic variability, heritability and association analysis for yield and related characters in rice germplasm (*Oryza sativa* L). The Pharma Innov. J. 2023;12 (4):2236-2240
28. Nazir A, Farooq J, Mahmood A, Shahid M, Riaz M. Estimation of genetic diversity for CLCuV, earliness and fiber quality traits using various statistical procedures in different crosses of *Gossypium hirsutum* L. Vestnik Orel Gau. 2013;43:4: 2-9.
29. Satyanaryan PV, Srinivas T, Reddy PR, Madhavalatha L, Suneetha Y. Studies on variability, correlation and path coefficient analysis for restorer lines in rice (*Oryza sativa* L.). Res Crops. 2005;6:80-84.

© 2024 Sao et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/112515>