



Edible Flowers: Super Foods with Potential Health Benefits

**Yachna Sood ^a, Manmohan Lal ^{a*}, Ayush Kalia ^a
and Sakshi Verma ^a**

^a *University Institute of Agricultural Sciences, Chandigarh University, India.*

Authors' contributions

This work was carried out in collaboration among all authors. Author YS designed the study and wrote the first draft of the manuscript. Author ML and AK managed the analyses the study and prepared the final draft. Author AK and SV managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2024/v36i34417

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

<https://www.sdiarticle5.com/review-history/112264>

Review Article

Received: 16/11/2023

Accepted: 20/01/2024

Published: 09/02/2024

ABSTRACT

Edible flowers have become popular in recent years as a unique and visually tempting addition to culinary creations. This review article explores the assorted world of edible flowers, focusing on their potential health benefits, bioactive compounds, and associated toxicological risks, with particular attention to the anti-nutritional compounds and toxic flowers. Edible flowers from different plant species provide a variety of bioactive compounds that contribute to their flavor, aroma, and potential health benefits. These bioactive compounds, including antioxidants, vitamins, minerals and phytochemicals, make edible flowers not only beautiful but also nutritious. Incorporating them into your diet can improve the overall nutritional content of your meals. However, consuming edible flowers is not without risks. Some flowers contain anti-nutrient compounds, such as oxalates and phytates, which can interfere with the absorption of essential minerals and lead to nutritional deficiencies over time. Therefore, it is important to practice moderation when consuming certain edible flowers, especially for those with pre-existing medical conditions or prone to mineral imbalances. However, it is essential to pay attention to anti-nutritional compounds that can

*Corresponding author: E-mail: manmohanhort@gmail.com;

affect mineral absorption and the potential toxicity of certain flowers. When used judiciously and safely, edible flowers can be a delicious and healthy addition to a balanced diet. Careful selection and proper preparation are essential to reap the benefits while minimizing the risks associated with these beautiful flowers. This review serves as a valuable resource for consumers, chefs and nutritionists, providing insight into the world of edible flowers and their bioactive compounds as well as considerations regarding anti-nutritional compounds and their toxicity.

Keywords: Edible flowers; bioactive compounds; anti-nutritional compounds; toxicity.

1. INTRODUCTION

Edible flowers are the flowers of any plant that are generally safe for humans. These flowers are becoming increasingly popular, as evidenced by the increasing number of edible flowers used in cookbooks, cooking magazine articles, and political television shows. Many flowers play a prominent role in food preparation, adding fragrance, flavor and aesthetic value, such as roses (*Rosa* spp. L.), which were used in ancient Rome to increase the flavor and sweetness of foods in drinks, salads, purees, desserts, etc. People in China, the Middle East, and India also use this flower because it originated in ancient civilizations [1]. Today, sales of fresh flowers for human consumption are increasing around the world. In addition, these products, suitably packaged in bags, boxes, etc., will be sold directly in farm stores or through various specialized sales outlets [2]. Curiosity about edible flowers is a result of the search for new natural food products. Consumers are increasingly choosing food products containing natural ingredients due to concerns about the negative health effects of synthetic compounds. However, processing and preservation aspects have only been explored for a few edible flowers and could be applied more widely. The commercialization of value-added products from this flower is still very far away due to the lack of scientific reports. The potential of these flowers as superfoods can also be explored.

2. EDIBLE FLOWERS

2.1 Edible Flowers as a Source of Bioactive Substances

Edible flowers typically contain 70-95% water, petals, nectar, and pollen, making them an important source of metabolites, vitamins, and minerals. Edible flowers are also rich in antioxidant molecules (such as polyphenols and pigments) that can help prevent certain diseases. Edible flowers are important for human health as they are rich in nutrients and bioactive

compounds and offer opportunities for further commercialization [3]. Previous studies have reported many species, genera, and families that produce edible flowers, but no official list has been published by any international organization, and to date only a small number of them have been published. is being researched. Several scientific reports have focused on bioactive substances used as food sources. Some of them are shown in Table 1.

2.2 Health Benefits of Edible Flowers

Edible flowers provide a palatable blend of aesthetics and nourishment, making them a one-of-a-kind harmonize to culinary creations. Aside from their visual beauty, these blossoms can benefit our overall health and well-being in different ways. First, most edible flowers are rich in antioxidants, such as flavonoids and polyphenols, which fight oxidative stress and reduce the risk of chronic disease [23]. In addition, these flowers often contain vitamins and minerals like as vitamin C, which benefits in immune function, and potassium, which aids in the maintenance of appropriate blood pressure levels [3].

Furthermore, a number of edible flowers, such as calendula and chamomile, have anti-inflammatory effects and can be ingested or applied topically to relieve digestive discomfort or skin irritations [1]. Edible flowers add a rush of colourful flavours and smells to dishes, improving the sensory experience and potentially decreasing the need for excessive salt or sugar [24]. Furthermore, integrating edible flowers into your diet can develop a connection to nature and a deeper appreciation for the beauty and diversity of the natural world, both of which can benefit your mental health. However, while hunting or purchasing edible flowers, it is critical to exercise caution to ensure that they are pesticide-free and safe for consumption. Edible flowers, when utilized wittily and correctly, may be a delectable and health-enhancing addition to the culinary selection.

Table 1. List of common edible flowers, their bioactive compounds

Scientific Name	Common Name	Bioactive Compound	References
<i>Sambucus nigra</i> L.	Black elder	(poly) phenolic compounds, terpenoid.	[4]
<i>Viola tricolor</i> L.	Pansy	Flavones (apigenin glycoside), anthocyanin (cyanidin vâdelphi -nidin)	[5,6]
<i>Centaurea cyanus</i> L.	Cornflower	Cinnamic acid, Flavanols, Benzoic acid, Vitamin C, Tocopherols	[7]
<i>Dendranthema</i> spp.	Chrysanthemum	Caffeoylquinic acids,	[8]
<i>Cucurbita pepo</i> var. giromontiina	Zucchini	Carotenoids, Polyphenols	[9,10]
<i>Cichorium intybus</i> L.	Chicory	Polyphenols, inulin, oligofruc- tose, sesquiterpene lactones	[11]
<i>Cucurbita pepo</i> L.	Pumpkin	carotenoids, polyphenols	[9,10]
<i>Viola odorata</i> L.	Sweet violet	Eugenol, gamma-sitosterol, tetradecanoic acid, Phytol, Octacosanol.	[6]
<i>Forsythia Vah</i>	Forsythia	Forsythiaside, Rutin and Forsythin	[11,12]
<i>Fuchsia</i> L.	Fuchsia	Flavonoids, Phenolic acids, Anthocyanins, Tannins	[13]
<i>Hibiscus</i> L.	Hibiscus	Hydroxycitric acid, Anthocyanins, Flavonoids, Phenolic acid	[14]
<i>Lamium</i> L.	Dead-nettle	Phenolic acids (chlorogenic, caffeic acids), Flavonoids (rutin).	[15]
<i>Jasminum</i> L.	Jasmine	Saponin, Flavonoid, Quinine, Protein, Steroids	[16]
<i>Echinacea Moench</i>	Coneflower	Alkylamides, Polysaccharides, Glycoproteins, Flavonoids and Phenolic compounds	[17]
<i>Trifolium</i> L.	Clover	Flavones, Flavonols, Isoflavones, Pterocarpan, Cyanogenic gluco -sides, Saponins, Condensated tannins	[18]
<i>Lavandula</i> L.	Lavender	Linalool, Limonene, Perillyl alcohol, Linalyl acetate, Cis-smine, Terpene, Coumarin, Tannin, Caffeic acid	[19]
<i>Papaver rhoeas</i> L.	Common poppy	Phenolic compounds, alkaloids, flavonoids, organic acids and coumarins	[20]
<i>Calendula off.</i> L.	Common marigold	Terpenoids, Terpenes, Carotenoids, Flavonoids	[21]
<i>Tropaeolum</i> L.	Nasturtium	Anthocyanins, polyphenols and vitamin C	[22]
<i>Impatiens</i> L.	Impatiens	Phenolic acids and flavonoids	[7]

Table 2. Potential health benefits of common edible flowers

Common name	Scientific name	Health benefits	References
Black elder	<i>Sambucus nigra</i> L. (Black elder)	Antioxidant, antibacterial, bone protection, healing, anti-inflammatory and pain relief	[4]
Pansy	<i>Viola tricolor</i> L.	It is used to treat scabs, itching, ulcers, eczema, and psoriasis, as well as lung and chest infections such as bronchitis and asthma.	[6]
Cornflower	<i>Centaurea cyanus</i> L.	It has antibacterial, anti-inflammatory, neurological, anti-oxidant, diuretic, gastroprotective, and many other effects.	[25]
Chrysanthemum	<i>Dendranthema spp.</i>	Easing stress and anxiety. Improving cardiovascular, health and protect against oxidative damage. Inhibit inflammation and support healthy immune function.	[8]
Zucchini	<i>Cucurbita pepo var. giromontiina</i>	The antioxidant activity of Cucurbitaceae appears to be of great interest in many application areas, including CVD (cardiovascular diseases) associated with oxidative stress.	[9]
Chicory	<i>Cichorium intybus</i> L.	Relieves symptoms associated with mild digestive disorders (such as bloating, flatulence, and indigestion) and temporary loss of appetite.	[11]
Pumpkin	<i>Cucurbita pepo</i> L.	Heal and purify fatigue and thirst blood. Cure colds and relieve pain. Seeds treat irritable bladder, prostate complain, anthelmintic, acaricide. Treatment of gastritis, burns, enteritis, and fever disease. Increase hemoglobin content in the blood.	[10]
Sweet violet	<i>Viola odorata</i> L.	Anti-inflammatory, analgesic, antioxidant, and diuretic effects, as well as other positive effects such as antihypertensive, antihyperlipidemic, diaphoretic, antipyretic, and antifungal effects, as well as weight loss effects, etc.	[5]
Phlox	<i>Phlox</i> L.	Antioxidant and antimicrobial activity	[26]
Forsythia	<i>Forsythia Vah</i>	Antipyretic and anti-inflammatory activity in the treatment of bacterial infections and upper respiratory ailments	[12]
Freesia	<i>Freesia sp.</i>	inhibiting sympathetic nervous excitement and reducing blood pressure	[27]
Hibiscus	<i>Hibiscus</i> L.	Antihypertensive effect, antilipidemia effect, hypoglycemic effect, body fat reduction effect, renal protection effect, antianemia effect, antioxidant effect, anti-inflammatory effect, antistromal effect	[28]
Dead-nettle	<i>Lamium</i> L.	Antimicrobial, antiviral, anti-inflammatory, anti-nociceptive activity, and pain therapy and cytotoxicity and cytoprotective activity	[29]
Jasmine	<i>Jasminum</i> L.	Anti-oxidant, anti-inflammatory, anti-cancer, anti-obesity, and neuroprotective effect	[4]
Coneflower	<i>Echinacea Moench</i>	Antioxidant potential, immune-modulatory effect, therapeutical uses	[17]
Clover	<i>Trifolium</i> L.	estrogenic and progestogenic activities, antioxidant, anti-cancer and others	[30]
Lavender	<i>Lavandula</i> L.	Anti-inflammatory, antiseptic, anti-bacterial, antifungal antimicrobial, antidepressant properties	[31]
Common poppy	<i>Papaver rhoeas</i> L.	Polyphenols content, antioxidant power and antibacterial effects.	[32]

Table 3. List of edible flowers having anti-nutritional compounds

Name of Flower	Toxic Compound
<i>Allium cepa</i>	Phytates, Oxalates, Alkaloids and Saponin
<i>Carica papaya</i>	Tannins, Phytates, Oxalates and Alkaloids
<i>Cucurbita maxima</i>	Tannins, Oxalates, Alkaloids and Saponin
<i>Moringa oleifera</i>	Alkaloids and Saponin
<i>Musa paradisiaca</i>	Tannins, Phytates, Oxalates,
<i>Musa acuminata</i>	Tannins, Alkaloids and Saponin
<i>Wood fordia fruticosa</i>	Phytates, Alkaloids and Saponin

[18,32,33,34]




Fig. 1. Common edible flowers of fruit crops



Fig. 2. Common edible ornamental crops

2.3 Anti-nutritional Compounds in Edible Flowers

Antinutritional factors are substances or compounds found in fruits, flowers, and vegetables that are generally toxic to humans. They reduce the availability of nutrients such as minerals, vitamins, and even proteins found in plants. This affects the nutritional value of these plants. Antinutrients include amino acids, proteins, simple amines, alkaloids, glycosides, and phenolic compounds. When plant foods are consumed as a food source, antinutrients are

absorbed, posing health risks to consumers [35]. Antinutritional factors are natural compounds that are present in varying amounts in different foods, depending on the type of food, how it is grown, and the chemicals used to store it. These antinutritional factors are also called "secondary metabolites" of plants and have been shown to be highly biologically active. These secondary metabolites are secondary compounds that occur as byproducts of processes that lead to the synthesis of primary metabolites. Antinutrients are chemicals that plants produce for self-defense. These anti-nutritional factors are

produced by plants to protect themselves from fungi, insects and predators and represent a plant defense mechanism. These anti-nutritional factors are known to impair metabolism, thereby negatively impacting growth and bioavailability of nutrients [36]. Many edible flowers are also known to contain these anti-nutritional components, and most of these edible flowers are directly consumed by rural people. Although these are not necessarily toxic plant substances, they reduce the nutritional value of plant foods, often making essential nutrients unavailable or indigestible to humans and animals. Animal Consumption [37]. Some vitamins found in foods can be destroyed by antinutrients. Below are some of them and the compounds they contain.

3. CONCLUSION

This review has focused light on the numerous health benefits linked with eating edible flowers. Edible flowers provide a unique and attractive addition to the diet, delivering both flavour and health benefits, thanks to their rich nutritional profiles and potential antioxidant and anti-inflammatory capabilities. However, when selecting edible flowers for food, it is critical to use caution and discretion. The disadvantage is the presence of poisonous blooms, which, if consumed inadvertently, can cause severe health difficulties or even death. To discern between safe and dangerous floral kinds, it is critical to be well-informed and rely on expert help.

ACKNOWLEDGEMENTS

All authors are very thankful to the University Institute of Agricultural Sciences, Chandigarh University for timely support and coordination.

COMPETING INTERESTS

Authors do not have any competing interests.

REFERENCES

1. Swarna T. Nutritional and medicinal value of edible flowers. *Foot*. 2023;43:47.
2. Friedman H, Rot I, Agami O, Vinokur Y, Rodov V, Reznick N, Umiel N, Dori I, Ganot L, Shmuel D, Matan, E. Edible flowers: New crops with potential health benefits. In *International Conference on Quality Management in Supply Chains of Ornamentals*. 2007;755:283-290.
3. De LC. Edible seeds and nuts in human diet for immunity development. *Int. J. Recent Sci. Res.* 2020;6(11):38877-38881.
4. Lu B, Li M, Yin R. Phytochemical content, health benefits, and toxicology of common edible flowers: A review (2000–2015). *Critical Reviews in Food Science and Nutrition*. 2016;56(sup1):130-148.
5. Feyzabadi Z, Jafari F, Kamali SH, Ashayeri H, Aval SB, Esfahani MM, Sadeghpour O. Efficacy of *Viola odorata* in treatment of chronic insomnia. *Iranian Red Crescent Medical Journal*. 2014;16(12).
6. Hellinger R, Koehbach J, Fedchuk H, Sauer B, Huber R, Gruber CW, Gründemann C. Immunosuppressive activity of an aqueous *Viola tricolor* herbal extract. *Journal of Ethnopharmacology*. 2014;151(1):299-306.
7. Pires Jr. EDO, Pereira E, Pereira C, Dias MI, Calhelha RC, Ćirić A, Soković M, Hassemer G, Garcia CC, Caleja C, Barros L. Chemical composition and bioactive characterisation of *Impatiens walleriana*. *Molecules*. 2021;26(5):1347.
8. Shahrajabian MH, Sun W, Zandi P, Cheng Q. A review of chrysanthemum, the eastern queen in traditional Chinese medicine with healing power in modern pharmaceutical sciences. *Applied Ecology & Environmental Research*. 2019;17(6).
9. Rolnik A, Kowalska I, Soluch A, Stochmal A, Olas B. Comparative phytochemical, antioxidant and haemostatic studies of preparations from selected vegetables from Cucurbitaceae family. *Molecules*. 2020;25(18):4326.
10. Ratnam N, Najjibullah M, Ibrahim MD. A review on *Cucurbita pepo*. *Int J Pharm Phytochem Res*. 2017;9:1190-1194.
11. Street RA, Sidana J, Prinsloo G. *Cichorium intybus*: Traditional uses, phytochemistry, pharmacology, and toxicology. *Evidence-Based Complementary and Alternative Medicine*; 2013.
12. Zeng MJ, Zhong YJ, Diao Y. Promoting mechanism of plant growth-promoting rhizobacteria in medicinal plants. *Biotechnology Bulletin*. 2017;33(11):13.
13. Csepregi R, Temesfői V, Das S, Alberti Á, Tóth CA, Herczeg R, Papp N, Kőszegi T. Cytotoxic, antimicrobial, antioxidant properties and effects on cell migration of phenolic compounds of selected transylvanian medicinal plants. *Antioxidants*. 2020;9(2):166.

14. Ojulari OV, Lee SG, Nam JOS. Beneficial effects of natural bioactive compounds from *Hibiscus sabdariffa* L. on obesity. *Molecules*. 2019;24(1):210.
15. Danila D, Adriana T, Camelia SP, Valentin G, Anca M. Antioxidant activity of methanolic extracts of *Lamium album* and *Lamium maculatum* species from wild populations in the Romanian eastern Carpathians. *Planta Medica*. 2015;81(16):192.
16. Prakkash MJ, Ragunathan R, JESTEENA J. Evaluation of bioactive compounds from *Jasminum polyanthum* and its medicinal properties. *Journal of drug delivery and therapeutics*. 2019;9(2):303-310.
17. Burlou-Nagy C, Bănică F, Jurca T, Vicaș LG, Marian E, Muresan ME, Bácskay I, Kiss R, Fehér, P, Pallag A. *Echinacea purpurea* (L.) Moench: Biological and pharmacological properties. A review. *Plants*. 2022;11(9):1244.
18. Ahmad S, Zeb A. Phytochemical profile and pharmacological properties of *Trifolium repens*. *Journal of Basic and Clinical Physiology and Pharmacology*. 2020;32(1):20200015.
19. Habán M, Korczyk-Szabó J, Čerteková S, Ražná K. *Lavandula* species, their bioactive phytochemicals, and their biosynthetic regulation. *International Journal of Molecular Sciences*. 2023;24(10):8831.
20. Hmamou A, Kara M, Khomsi ME, Saleh A, Al Kamaly O, Bendaoud A, El Ouadrhiri F, Adachi A, Tlemcani S, Eloutassi N, Lahkimi A. Comparative Study on the Total Phenolics, Total Flavonoids, and Biological Activities of *Papaver rhoeas* L. extracts from different geographical regions of Morocco. *Applied Sciences*. 2023;13(4):2695.
21. Silva D, Ferreira MS, Sousa-Lobo JM, Cruz MT, Almeida IF. Anti-inflammatory activity of *Calendula officinalis* L. Flower extract. *Cosmetics*. 2021;8(2):31.
22. Jakubczyk K, Janda K, Watychowicz K, Lukasiak J, Wolska J. Garden nasturtium (*Tropaeolum majus* L.)-a source of mineral elements and bioactive compounds. *Roczniki Państwowego Zakładu Higieny*. 2018;69(2).
23. Zhang YJ, Gan RY, Li S, Zhou Y, Li AN, Xu DP, Li HB. Antioxidant phytochemicals for the prevention and treatment of chronic diseases. *Molecules*. 2015;20(12):21138-21156.
24. Mouritsen OG. Tsukemono—crunchy pickled foods from Japan: A case study of food design by gastrophysics and nature. *International Journal of Food Design*. 2018;3(2):103-124.
25. Al-Snafi AE. The pharmacological importance of *Centaurea cyanus*-A review. *Int J of Pharm Rev & Res*. 2020;5(4):379-384.
26. Farinas C, Jourdan P, Paul PA, Peduto Hand F. Development and evaluation of laboratory bioassays to study powdery mildew pathogens of Phlox *In vitro*. *Plant Disease*. 2019;103(7):1536-1543.
27. Xu W, Wang D, Matina P, Lin B, Liu J, Huang J, Li, X. Biological effect of gamma and electron beam to target turning X-ray irradiation on two varieties of freesia in M3 generation. *Journal of Radiation Research and Applied Sciences*. 2023;16(3):100622.
28. Montalvo-González E, Villagrán Z, González-Torres S, Iñiguez-Muñoz LE, Isiordia-Espinoza MA, Ruvalcaba-Gómez JM, Arteaga-Garibay RI, Acosta JL, González-Silva N, Anaya-Esparza LM. Physiological effects and human health benefits of *Hibiscus sabdariffa*: a review of clinical trials. *Pharmaceuticals*. 2022;15(4):464.
29. Salehi B, Armstrong L, Rescigno A, Yeskalyeva B, Seitimova G, Beyatli A, Sharmeen J, Mahomoodally MF, Sharopov F, Durazzo A, Lucarini M. *Lamium* plants—A comprehensive review on health benefits and biological activities. *Molecules*. 2019;24(10):1913.
30. Ramos GP, Apel MA, Morais CBD, Ceolato PC, Schapoval EE, Dall'Agnol M, Zuanazzi JA. In vivo and in vitro anti-inflammatory activity of red clover *Trifolium pratense* dry extract. *Revista Brasileira de Farmacognosia*. 2012;22:176-180.
31. Sharma L, Chandra M, Puneeta A. Health benefits of lavender (*Lavandula angustifolia*). *International Journal of Physiology, Nutrition and Physical Education*. 2020. 4(1):1274-1277.
32. Marsoul A, Ijjaali M, Oumous I, Bennani B, Boukir A. Determination of polyphenol contents in *Papaver rhoeas* L. flowers extracts (soxhlet, maceration), antioxidant and antibacterial evaluation. *Materials Today: Proceedings*. 2020;31:183-189.
33. Costa Marques TH, Santos De Melo CH, Fonseca De Carvalho RB, Costa LM, De Souza AA, David JM, De Lima David JP, De Freitas RM. Phytochemical profile and

- qualification of biological activity of an isolated fraction of *Bellis perennis*. Biological Research. 2013;46(3): 231-238.
34. Borowy A, Chwil, M., Kaplan, M. Biologically active compounds and antioxidant activity of borage (*Borago officinalis* L.) flowers and leaves. Acta Scientiarum Polonorum. Hortorum Cultus. 2017;16(5).
35. Nie R, Zhang Y, Jin Q, Zhang S, Wu G, Chen L, Zhang H, Wang X. Identification and characterisation of bioactive compounds from the seed kernels and hulls of *Paeonia lactiflora* Pall by UPLC-QTOF-MS. Food Research International. 2021;139:109916.
36. Karimi E, Oskoueian E, Karimi A, Noura R, Ebrahimi M. *Borago officinalis* L. flower: A comprehensive study on bioactive compounds and its health-promoting properties. Journal of Food Measurement and Characterization. 2018;12:826-838.
37. Singh O, Khanam Z, Misra N, Srivastava MK. Chamomile (*Matricaria chamomilla* L.): An overview. Pharmacognosy Reviews. 2011.5(9):82.

© 2024 Sood et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/112264>