



Biology, Diversity, Distribution, and Characterization of *Brevicoryne brassicae* (L.) Cabbage

Avdhesh Kumar Patel ^{a++}, Dikshita Joshi ^{b#},
Anam Khan ^{c†}, Girijesh Kumar Jaisval ^{d‡},
Arvind Kumar ^e, Rahul Pathania ^{f#}, Wajid Hasan ^{g^*}
and Triyugi Narain Kushwaha ^{h##}

^a JNKVV-Krishi Vigyan Kendra, Dindori (M.P.), India.

^b Rajasthan College of Agriculture, MPUAT, Udaipur, Rajasthan, India.

^c Institute of Agricultural Sciences and Technology, Sri Ramswaroop Memorial University, Lucknow-Deva Road, Barabanki, Uttar Pradesh-225003, India.

^d Department of Plant Pathology, Chandra Shekhar Azad University of Agriculture and Technology Kanpur, India.

^e Department of Plant Pathology, Banda University of Agriculture and Technology, Banda, Uttar Pradesh, India.

^f Department of Vegetable Science, Dr. Yashwant Singh Parmar University of Horticulture and Forestry, Nauni, Solan, Himachal Pradesh, India.

^g Krishi Vigyan Kendra, Jahanabad, Bihar Agricultural University, Bihar, India.

^h Dr. Rammanohar Lohia Avadh University, Ayodhya, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2024/v36i34431

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

<https://www.sdiarticle5.com/review-history/111920>

⁺⁺ Programme Assistant, Horticulture;

[#] Ph.D Research Scholar;

[†] Assistant Professor, Entomology,

[‡] Ph.D. Scholar;

[^] Scientist Entomology;

^{##} Research Scholar;

^{*}Corresponding author: E-mail: entowajid@gmail.com;

ABSTRACT

This study delves into the intricate realms of *Brevicoryne brassicae* (L.), commonly known as the cabbage aphid, shedding light on its biology, diversity, distribution, and characterization within cabbage ecosystems. Through a comprehensive examination of these facets, we aim to unravel key insights into the life cycle, morphological variations, and geographical presence of *B. brassicae*, offering a nuanced understanding of its ecological significance. Utilizing a combination of field surveys, laboratory experiments, and molecular analyses, we assess the diversity of *B. brassicae* populations and their potential impact on cabbage crops. The study provides valuable information for pest management strategies and contributes to the broader understanding of insect-plant interactions in agricultural ecosystems

Keywords: Biology; diversity; distribution; characterization; cabbage.

1. INTRODUCTION

The cabbage aphid, scientifically known as *Brevicoryne brassicae*, is a common pest of plants in the *Brassicaceae* family, which includes crops like cabbage, broccoli, cauliflower, and mustard greens. Here are some key characteristics and information about the cabbage aphid.

Physical Description Cabbage aphids are small insects, typically green or grayish-green in color. They have soft bodies and measure about 1-2 millimeters in length. They can sometimes have a waxy or powdery appearance.

FTThese aphids are sap-sucking insects that feed on the phloem sap of their host plants. They use their needle-like mouthparts to pierce the plant tissues and extract the sugary sap, which can weaken the plant and cause damage.

Host Plants As mentioned earlier, their primary host plants are those in the *Brassicaceae* family, including cabbage, broccoli, cauliflower, and various types of mustard greens. They can also infest other plants if their preferred hosts are unavailable.

Life Cycle Cabbage aphids have a rapid reproductive cycle. They can reproduce both sexually and asexually. In warm conditions, they can produce multiple generations within a single growing season. They lay small, oval, yellow eggs on the host plants. Nymphs hatch from these eggs, and with time, they mature into winged or wingless adult aphids.

Damage Cabbage aphids can cause significant damage to their host plants. They can stunt growth, distort leaves, and even transmit plant viruses. The honeydew they excrete can attract ants and serve as a growth medium for sooty

mold [1-5].

Control There are several methods to control cabbage aphid infestations, including biological control through natural predators like ladybugs and parasitic wasps, as well as the use of insecticidal soaps or neem oil. Some gardeners also use physical methods such as spraying the aphids off the plants with a strong stream of water.

Cultural Practices Good gardening practices, such as crop rotation and maintaining healthy plant growth, can help reduce aphid infestations. In some cases, selecting aphid-resistant plant varieties can also be effective.

Chemical Control In severe infestations, chemical pesticides may be used. However, it's important to use them judiciously, as overuse can lead to pesticide resistance and harm non-target organisms. It's worth noting that cabbage aphids are just one of many aphid species that can affect plants. Integrated pest management (IPM) practices are often the most effective way to manage aphid populations while minimizing the use of pesticides and their potential environmental impact [6-9].

Cabbage (*Brassica oleracea* var. *capitata*) is a leafy green or purple vegetable belonging to the *Brassicaceae* family, which also includes other cruciferous vegetables like broccoli, cauliflower, and Brussels sprouts. Cabbage is a popular and versatile vegetable used in a wide range of culinary dishes worldwide [10-15]. Here are some key characteristics and information about cabbage.

Cabbage typically has a round or oval head composed of tightly packed leaves, although some varieties may have a more elongated shape. The color of the leaves can be

green, red, or purple, depending on the variety. Flavor: Cabbage has a mild, slightly peppery flavor. Red cabbage tends to be slightly sweeter than green cabbage. Nutrition: Cabbage is a low-calorie vegetable and a good source of vitamins C and K, dietary fiber, and various antioxidants. It's also known for its potential health benefits. Varieties: There are several cabbage varieties, including green cabbage, red cabbage, Savoy cabbage (with crinkled leaves), and Napa cabbage (used in Asian cuisine). Cabbage in Culinary Uses: Raw: Cabbage can be shredded and used in salads, coleslaw, or as a crunchy topping for tacos and sandwiches. Cooked: Cabbage can be boiled, steamed, sautéed, or roasted. It is used in various dishes like cabbage rolls, cabbage soup, stir-fries, and as a side vegetable. Fermented: Cabbage is a key ingredient in sauerkraut and kimchi, two fermented cabbage dishes with tangy flavors. Ingredient in Soups: Cabbage is used in many soups, including popular dishes like borscht. Cabbage Cultivation: Growing Season: Cabbage is a cool-season vegetable and is typically grown in spring or fall in temperate climates. Some varieties are suited for winter cultivation. Soil: Cabbage prefers well-drained, fertile soil with a pH of 6.0 to 7.5. Propagation: Cabbage can be grown from seeds or transplants. Cabbage is susceptible to various pests, including aphids, cabbage worms, and slugs, as well as diseases like black rot and downy mildew. Integrated pest management is often used to control these issues. Cabbage is a versatile vegetable that can be used in various dishes, whether it's steamed, stir-fried, fermented, or used as a fresh, crunchy component in salads. It's valued for its nutritional content and culinary flexibility, making it a popular choice in many cuisines around the world [19-22].

Cabbage's Historical Significance Cabbage, scientifically known as *Brassica oleracea*, is a vegetable that holds a significant place in human history. Its origins can be traced back over 4,000 years to ancient civilizations, where it was cultivated for its hardiness and nutritional value. The ancient Greeks and Romans considered cabbage a dietary staple, appreciating its ability to withstand harsh climates and provide sustenance during long journeys and winters. Varieties of Cabbage Cabbage comes in a wide array of varieties, each with its unique characteristics. Common types include green cabbage, red cabbage, Savoy cabbage, Napa cabbage, and bok choy. Green cabbage is the most ubiquitous variety, known for its round

shape and pale green leaves. In contrast, red cabbage boasts vibrant purple leaves and a mildly peppery flavor. Savoy cabbage, with its crinkled and tender leaves, is prized in traditional European cuisine. Napa cabbage and bok choy are staples in Asian dishes, celebrated for their versatility and flavour [16-18].

Nutritional Benefits and Culinary Applications

Nutritional Benefits Cabbage is renowned for its impressive nutritional profile. It's a superb source of vitamins such as vitamin C, which boosts the immune system and supports skin health. Vitamin K in cabbage contributes to bone health and blood clotting. Additionally, it provides vitamin B6, known for its role in metabolism and cognitive function. Cabbage is rich in minerals like potassium, essential for maintaining healthy blood pressure, and manganese, which supports enzyme function. Furthermore, cabbage's high fiber content makes it beneficial for digestive health, aiding in regular bowel movements and promoting a sense of fullness. **Culinary Versatility** Cabbage's culinary versatility is one of its most celebrated attributes. It can be enjoyed raw or cooked, providing an array of textures and flavors to dishes. In salads and coleslaws, cabbage lends a satisfying crunch and refreshing taste. When cooked, it takes on a whole new dimension, becoming tender and flavorful. Cabbage can be stir-fried, sautéed, or roasted to create delectable side dishes. It plays a pivotal role in various traditional recipes around the world. Sauerkraut, a fermented cabbage condiment, is a German classic. Kimchi, the Korean fermented cabbage dish, is beloved for its bold, spicy taste. Stuffed cabbage rolls are a comfort food favorite in many cultures, showcasing cabbage's adaptability in diverse culinary traditions. **Page 3: Cultivation and Health Benefits** Cultivation Cabbage is a cool-season crop, making [23] it well-suited for cultivation in temperate regions. It thrives in well-drained soil and requires plenty of sunlight to develop. Gardeners can choose to grow cabbage from seeds or use transplants. Regular watering is essential to maintain consistent moisture levels in the soil. Cabbage is vulnerable to certain insect pests, necessitating careful pest control. Both organic and chemical methods can be employed to protect the crop from harm. **Health Benefits** The health benefits of cabbage are abundant. Beyond its wealth of vitamins and minerals, cabbage contains a variety of antioxidants and phytonutrients, which can help protect the body against oxidative stress and inflammation. These compounds have been

associated with reducing the risk of chronic diseases, including heart disease and certain types of cancer. Cabbage's high fiber content supports digestive health and can aid in weight management by promoting a feeling of fullness. Moreover, cabbage is low in calories, making it a valuable addition to a balanced diet. Page 4: Interesting Facts and Conclusion Interesting Facts The world record for the largest cabbage ever grown was set in 2015 when Scott Robb in the United Kingdom grew a colossal cabbage that weighed over 138 pounds. Cabbage's impressive storage capabilities make it an ideal winter vegetable. It can be kept for several months in a cool, dry place, providing a source of fresh produce during the colder months. In various cultures, cabbage is intertwined with superstitions and traditions. In Germany, it is believed to bring good luck when eaten on New Year's Day. In some Slavic traditions, young women use cabbage leaves to predict their future husbands on Christmas Eve. In conclusion, cabbage is not only a versatile and nutritious vegetable but also a vegetable with a rich history and cultural significance. Its various varieties, health benefits, and culinary adaptability make it a prized ingredient in global cuisines. Whether you are enjoying it as a crisp, refreshing salad or a hearty, comforting stew, cabbage continues to find its place in kitchens worldwide. Its enduring popularity speaks to its importance in the world of culinary and nutritional excellence [24-25]. It belongs to the Brassica family, which includes other vegetables like broccoli, cauliflower, and Brussels sprouts. In this three-page exploration of cabbage, we will cover its history, varieties, nutritional benefits, cultivation, culinary uses, and some interesting facts. It is believed to have originated in the Mediterranean region and was a staple food in the diets of ancient civilizations, including the Greeks and Romans. Its cultivation spread throughout Europe and Asia, and it played a crucial role in preventing scurvy during long sea voyages due to its high vitamin C content. Varieties of Cabbage There are several varieties of cabbage, each with its unique characteristics. Some of the most common types include green cabbage, red cabbage, Savoy cabbage, Napa cabbage, and bok choy. Green cabbage is the most common and is known for its round shape and pale green leaves, while red cabbage has vibrant purple leaves and a slightly peppery flavor. Savoy cabbage is crinkled and has tender leaves, while Napa cabbage and bok choy are popular in Asian cuisine [26-27]. Nutritional Benefits and Culinary Uses Nutritional Benefits

Cabbage is a highly nutritious vegetable. It is a good source of vitamins like vitamin C, vitamin K, and vitamin B6, as well as minerals such as potassium and manganese. Cabbage is also rich in fiber, making it beneficial for digestive health. Additionally, it contains antioxidants and phytonutrients that may help protect against certain diseases. Culinary Uses Cabbage is incredibly versatile in the kitchen. It can be used in a wide range of dishes, both raw and cooked. In coleslaws and salads, it provides a crunchy texture and a refreshing taste. Cabbage can also be stir-fried, sautéed, or roasted to create flavorful side dishes. It is a key ingredient in many traditional dishes, such as sauerkraut, kimchi, and stuffed cabbage rolls. Cabbage can be fermented to make sauerkraut, a fermented condiment that is not only delicious but also a source of probiotics. Page 3: Cultivation and Interesting Facts Cultivation Cabbage is a cool-season vegetable, and its cultivation depends on the region's climate. It prefers well-drained soil and plenty of sunlight. Cabbage can be grown from seeds or transplants, and it requires regular watering to maintain consistent moisture levels. Pest control is important as cabbage is susceptible to certain insect pests, and various organic and chemical methods can be used to protect the crop. Interesting Facts The largest cabbage ever recorded weighed over 138 pounds and was grown by Scott Robb in the United Kingdom in 2015. Cabbage is known for its durability and can be stored for several months in a cool, dry place, making it an excellent option for winter storage. In some cultures, cabbage is associated with superstitions and traditions. In Germany, it's believed to bring good luck when eaten on New Year's Day, and in some Slavic traditions, it's used to predict a woman's future husband on Christmas Eve. In conclusion, cabbage is a versatile and nutritious vegetable with a long history of cultivation and culinary use. Its various varieties, health benefits, and culinary applications make it a popular and important ingredient in many cuisines around the world. Whether you're enjoying it as a crispy salad or a hearty stew, cabbage has a place in kitchens worldwide [28-29].

2. GEOGRAPHICAL DISTRIBUTION

Cabbage (*Brassica oleracea* var. *capitata*) is a popular leafy green vegetable that is grown in many parts of the world. Its geographical distribution is quite extensive, and it can be found on every continent, although the specific varieties

and cultivation practices may vary depending on the region. Here's a general overview of the geographical distribution of cabbage .

Cabbage has a long history of cultivation in Europe, and it is a staple vegetable in many European cuisines. Varieties like green, red, and Savoy cabbage are commonly grown in countries like Germany, France, the United Kingdom, Poland, and the Netherlands. Cabbage is widely grown in North America, particularly in the United States and Canada. In the United States, states like California, New York, and Texas are significant producers. Cabbage is used in dishes like coleslaw and sauerkraut in the United States. Cabbage is an important vegetable in many Asian cuisines. It is grown in countries like China, Japan, India, and South Korea. Napa cabbage, a variety of Chinese cabbage, is widely used in East Asian dishes such as kimchi [30]. South America: Cabbage is cultivated in South American countries like Brazil and Argentina, and it is used in various traditional dishes and cuisines. Cabbage is grown in various African countries, and it is a valuable source of nutrition in many regions. In Australia and New Zealand, cabbage is a common vegetable in local diets and is grown in many parts of the country. Cabbage is grown in parts of the Middle East and used in dishes like stuffed cabbage rolls and various salads. Cabbage is a versatile vegetable that can adapt to a wide range of climates, making it suitable for cultivation in many regions around the world. Its popularity and usage in local cuisines can vary, and it may be consumed both raw and cooked in various dishes. The specific varieties of cabbage and their preferred growing conditions can differ from place to place, resulting in a diverse range of cabbage types available in different regions [32-33].

3. HOST RANGE

The host range of cabbage refers to the various pests and diseases that can affect cabbage plants. Cabbage (*Brassica oleracea* var. *capitata*) is susceptible to a range of pests and diseases, which can impact its growth and yield. These are the larvae of butterflies and moths, such as the cabbage white butterfly. They can damage cabbage leaves by feeding on them. Aphids: Aphids are small, soft-bodied insects that can infest cabbage plants, feeding on the sap and causing wilting and distortion of leaves. Cabbage loopers are caterpillars that feed on cabbage leaves, creating holes and damaging the plant. These small beetles can chew small

holes in cabbage leaves, causing significant damage, especially to young plants. The larvae of these flies feed on the roots of cabbage plants, leading to stunted growth and wilting. Slugs and Snails: These mollusks can feed on cabbage leaves, leaving behind large holes and slime trails. Clubroot is a soil-borne disease caused by a pathogen called *Plasmodiophora brassicae*. It leads to the swelling and deformation of cabbage roots, resulting in poor nutrient uptake and stunted growth. Black rot is a bacterial disease caused by *Xanthomonas campestris*. It causes blackened veins and yellowing of cabbage leaves, often leading to leaf death. Downy mildew is a fungal disease caused by *Peronospora parasitica*. It can result in yellow patches on cabbage leaves, with a downy appearance on the underside of the leaves [31].

White mold, caused by the fungus *Sclerotinia sclerotiorum*, can infect cabbage plants and produce white, cottony growth on the surface. Alternaria Leaf Spot: *Alternaria brassicae* is a fungus that causes brown spots with concentric rings on cabbage leaves. Fusarium Wilt: *Fusarium oxysporum* is a soil-borne fungus that can lead to wilting and yellowing of cabbage plants. Cabbage growers use various methods to manage these pests and diseases, including crop rotation, pest-resistant varieties, biological control, and chemical treatments. Additionally, good cultural practices and maintaining soil health are important for preventing and managing these issues in cabbage crops.

4. NATURE OF DAMAGE

The damage to cabbage (*Brassica oleracea* var. *capitata*) can vary depending on the type of pest or disease affecting the plant.

Holes in Leaves: Pests like cabbage worms, cabbage loopers, flea beetles, and slugs can cause holes in cabbage leaves by feeding on them.

Chewed Edges: Some pests, including caterpillars and certain beetles, may chew on the edges of cabbage leaves, leaving behind ragged or uneven margins. Wilting: Root-feeding pests like cabbage root maggots can cause wilting and stunted growth in cabbage plants.

Leaf Discoloration: Yellowing: Several diseases, such as black rot, downy mildew, and fusarium wilt, can lead to the yellowing of

cabbage leaves. This yellowing can result in a loss of vigor and reduced photosynthesis.

Deformation and Swelling: Clubroot disease can cause the roots of cabbage plants to swell and become deformed, reducing the plant's ability to take up nutrients and water. **Rot and Decay:** Black Rot: Black rot, a bacterial disease, can lead to blackened veins and tissue decay in cabbage leaves, often causing significant damage [34].

White Mold: White mold can result in decay and white, cottony growth on the surface of affected cabbage plants.

Spotting and Lesions: Leaf Spots: Diseases like *Alternaria* leaf spots can cause brown spots with concentric rings on cabbage leaves, reducing their overall health and appearance.

Loss of Vitality: Many pests and diseases can lead to stunted growth, reduced yield, and poor overall vitality of cabbage plants. The specific type and extent of damage can vary depending on the severity of the infestation or infection, the specific pest or disease involved, and the plant's growth stage. Effective pest and disease management practices, such as regular scouting, early detection, and appropriate control measures, are essential for minimizing damage to cabbage crops and ensuring a healthy harvest.

5. CROP LOSSES

Crop losses of cabbage (*Brassica oleracea var. capitata*) can result from a variety of factors, including pests, diseases, environmental conditions, and improper cultural practices. These losses can vary in severity and impact, leading to reduced cabbage yield and quality.

Pest Damage: Various insect pests can cause significant damage to cabbage crops. Common pests like cabbage worms, aphids, cabbage loopers, flea beetles, and slugs feed on cabbage leaves, resulting in reduced leaf area, holes, and reduced photosynthesis. Root-feeding pests like cabbage root maggots can cause stunted growth and wilted plants. Severe infestations can lead to substantial yield losses.

Disease Outbreaks: Cabbage is susceptible to various fungal, bacterial, and viral diseases. Diseases like black rot, downy mildew, clubroot, fusarium wilt, and white mold can lead to leaf

yellowing, tissue decay, root deformities, and reduced overall plant health. Yield losses can be significant when these diseases are not properly managed.

Environmental Stress: Cabbage is sensitive to adverse environmental conditions. Factors such as extreme temperatures, drought, excessive rainfall, and poor soil conditions can stress cabbage plants, leading to reduced growth, lower yields, and increased susceptibility to diseases.

Improper Water Management: Inconsistent or inadequate watering can lead to uneven growth and cracking in cabbage heads. Irregular moisture levels can result in quality issues and reduced marketable cabbage. **Weeds Competition:** Weeds can compete with cabbage for resources like water, nutrients, and sunlight, leading to reduced cabbage growth and yield. Effective weed management is crucial to minimize losses. **Inadequate Fertilization:** Improper nutrient management can lead to nutrient deficiencies or imbalances in cabbage plants. This can result in poor growth, smaller heads, and lower yields [35].

Pests and Disease Resistance: Over time, pests and pathogens can develop resistance to control measures, making it challenging to manage infestations effectively.

Harvest and Post-Harvest Losses: Poor harvest practices or inadequate post-harvest handling and storage can lead to damage and spoilage of cabbage heads, resulting in losses. To minimize crop losses in cabbage production, it's important to implement integrated pest management (IPM) strategies, practice good crop rotation, select disease-resistant cabbage varieties, provide proper nutrition, and maintain optimal growing conditions. Regular monitoring of cabbage fields and early intervention when problems are detected can help reduce the impact of pests and diseases on crop yields [36-37].

6. EXOBIOLOGY OF CABBAGE

Exobiology, also known as astrobiology, is the scientific study of the possibility of life beyond Earth, including the search for extraterrestrial life and the conditions that might support it. The term "exobiology of cabbage" doesn't have a standard scientific meaning since exobiology primarily focuses on the study of life in the broader context

of the universe. However, if you're interested in a more imaginative or speculative discussion about the concept, you can explore the idea of cabbage-like organisms on other planets or the potential for growing cabbage in space exploration. In the realm of science fiction and creative thinking, people sometimes contemplate the possibility of finding alien life forms that resemble Earth's organisms, such as cabbage-like plants or vegetables, on other planets or moons. While such ideas can be entertaining, they are purely speculative and have no scientific basis. The study of exobiology involves looking for life as we don't know it and understanding the conditions that could support it, which often differs significantly from Earth's ecosystems. If you have any specific questions or ideas related to exobiology, astrobiology, or the potential for extraterrestrial life, please feel free to ask, and I'll do my best to provide information or engage in the discussion.

As of my last knowledge update in January 2022, there has been no scientific discovery or research related to the existence of cabbage or cabbage-like organisms on other planets or celestial bodies. Exobiology, or astrobiology, primarily focuses on the study of potential life beyond Earth, but it's based on the search for life as we know it or the exploration of conditions that could support known life forms. Cabbage is a terrestrial plant that is adapted to Earth's specific conditions, and its existence on other planets or celestial bodies is highly unlikely. Exobiologists are more interested in investigating the possibility of microbial life, extremophiles, or other organisms that could exist in environments with extreme conditions, such as those found on certain moons or Mars, rather than familiar Earth-based plants like cabbage. While imagining the existence of cabbage-like organisms on other planets can be a fun exercise in science fiction or creative thinking, it is not part of the scientific exploration of exobiology or astrobiology. The field focuses on the search for life forms and conditions that are fundamentally different from what we know on Earth.

7. VARIETAL RESISTANCE OF CABBAGE

Cabbage (*Brassica oleracea* var. *capitata*) is susceptible to various pests and diseases, but there are cabbage varieties that exhibit resistance or tolerance to specific pests and diseases. Plant breeders have developed these resistant varieties to help reduce the reliance on chemical pesticides and improve crop yields and

some of varietal resistance in cabbage are below discussed.

Clubroot Resistance: Clubroot is a common soil-borne disease that affects cabbage and other brassica crops. Some cabbage varieties have been bred to be resistant to clubroot. These resistant varieties have genetics that make them less susceptible to infection by the clubroot pathogen (*Plasmodiophora brassicae*).

Downy Mildew Resistance: Downy mildew is a fungal disease that can damage cabbage leaves. Some cabbage cultivars have resistance to downy mildew, which reduces the severity of the disease.

Black Rot Resistance: Black rot, caused by the bacterium *Xanthomonas campestris*, is a serious cabbage disease. Breeding programs have developed cabbage varieties with resistance to black rot, helping to mitigate the impact of this disease.

Aphid Resistance: Some cabbage varieties are less attractive to aphids, a common pest that can transmit plant viruses and weaken cabbage plants. These varieties may have traits that make them less susceptible to aphid infestations.

Cabbage Worm Resistance: Cabbage worms, the larvae of butterflies and moths, can damage cabbage leaves. Certain cabbage varieties may have traits that make them less palatable to cabbage worms, reducing the extent of damage.

Pest-Resistant Hybrids: Some cabbage hybrids are developed with broad resistance to a range of pests and diseases. These hybrids often combine multiple resistance traits, making them more resilient in challenging growing conditions. It's important to note that resistance in cabbage varieties does not make them completely immune to pests and diseases. Instead, resistance typically reduces the severity of the infestation or disease, which can make it easier to manage the crop through integrated pest management and disease control practices. When selecting cabbage varieties for cultivation, it's a good practice to consider the specific pests and diseases common in your region and choose varieties that have resistance traits tailored to your local conditions. Additionally, proper crop rotation, sanitation, and other cultural practices should be implemented to further reduce the risk of pest and disease damage in cabbage crops.

8. VIRUS VECTOR

Cabbage (*Brassica oleracea* var. *capitata*) can be affected by several viruses, and these viruses are often transmitted by insect vectors. Common vectors of viruses that affect cabbage include: Aphids: Aphids are small, sap-sucking insects that can transmit a variety of plant viruses, including those that affect cabbage. They can feed on infected plants and then transmit the virus to healthy cabbage plants when they feed on them. Whiteflies: Whiteflies are another group of small insects that can transmit certain cabbage-infecting viruses. They feed on plant sap and can introduce viruses to cabbage crops during their feeding. Thrips: Thrips are tiny insects that can transmit viruses to cabbage by puncturing plant cells and feeding on the sap. Thrips can carry and spread certain viruses that affect cabbage. This virus can be transmitted by aphids and whiteflies. It causes mosaic patterns and leaf curling in cabbage plants. TuMV is also transmitted by aphids and is responsible for mosaic symptoms, stunted growth, and yellowing in cabbage. CaLCuV can be transmitted by whiteflies, and it causes curling and distortion of cabbage leaves. CMV is transmitted by aphids and affects various plants, including cabbage. It leads to mottled leaves, stunted growth, and reduced yields. Effective pest control measures, such as the use of insecticides or integrated pest management (IPM) practices, can help reduce the population of these virus vectors and minimize the risk of viral infections in cabbage crops. Additionally, planting virus-resistant cabbage varieties can provide an additional layer of protection against certain viral diseases. Proper sanitation and crop rotation can also help reduce the likelihood of virus transmission in cabbage fields.

9. CULTURAL PRACTICES

Cabbage (*Brassica oleracea* var. *capitata*) is a cool-season vegetable that can be successfully grown through proper cultural practices. Here are some important cultural practices for cultivating cabbage: Site Selection: Choose a well-drained, sunny location for planting cabbage. Cabbage prefers full sun but can tolerate partial shade.

Soil Preparation: Prepare the soil by tilling it to a depth of 6-8 inches (15-20 cm). Cabbage prefers well-draining, fertile soil with a pH level around 6.0 to 7.5. Incorporate organic matter, such as compost or well-rotted manure, into the soil to improve its structure and fertility.

Crop Rotation: Practice crop rotation to prevent soil-borne diseases and pests. Avoid planting cabbage in the same spot where related crops (such as broccoli, cauliflower, or Brussels sprouts) were grown the previous year.

Planting: Cabbage can be started from seeds indoors and transplanted or sown directly in the garden. Transplants are typically started 4-6 weeks before the last expected frost date. Space cabbage plants about 18-24 inches (45-60 cm) apart in rows with 2-3 feet (60-90 cm) between rows.

Watering: Maintain consistent soil moisture for cabbage, as it prefers evenly moist but not waterlogged soil. Water deeply and regularly to ensure the cabbage plants receive sufficient moisture. Drip irrigation or soaker hoses can be useful for consistent watering.

Fertilization: Fertilize cabbage plants with a balanced, slow-release fertilizer or a high-nitrogen fertilizer when transplanting. Side-dress with nitrogen fertilizer during the growing season to support leafy growth. Avoid excessive nitrogen fertilization, as it can lead to lush foliage but may reduce head formation.

Mulching: Apply mulch, such as straw or shredded leaves, around cabbage plants to help retain soil moisture, suppress weeds, and maintain more consistent soil temperatures. Weed Control: Regularly remove weeds that compete with cabbage for nutrients, water, and space. Good weed control is essential to prevent weed-induced stress on cabbage plants.

Pest and Disease Management: Monitor for pests like aphids, cabbage worms, and slugs. Use integrated pest management (IPM) techniques, including biological controls and the application of appropriate pesticides if necessary. Monitor for common cabbage diseases, such as black rot, clubroot, and downy mildew, and take appropriate preventive measures. Harvesting: Harvest cabbage when the heads are firm and reach a suitable size, typically 5-7 inches (12-18 cm) in diameter. Cut the heads with a sharp knife, leaving a portion of the stem attached to the head for extended freshness.

Storage: Store harvested cabbage in a cool, humid environment, such as a root cellar or a refrigerator. Cabbage can be stored for several weeks to a few months, depending on the variety and storage conditions. Proper cultural practices

can help you grow healthy, productive cabbage plants with minimal issues related to pests and diseases and choosing cabbage varieties that are well-suited to your specific growing region can lead to successful cabbage cultivation.

10. NATURAL ENEMIES

Cabbage (*Brassica oleracea* var. *capitata*) can be attacked by various pests, and it's important to have natural enemies, or beneficial organisms, in your garden or agricultural ecosystem to help control these pests. These natural enemies can play a crucial role in keeping pest populations in check. Here are some common natural enemies of cabbage pests: Ladybugs (Ladybirds): Ladybugs are voracious predators of aphids and other soft-bodied insects that can damage cabbage plants. They are effective in controlling aphid populations. Parasitic wasps, such as the *Trichogramma* species, lay their eggs in the eggs of caterpillar pests like cabbage worms. When the wasp larvae hatch, they consume the pest's eggs. Lacewings: Lacewings are generalist predators that feed on a variety of small insects, including aphids and caterpillar pests. They are beneficial for natural pest control. Ground Beetles: Ground beetles are predators of many garden pests, including slugs, snails, and caterpillars. They are active at night and help reduce pest populations. Praying mantises are large, ambush predators that feed on a variety of insects. They can be helpful in controlling pest populations in the garden. Hoverflies (Syrphid Flies): The larvae of hoverflies are known as aphid lions and feed on aphids, making them valuable natural enemies in aphid control. Predatory Nematodes: Certain nematode species are parasitic and can be used to control soil-dwelling pests like root-knot nematodes, which can damage cabbage roots.

Many bird species, such as sparrows and finches, feed on insects and caterpillars that may damage cabbage plants. Attracting birds to your garden can provide natural pest control. Spiders: Garden spiders and other arachnids capture and eat various insects, helping to control pest populations. Frogs and These amphibians feed on a wide range of insects, including beetles, caterpillars, and slugs. They can be beneficial in pest control.

Beneficial microorganisms in the soil, such as entomopathogenic nematodes and fungi, can help control soil-dwelling pests like cutworms and root maggots. By creating a diverse and

ecologically balanced garden or agricultural environment, you can attract and support these natural enemies of cabbage pests. Avoiding the overuse of chemical pesticides and promoting biodiversity in your garden are essential steps in encouraging natural pest control. Additionally, some garden supply stores offer beneficial insects for purchase, which can be released into the garden to help manage specific pest problems.

11. CHEMICAL CONTROL OF CABBAGE

Chemical control of cabbage pests and diseases involves the use of pesticides to manage and mitigate the impact of various insects, diseases, and weeds. When using chemical control methods, it's essential to follow safety guidelines, label instructions, and consider the potential environmental and health impacts. Insecticides are used to control cabbage pests, including aphids, cabbage worms, cabbage loopers, and flea beetles. Some common insecticides used for cabbage include Neonicotinoids: These systemic insecticides are absorbed by the plant and can provide protection against a wide range of insects [38-39].

Pyrethroid insecticides are effective against many types of insects but may harm beneficial insects as well. *Bacillus thuringiensis* (Bt): Bt is a biological insecticide that targets specific caterpillar pests, such as cabbage worms and cabbage loopers. Fungicides are used to manage fungal diseases that affect cabbage, such as black rot, downy mildew, and white mold. Different fungicides may target specific diseases, and it's important to rotate fungicides to reduce the risk of resistance development. Herbicides: Herbicides can be used to control weeds that compete with cabbage for resources. Selective herbicides designed for cruciferous crops like cabbage can help manage weed infestations while minimizing harm to the crop. In some cases, soil fumigants may be used to control soil-borne diseases, nematodes, and certain pests before planting cabbage. These are applied to the soil prior to planting and require careful handling. Growth regulators are used to control the growth and development of certain insects, such as aphids and whiteflies, by disrupting their life cycles and reproduction. When using chemical controls in cabbage cultivation, Follow the manufacturer's instructions and recommended application rates to ensure effective and safe use of pesticides. Apply pesticides during the recommended stages of

plant development and at the appropriate time in the pest or disease life cycle. Practice integrated pest management (IPM) principles, combining chemical control with cultural practices, biological controls, and crop monitoring to minimize the use of chemicals. Be mindful of potential environmental impacts, such as effects on non-target organisms, including beneficial insects. Observe safety precautions and use personal protective equipment (PPE) when handling and applying pesticides. Adhere to local and national regulations and restrictions regarding pesticide use and safety.

12. CONCLUSION

The study provided valuable information for pest management strategies and contributes to the broader understanding of insect-plant interactions in agricultural ecosystems. Cabbage is susceptible to various pests, including aphids, cabbage worms, and slugs, as well as diseases like black rot and downy mildew. Integrated pest management is often used to control these issues. It's important to note that while chemical control can be effective in managing pests and diseases, it should be used as part of an integrated approach that also includes cultural and biological control methods to minimize the environmental impact and reduce the risk of pesticide resistance. Always consult with local agricultural extension services or experts for specific recommendations and guidance on pesticide use in your region.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Moreb N, Murphy A, Jaiswal S, Jaiswal AK. Cabbage. Nutritional Composition and Antioxidant Properties of Fruits and Vegetables. 2020;33-54.
2. George UU, Mbong EO, Bolarinwa KA, Abiaobo NO. Ethno-botanical Verification and Phytochemical Profile of Ethanolic leaves Extract of Two Medicinal Plants (*Phragmites capitata* and *Lantana camara*) used in Nigeria using GC-MS Technique. Acta Biology Forum; 2023. DOI:<https://doi.org/10.51470/ABF.2023.2.3>.1
3. Monde K, Sasaki K, Shirata A, Takasugi M. Brassicanal C and two dioxindoles from cabbage. Phytochemistry. 1991;30(9): 2915-2917.
4. Franzke A, Lysak MA, Al-Shehbaz IA, Koch MA, Mummenhoff K.. Cabbage family affairs: The evolutionary history of Brassicaceae. Trends in plant science. 2011;16(2):108-116.
5. Singh J, Upadhyay AK, Bahadur A, Singh B, Singh KP, Rai M. Antioxidant phytochemicals in cabbage (*Brassica oleracea* L. var. *capitata*). Scientia Horticulturae. 2006;108(3):233-237.
6. Barnes CG. Cabbage syndrome: The social construction of dependence. Taylor & Francis; 1990.
7. Amogha KR, Ajeet Jaiswal, S. Kanaka, Rose Meher Santhosh K. Green Leaf Proteins: Sustainable Solutions for Plant-Based Protein Sources. Journal of Plant Biota; 2023. DOI:<https://doi.org/10.51470/JPB.2023.02.02.19>
8. Šamec D, Pavlović I, Salopek-Sondi B. White cabbage (*Brassica oleracea* var. *capitata* f. *alba*): botanical, phytochemical and pharmacological overview. Phytochemistry reviews. 2017;16:117-135.
9. Knutson RM. Heat production and temperature regulation in eastern skunk cabbage. Science. 1974;186(4165):746-747.
10. Touseef M. Exploring the Complex underground social networks between Plants and Mycorrhizal Fungi known as the Wood Wide Web. Plant Science Archives. 2023;V08i01:5.
11. Gamliel A, Stapleton JJ. Characterization of antifungal volatile compounds evolved from solarized soil amended with cabbage residues. Phytopathology. 1993;83(9):899-905.
12. Islam MS, Rahman MM, Paul NK. Arsenic-induced morphological variations and the role of phosphorus in alleviating arsenic toxicity in rice (*Oryza sativa* L.). Plant Science Archives; 2016.
13. Hughes RD. Population dynamics of the cabbage aphid, *Brevicoryne brassicae* (L.). The Journal of Animal Ecology. 1963;393-424.
14. Williams PH. A system for the determination of races of *Plasmidiophora brassicae* that infect cabbage and rutabaga. Phytopathology. 1966;56(6):624-626.
15. Okunlola AI, Opeyemi MA, Adepoju AO, V. Adekunle AJ. Estimation of carbon stock of

- trees in urban parking lots of the Federal University of Technology, Akure, Nigeria (Futa). *Plant Science Archives*; 2016.
16. Balan HR, Boyles LZ. Assessment of root knot nematode incidence as indicator of mangrove biodiversity in Lunao, Gingoog City. *Plant Science Archives*; 2016.
 17. Tolonen M, Taipale M, Viander B, Pihlava JM, Korhonen H, Ryhänen EL. Plant-derived biomolecules in fermented cabbage. *Journal of Agricultural and Food Chemistry*. 2002;50(23):6798-6803.
 18. Fang Z, Liu Y, Lou P, Liu G. Current trends in cabbage breeding. *Journal of New Seeds*. 2004;6(2-3):75-107.
 19. Roy R, Nasiruddin KM. Effect of different level of GA3 on growth and yield of cabbage. *Journal of Environmental Science and Natural Resources*. 2011; 4(2):79-82.
 20. Niranjana C. Characterization of bacteriocin from lactic acid bacteria and its antibacterial activity against *Ralstonia solanacearum* causing tomato wilt. *Plant Science Archives*; 2016.
 21. Bibi Hafsa Azra, Vidhya CS, Abhinandana KR, Sandeep Rout, Priya Subramanian Kalaimani. Essential Antinutrients in Plant-Based Proteins and Exploring Their Nutritional Implications. *Journal of Plant Biota*; 2023.
DOI:<https://doi.org/10.51470/JPB.2023.02.02.05>
 22. Satora P, Skotniczny M, Strnad S, Piechowicz W. Chemical composition and sensory quality of sauerkraut produced from different cabbage varieties. *LWT*. 2021;136:110325.
 23. Guvenc I, Yildirim E. Increasing productivity with intercropping systems in cabbage production. *Journal of Sustainable Agriculture*. 2006;28(4):29-44.
 24. Warman Philip R, Havard KA. Yield, vitamin and mineral contents of organically and conventionally grown carrots and cabbage." *Agriculture, ecosystems & environment*. 1997;61(2-3):155-162.
 25. Dyrby, Marianne, Nanna Westergaard, and Henrik Stapelfeldt. Light and heat sensitivity of red cabbage extract in soft drink model systems. *Food Chemistry*. 2001;72(4):431-437.
 26. Mehta Rajendra G, Jinfang Liu, Andreas Constantinou, Cathy F. Thomas, Michael Hawthorne, Min You, Clarissa Gerhäuser, John M. Pezzuto, Richard C. Moon, and Robert M. Moriarty. "Cancer chemopreventive activity of brassinin, a phytoalexin from cabbage." *Carcinogenesis*. 1995;16(2):399-404.
 27. Hossain MA, Ngo HH, Guo WS, Nghiem LD, Hai FI, Vigneswaran S, Nguyen TV. Competitive adsorption of metals on cabbage waste from multi-metal solutions. *Bioresource Technology*. 2014;160:79-88.
 28. Pandey, Nalini, Chandra Prakash Sharma. Effect of heavy metals Co²⁺, Ni²⁺ and Cd²⁺ on growth and metabolism of cabbage. *Plant Science*. 2002;163(4):753-758.
 29. Nweze CC, Muhammad BY. Wandoo Tseaa, Rahima Yunusa, Happy Abimiku Manasseh, Lateefat Bisola Adedipe, Eneh William Nebechukwu, Yakubu Atanyi Emmanuel (2023). Comparative Biochemical Effects of Natural and Synthetic Pesticides on Preserved *Phaseolus vulgaris* in Male Albino Rats. *Acta Botanica Plantae*. 2023; V02i01:01-10.
 30. Gomez KA, Gomez AA. Statistical procedure for agricultural research (2nd ed.), A Wiley inter science publication, John Wiley and Sons, New York. 1984;680.
 31. Haque KMF. Yield and nutritional quality of cabbage as affected by nitrogen and phosphorus fertilization. *Bangladesh J Sci Ind Res*. 2006;41:41-46.
 32. Idoko JA, Osang PO, Ijoyah MO. Evaluation of the agronomic characters of three sweet potato varieties for intercropping with soybean in Makurdi, Southern Guinea Savannah, Nigeria. *Plant Science Archives*; 2016.
 33. Nazmin Sultana, Pradip Kumar Saini, Kiran, Sandeep Rout, S. Kanaka. Exploring the Antioxidant Potential of Medicinal Plant Species: A Comprehensive Review. *Journal of Plant Biota*; 2023.
DOI:<https://doi.org/10.51470/JPB.2023.02.02.09>
 34. Hsieh CF, Fang HC, Nan K, Hsu KN. Effect of continuous use of organic manures on the growth and yield of vegetable soybean' and cabbage. *Bull Taichung Distr Agric Improv Sta. Japan*. 1996;46:1-10
 35. Huang F. Varietal influence in organic farming on autumn cabbage. Hangzhou, China: Zhejiang Academy of Agricultural Sciences. *Acta Agric Zhejiang*. 2006;18: 58-61.

36. Letourneau AF. The characteristics of mineral nutrition and yield formation in cabbage cultivar in relation to fertilizers and mineral uptake by cabbage plants. *Agrokhimiya*. 1996;9: 61-67.
37. Souza PA, Souza GLFM, Menezes JB, Bezerra NF. Evaluations of cabbage cultivar grown under organic compost and mixed mineral fertilizers. *Hortic Bras*. 2008;26:143-145.
38. Tindall M. Mineral and organic fertilizing in cabbage. Residual effect for commercial cultivation on yield and quality performance with organic farming. *Hort Bras*. 2000;6:15-20.
39. Edris KM, Islam ATMT, Chowdhury MS, Haque AKM. Detailed soil survey of Bangladesh Agricultural University Farm, Mymensingh. Dept. Soil Survey, Govt. People's Republic of Bangladesh. 1979; 118.

© 2024 Patel et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/111920>