



Optimizing Yield and Quality of Garlic (*Allium sativum* L.) cv. Anand Kesari through Planting Time and Plant Spacing Management

Kena Patel ^{a*}, A. V. Kotecha ^{b++} and M. M. Masu ^{c#}

^a Department of Horticulture, B. A. College of Agriculture, Anand Agricultural University, Anand, Gujarat, India.

^b Department of Horticulture, College of Agriculture, Anand Agricultural University, Jabugam-39115, India.

^c Anand Agricultural University, Anand- 388110, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ijpss/2024/v36i74762>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/117874>

Original Research Article

Received: 02/05/2024
Accepted: 13/06/2024
Published: 20/06/2024

ABSTRACT

The field experiment was conducted during rabi season of the year 2022-23 at B. A. College of Agriculture, Anand Agricultural University. There were nine treatments consisting of three planting time (P₁: 1st week of October, P₂: 3rd week of October and P₃: 1st week of November) and spacing (S₁: 15 x 10 cm, S₂: 15 x 15 cm and S₃: 20 x 10 cm). Results revealed that planting time P₂ (3rd week of October) led to the most favorable outcomes, with bulb diameter (4.46 cm), bulb weight

⁺⁺ Assistant Professor;

[#] Assistant Research Scientist, Directorate of Research;

*Corresponding author: E-mail: kenapatel78@gmail.com;

Cite as: Patel, Kena, A. V. Kotecha, and M. M. Masu. 2024. "Optimizing Yield and Quality of Garlic (*Allium Sativum* L.) Cv. Anand Kesari through Planting Time and Plant Spacing Management". *International Journal of Plant & Soil Science* 36 (7):529-34. <https://doi.org/10.9734/ijpss/2024/v36i74762>.

(13.18), yield (1.93 kg/plot and 60.70 q/ha). Spacing S₂ (15 x 15 cm) also exhibited positive effects on bulb quality like bulb diameter (4.42 cm) and bulb weight (12.01 g), while yield (1.934 kg/plot and 59.20 q/ha) were recorded to be significantly the highest at spacing S₁ (15 x 10 cm). The P₂ S₃ recorded significantly higher bulb diameter (4.84 cm), weight of bulb (13.61 g). While, yield (2.341 kg/plot and 70.93 q/ha) were recorded to be significantly highest at P₂S₂. These findings provide valuable insights for garlic cultivation, helping farmers optimize planting strategies to enhance crop performance and yield.

Keywords: Planting time; spacing; garlic; bulb diameter; bulb weight; yield; sulphur content.

1. INTRODUCTION

Garlic (*Allium sativum* L.) is a significant bulb crop, second only to onions, and is believed to have originated from Central Asia and Southern Europe. It is grown in India on a large scale, with Gujarat, Madhya Pradesh, Maharashtra, Uttar Pradesh, and Rajasthan being major garlic-growing states. Garlic is known for its numerous medicinal properties and culinary uses, containing essential minerals, vitamins, and other beneficial substances [1,2,3]. The growth, yield, and quality of garlic bulbs are influenced by environmental factors and agricultural practices, such as planting time and spacing. Temperature, photoperiod, and soil conditions play crucial roles in garlic cultivation. Proper management of these factors is essential to optimize growth, maximize yield, and enhance the quality of garlic bulbs. Determining the ideal spacing between plants is crucial to achieving the best results. While wider spacing promotes larger bulb size, closer spacing increases overall yield by accommodating more plants per unit area. Additionally, selecting appropriate planting dates can significantly impact garlic's growth cycle and final yield. Research on specific planting time and spacing for garlic cv. Anand Kesari can provide valuable insights for farmers in Gujarat to enhance their garlic production.

2. MATERIALS AND METHODS

The field experiment was carried out during *rabi* season of the year 2022-23 at Horticulture Research Farm, Department of Horticulture, B.A. College of Agriculture, Anand Agricultural University, Anand. It was laid out in Randomized Block Design concept (Factorial). The soil of the experimental plot was sandy loam type. There were three levels of planting time viz., P₁ – 1st week of October, P₂ – 3rd week of October and P₃ – 1st week of November and three levels of spacing viz., S₁ - 15 x 10 cm, S₂ – 15 x 15 cm and S₃ – 20 x 10 cm. Recommended dose of farm yard manure and NPK fertilizers were given as common dose in all the treatments.

2.1 Yield Parameters

The weight of bulb was taken in grams on average basis. Bulb diameter were taken with the help of vernier caliper in centimeters. Bulb yield was taken on plot basis and then converted into quintals per hectare.

2.2 Quality Parameters

Chemical analysis was done using the standard procedures. T.S.S (⁰Brix) was recorded with Ernma pocket hand refractometer and the sulphur content of the bulb (%) was recorded using the Turbidimetric method with a Spectrophotometer.

3. RESULTS AND DISCUSSION

The planting time P₂ (3rd week of October) gave significantly highest bulb diameter (4.46 cm), which was found to be equal with the planting time P₁ (1st week of October) by recording the bulb diameter of (4.18 cm). The greater number of plant height and number of leaves, the larger bulb diameter. Similar results were reported by Nourbakhsh and Cramer [4] in onion and Saikia et al. [5], El-Magd [6] in broccoli.

The bulb diameter was found to be significantly highest at spacing S₂ (15 x 15 cm) i.e. (4.42 cm), it was equal with the spacing S₃ (20 x 10 cm) i.e. (4.25 cm). Each plant in the crop that was more widely spaced out had plenty of space around it, which prevented it from competing with other plants for food and nutrients. As a consequence, each plant performed better in terms of its own personality. The findings align with those of Gaikwad et al. [7], Nasir [8], Kahsay et al. (2014), Misra et al., [9] and Chanu et al. [10] in garlic and Biru [11] in shallot. The interaction effect of planting time and spacing evidenced significantly maximum bulb diameter in P₁S₂ (Planting in 1st week of October at spacing 15 x 15 cm) i.e. (4.84 cm), it was at par with P₂S₁

(Planting in 3rd week of October at spacing 15 x 10 cm) recording the bulb diameter of (4.79 cm) and P₂S₂ (Planting in 3rd week of October at spacing 15 x 15 cm) recording the bulb diameter of (4.27 cm). The significantly minimum bulb diameter was noted in P₁S₁ (Planting in 1st week of October at spacing 15 x 10 cm) *i.e.* (3.82 cm).

The planting time showed significant effect on the weight of bulb, the planting time P₂ (3rd week of October) gave significantly highest weight of bulb (12.29 g).

The increase in bulb weight might be due to better vegetative growth like plant height, number of leaves/plants there by increasing the sink size in terms of bulb size. Thus, robust growth leads to production of more leaves, which resulted in synthesis of more photosynthates and increased accumulation of carbohydrates and other metabolites which ultimately determined the weight of bulbs. These results are in conformity with the findings of Teshale and Tekeste [12], Negero [13], Savale et al. [14] in garlic. The weight of bulb was found to be significantly highest at spacing S₂ (15 x 15 cm) being (12.01 g). This might be due that a plant would be able to get more nutrients and air space with a wider spacing or a lower plant density. The findings aligned with that given by Singh and Singh [15] and Poovamma et al. [16]. The interaction effect of planting time and spacing evidenced significantly maximum weight of bulb in P₂S₃ (Planting in 3rd week of October at spacing 20 x 10 cm) *i.e.* (13.61 g), it was at par with P₁S₁ (Planting in 1st week of October at spacing 15 x 15 cm) recording the weight of bulb of (13.42 g). However, the minimum weight of bulb (9.58 g) was recorded in P₃S₁ (Planting in 1st week of November at spacing 15 x 10 cm). This might be as a result of the optimal plant densities and planting dates. Muhammad Jamroz et al. [17] in garlic.

The planting time P₂ (3rd week of October) gave significantly highest yield (1.93 kg/plot). Decrease in yield was observed in early planting which may be due to poor growth and development of plant. As yield potential of a plant depends upon the amount of vegetative growth made before bulbing commences. Number of workers reported the beneficial effects of date of planting in garlic on yield, *viz.* Savale et al. [18]. The yield was found to be significantly highest

(1.95 kg/plot) at spacing S₁ (15 x 10 cm). Similar results were reported by Mekonnen and Gadisa [19] and Hadiawati and Nazam [20] in garlic, Shock et al. [21] in onion. The most intriguing finding was that the bigger bulb size and better morphological character development at S₂ (15 cm x 15 cm) spacing were unable to offset the yield decline caused by the decreased plant population per unit area, Kun et al. [22] (Planting time 1st week of November with spacing 20 x 10 cm). Higher yield in 3rd week of October and closer spacing is because of more plants per hectare and vigorous growth of plants in early planting dates, and it was noticed that yield decreased as the spacing between plants increased and delayed in planting. The observation is in agreement with the findings of Devulkar et al. [18], Vidya et al. [23] in garlic and Kahsay et al. [24] in onion.

Different planting time has a significant effect on the yield of garlic. Planting at P₂ (3rd week of October) showed significantly the highest yield (60.7 q/ha). The yield was found to be significantly the highest (59.2 q/ha) at spacing S₁ (15 cm x 10 cm) recorded the highest yield (t/ha), despite having low bulb weight, diameter, and cloves per bulb. Consequently, yield was lower at low plant populations due to narrow spacing. Moravcevic et al. [25] and Alam et al. [26] gave similar outcome that the production was best with the narrower planting, but the bulb quality was lowest.

The interaction effect of planting time and spacing depicted significantly maximum yield of garlic in P₂S₁ (Planting time at 3rd week of October with spacing 15x 10 cm) *i.e.* (70.93 q/ha), minimum yield (44.99 q/ha) was found in P₃S₂ (Planting time 1st week of November with spacing 15 x15 cm).

The effect of different time of planting time and spacing was found non-significant with respect to TSS and the interaction effect of different planting time and spacing was found non-significant with respect to Total Soluble Solids (TSS). The effect of different time of planting time and spacing was found non-significant with respect to sulphur content of the bulb and interaction effect of different planting time and spacing was found non-significant with respect to sulphur content of the bulb.

Table 1a. Effect of planting time and spacing on yield attributes of garlic

Treatment	Weight of bulb (g)	Bulb diameter (cm)	Yield (kg/plot)	Yield (q/ha)
Planting time (P)				
P ₁ - 1 st week of October	11.14	4.17	1.73	54.43
P ₂ - 3 rd week of October	12.29	4.46	1.93	60.7
P ₃ - 1 st week of November	9.91	4.00	1.49	46.96
S. Em±	0.24	0.11	0.061	1.91
CD (P = 0.05)	0.72	0.33	0.182	5.72
Spacing (S)				
S ₁ - 15 x 10 cm	10.08	3.96	1.95	59.2
S ₂ - 15 x 15 cm	12.01	4.42	1.56	49.43
S ₃ - 20 x 10 cm	11.25	4.25	1.65	53.46
S. Em±	0.30	0.11	0.061	1.91
CD (P = 0.05)	0.89	0.33	0.182	5.72
S. Em±	0.51	0.19	0.11	3.31
CD (P = 0.05)	Sig	Sig	Sig	Sig
CV (%)	8.57	7.87	10.60	10.60

Table 1b. Interaction effect of planting time and spacing on yield attributes of garlic

Spacing Planting time	Weight of bulb (g)			Bulb diameter (cm)			Yield (kg/plot)			Yield (q/ha)		
	S ₁	S ₂	S ₁	S ₂	S ₃	S ₁	S ₁	S ₂	S ₁	S ₂	S ₃	S ₁
P ₁	9.75	13.42	10.25	3.78	4.79	3.93	1.91	1.74	1.54	57.8	55.36	50.13
P ₂	10.91	12.34	13.61	4.26	4.27	4.84	2.34	1.48	1.98	70.93	46.93	64.25
P ₃	9.58	10.26	9.89	3.82	4.19	3.98	1.61	1.45	1.42	48.87	46.01	46
S. Em±	0.41			0.19			0.11			3.31		
CD (P = 0.05)	1.24			0.57			0.32			9.91		
CV (%)	8.57			7.87			10.60			10.60		

Table 2. Effect of planting time and spacing on quality attribute of garlic

Treatment	TSS (° Brix)	Sulphur content of the bulb (%)
Planting time (P)		
P ₁ - 1 st week of October	20.65	0.1331
P ₂ - 3 rd week of October	20.80	0.1339
P ₃ - 1 st week of November	19.58	0.1274
S. Em±	0.44	0.0024
CD (P = 0.05)	NS	NS
Spacing (S)		
S ₁ - 15 x 10 cm	20.65	0.1339
S ₂ - 15 x 15 cm	20.58	0.1320
S ₃ - 20 x 10 cm	19.79	0.1285
S. Em±	0.44	0.0024
CD (P = 0.05)	NS	NS
P x S Interaction		
S. Em±	0.77	0.0042
CD (P = 0.05)	NS	NS
CV (%)	6.54	5.54

4. CONCLUSION

Based on the preceding analysis, it can be deduced that the optimal results for garlic cultivation were achieved by planting during the third week of October, which led to enhanced vegetative growth characteristics and greater yield. The utilization of a spacing configuration of 15 x 10 cm resulted in the highest yield due to the increased plant population per hectare. Consequently, the most favorable combination for maximizing yield without compromising quality involved the cultivation of the Anand Kesari garlic variety in the third week of October and spacing it at 15 x 10 cm

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Kumar A, Singh H, Kumar A. Effect of NPK through Chemical Fertilizers Combination with Farm Yard Manure (FYM) and Bio-fertilizers on Yield Attributes of Onion (*Allium cepa* L.) in Jaipur Region of Rajasthan, India. J. Exp. Agric. Int. 2024; 46(5):387-98.
[Accesson: 2024 May 21];
Available:<https://journaljeai.com/index.php/JEAI/article/view/2389>
2. Shinde S, Gaikwad RT, Khaire PB. Effect of Time of Transplanting and Spacing on Tulsi (*Ocimum sanctum*). J. Adv. Biol. Biotechnol. 2024;27(5):464-70.
[Accesson: 2024 May 21];
Available:<https://journaljabb.com/index.php/JABB/article/view/807>
3. Zulkarnain Z, Tapingkae T, Taji A. Applications of in vitro techniques in plant breeding. Advances in plant breeding strategies: Breeding, biotechnology and molecular tools. 2015:293-328.
4. Nourbakhsh SS, Cramer CS. Onion plant size measurements as predictors for onion bulb size. Horticulturae. 2022;8(8):682.
5. Saikia BR, Phookan DB, Brahma S. Effect of time of planting and planting densities on growth, yield and economic production of broccoli (*Brassica oleracea* (L.) var. italica Plenck) cv. Pusa broccoli KTS-1. Journal of Hill Agriculture. 2010; 1(2): 135-139.
6. El-Magd MMA. Evaluation of some broccoli cultivars growth, head yield and quality under different planting dates. Journal of Applied Sciences Research. 2013;9(11): 5730-5736.
7. Gaikwad DG, Shinde SJ, Panchal KN, Moklikar MS. Studies on influence response of plant density on growth and yield of garlic (*Allium sativum* L.). International Journal of Current Microbiology and Applied Sciences. 2018; 6:2891-2897.
8. Nasir S. Review on effect of garlic clove weight on yield and yield components of garlic (*Allium sativum* L.). Academic Research Journal of Agricultural Science and Research. 2018;6(9):554-557.
9. Misra AD, Kumar A, Meitei WI. Effect of spacing and planting time on growth and yield of onion var. N-53 under Manipur Himalayas. Indian J. Hort. 2014;71(2):207-210.
10. Chanu KI, Devi AB, Singh UC, Devi NS, Singh SNG. Effect of different levels of nitrogen and spacing on growth and yield of common onion (*Allium cepa* L.) cv. Prema 178 under Manipur condition. The Pharma Innovation Journal. 2022;11(4): 1268-1274
11. Biru FN. Effect of spacing and nitrogen fertilizer on the yield and yield component of shallot (*Allium ascalonium* L.). Journal of Agronomy. 2015;14(4):220.
12. Teshale M, Tekeste N. Growth and yield response of garlic to intra-row spacing and variety at selekeleka, Northern Ethiopia. The Open Biotechnology Journal. 2021;15(1).
13. Negero FW. Yield and yield components of potato (*Solanum tuberosum* L.) as influenced by planting density and rate of nitrogen application at Holeta, West Oromia region of Ethiopia. African Journal of Agricultural Research. 2017; 12(26): 2242-2254.
14. Savale SV, Patel BK, Nanaware PR, More SG. Effect of different methods and time of planting on growth, yield and quality of garlic (*Allium sativum* L.). Ecology,

- Environment and Conservation Paper. 2015;21(3):1431-1435.
15. Singh PK, Singh DR. Effect of spacing and levels of nitrogen on growth and yield of garlic. Indian Journal of Horticulture. 2010; 67(4):305-307.
 16. Poovamma BC, Devi AB, Singh UC, Herojit A. Influence of different levels of planting time and spacing on growth and yield of multiplier onion (*Allium cepa* L. var. aggregatum Don.) Cv. Meitei Tilhou under Manipur condition. IJCS. 2020;8(6):2653-2658.
 17. Muhammad Jamroz, Muhammad Ishtiaq, Nisar Naeem Niaz, Muhammad Buzerg Jamiher & Javed Iqbal. Effects of different planting dates and spacing on growth and yield of garlic cv. Bianco. On line Journal of Biological Sciences. 2001;1(4):206-208.
 18. Devulkar NG, Bhanderi DR, More SJ, Jeth BA. Optimization of yield and growth in onion through spacing and time of planting. Green farming International Journal. 2015;6(2):305-307.
 19. Mekonnen A, Gadisa N. Agronomic practices for improving garlic (*Allium sativum* L.) production and productivity in Ethiopia Review. World Journal of Agricultural Sciences. 2021;17 (6):469-477.
 20. Hadiawati L, Nazam M. Effect of spacing on growth and yield of Sangga Sembalun garlic variety of Sembalun Higland West Lombok. In IOP Conference Series: Earth and Environmental Science. 2021;637(1): 12065.
 21. Shock CC, Feibert EB, Riveira A, Saunders LD. Response of onion yield, grade, and financial return to plant population and irrigation system. HortScience. 2015;50(9):1312-1318.
 22. Kun X, Junhua Y, Hong QL. The influence of seed cloves and plant density on garlic yield. Journal of Shandon Agricultural University. 1999;30(4):409412.
 23. Vidya G, Padma M, Rajkumar M. Effect of planting time and plant densities on yield, quality and cost of production in garlic (*Allium sativum* L.) cv. Jamnagar. Asian Journal of Horticulture. 2013;8(2):552-555.
 24. Kahsay Y, Belew D, Abay F. Effect of intra-row spacing on yield and quality of some onion varieties (*Allium cepa* L.) at Aksum, Northern Ethiopia. African Journal of Plant Science. 2014;7(12):613-622.
 25. Moravcevic D, Bjelic V, Moravcevic M, Gyoždanovic J, Varga, Beatovic D, Jelacic S. The effect of plant density on bulb quality and yield of spring garlic (*Allium sativum* L.). 46th Croatian and 6th International Symposium on Agriculture. Opatija Croatia. 2011;554-557.
 26. Alam MS, Rahim MA, Bhuyan MAH, Simonand PW, Malek MA. Effect of spacing on growth and yield of two lines of garlic under dry land condition. Journal of Agroforestry and Environment. 2010;4: 151-154.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/117874>