



# Effect of Crop Establishment, Intermittent Submergence and Precision Nitrogen Management on Growth and Yield of Rice

Nihal Chandra Mahajan <sup>a++\*</sup>, Rajesh Kumar Singh <sup>a#</sup>,  
Anurag Upadhyay <sup>a++</sup>, Ram Kumar Singh <sup>a†</sup>,  
Amitava Rakshit <sup>b#</sup> and V. K. Tripathi <sup>c‡</sup>

<sup>a</sup> Department of Agronomy, Institute of Agricultural Sciences, Banaras Hindu University, Varanasi, 221005, U.P., India.

<sup>b</sup> Department of Soil Science & Agricultural Chemistry, Institute of Agricultural Sciences, Banaras Hindu University, Varanasi, 221005, U.P., India.

<sup>c</sup> Department of Farm Engineering, Institute of Agricultural Sciences, Banaras Hindu University, Varanasi, 221005, U.P., India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/ijpss/2024/v36i74789>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/119319>

Original Research Article

Received: 25/04/2024

Accepted: 27/06/2024

Published: 28/06/2024

<sup>++</sup> Research Scholar;

<sup>#</sup> Associate Professor;

<sup>†</sup> Senior Professor;

<sup>‡</sup> Assistant Professor;

\*Corresponding author: E-mail: [nihalchandra1996@gmail.com](mailto:nihalchandra1996@gmail.com);

**Cite as:** Mahajan, Nihal Chandra, Rajesh Kumar Singh, Anurag Upadhyay, Ram Kumar Singh, Amitava Rakshit, and V. K. Tripathi. 2024. "Effect of Crop Establishment, Intermittent Submergence and Precision Nitrogen Management on Growth and Yield of Rice". *International Journal of Plant & Soil Science* 36 (7):761-70. <https://doi.org/10.9734/ijpss/2024/v36i74789>.

## ABSTRACT

The traditional production practices of rice are water-wasting, expensive, labor-intensive, energy-consuming, and time-consuming, making rice cultivation less profitable. Effective water and nitrogen management practices are essential for better rice growth and yield. Keeping these facts in view, the present investigation was conducted at the Agricultural Research Farm, Institute of Agricultural Sciences, Banaras Hindu University, Varanasi during *kharif* season of 2019 and 2020. The experimental soil was Gangetic alluvial sandy clay loam. The experiment was laid out in a split-plot design during both years with three replications. Crop establishment (as factor A) and Irrigation scheduling (as factor B) were allocated to the main plot, and nitrogen management in subplots. In main plot, two crop establishment methods (Factor A) *i.e.* CE<sub>1</sub>: Puddle Transplanted Rice (PTR) and CE<sub>2</sub>: Direct Seeded Rice (DSR) and three irrigation regime (Factor B), *i.e.* I<sub>1</sub>: Continuously submerged (CS) of 5 ± 2 cm depth, I<sub>2</sub>: Intermittent submergence (IS) of 5 ± 2 cm and irrigation after five days of disappearance of water from the soil surface & I<sub>3</sub>: Intermittent submergence of 5 ± 2 cm and irrigation after ten days of water disappearance from the soil surface were taken. In sub plot treatments, four nitrogen management practices *i.e.* N<sub>1</sub>: recommended dose of nitrogen (RDN), N<sub>2</sub>: LCC threshold ≤ 4, N<sub>3</sub>: SPAD 30 & N<sub>4</sub>: Rice-Wheat Crop Manager recommendation. Results showed that Direct Seeded Rice (DSR) and Intermittent submergence of 5 ± 2 cm and irrigation after 5 days of disappearance of water among the main plot and nitrogen recommendation based on Rice-Wheat Crop Manager (RWCM) recorded higher plant height, number of tillers, chlorophyll content and grain yield of rice over other treatment. DSR recorded a higher grain yield by 9.5 % over Conventional method, whereas I<sub>2</sub> recorded 7.5 % more grain yield over I<sub>1</sub> and RWCM (N<sub>4</sub>) recorded 17% more grain yield over Recommended Nitrogen Dose (RDN). It was concluded that Direct-Seeded Rice with intermittent submergence and precision nitrogen management improved the growth and yield of rice under the climatic conditions of Eastern Uttar Pradesh. This approach offers a more profitable and sustainable method for rice cultivation.

**Keywords:** Direct seeded rice (DSR); puddle transplanted rice (PTR); intermittent submergence; precision agriculture.

## 1. INTRODUCTION

“Rice is the world's second-largest cultivated after wheat. In India, rice occupies 45 million hectares with a production of 127.9 million tonnes and an average yield of 2.84 t ha<sup>-1</sup>” [1]. “In India, it accounts for more than 55 per cent of cereal production, providing direct employment to 70% of the people in rural areas. As it is the staple food for more than 65% of the population, our national food security hinges on the growth and stability of its production” [2].

“There are several ways for rice cultivation worldwide: wet seeding, which involves sowing pre-germinated seeds in wet puddled soils; and transplanting, which involves replanting rice seedlings grown in a nursery to puddled soils” [3]. “Generally, in India, rice is grown as a transplanted crop; it is traditionally grown by transplanting into puddled soil (land preparation with wet tillage) in June/July” [4]. “This production system not only leads to wastage of water but it is also costly, requires a significant labour input (300–350 man-h ha<sup>-1</sup>), energy-intensive (2016–2390 MJ ha<sup>-1</sup>), and it is a cumbersome and time-consuming process, which makes rice production

less profitable” [5]. “Increasing labour, diesel, and electricity costs are also considered as severe issues” [6]. “All these factors have threatened the productivity and sustainability of rice-based systems and demand a significant shift from the current transplanted rice production system. Aerobic or partially-aerobic rice cultivation techniques like direct-seeding of rice (DSR) are gaining recognition for resource conservation (water in particular) and environmental gains” [7]. “There are three principal methods of direct seeding of rice (DSR): dry seeding (sowing dry seeds into dry soil), wet seeding (sowing pre-germinated seeds on wet puddled soils), and water seeding (seeds are sown into standing water). Direct seeding offers such advantages compared to traditional transplanting as faster planting, reduced labour by up to 60%, less drudgery, earlier crop maturity by 7 to 10 days with less water requirement (12–36%), less methane emission, often higher profitability, and the rising interest in conservation agriculture” [8,6].

“Recent projections suggest that there will be a severe water shortage in the following decades. It is suggested that rice farmers use water more

effectively because of the rise in irrigation water costs and a decrease in return. These necessitate the adoption of water-efficient techniques for rice production to reduce water use in the agricultural sector while maintaining or increasing yield to support a growing population" [9,10]. "Alternate wetting and drying (AWD) are among the most widely promoted water-saving irrigation technique introduced by the International Rice Research Institute (IRRI) to cope with the increasing threat of water scarcity in rice cultivation" [11]. "Under this system, fields are subjected to intermittent flooding (alternate cycles of saturated and unsaturated conditions) where the water of about 2–5 cm is applied at an interval of 1 to 10 days, depending on the soil type and weather conditions, followed by the disappearance of pond water from the soil surface and appearance of visible signs of some fine cracks on the soil surface and then re-flooded up to 5 cm above the surface" [12]. This cycle is repeated until the rice harvest, except for the flowering stage.

As fertilizer N has been generally managed following blanket recommendations consisting of two or three split applications of present rates of the total amount of N, improvement in N use efficiency could only be achieved by a limit. Various factors such as spatial soil N variability, agro-climatic circumstances, time and rate of N fertilizer application, and varietal differences significantly hamper the efficient use of fertilizer N when broad-based general or blanket recommendations are followed. Therefore, effective N management during crop growth is critical to efficient crop production and enhancing crop yield and quality [13,14]. Adjusting N input to an economically and ecologically compatible level would require quantitative information on the N status of rice plants. An ideal indicator of crop N status should be able to detect deficiencies and excesses of N supply and provide a fast diagnosis to allow dressing correction for efficient fertilizer management during crop growth [15].

## 2. MATERIALS AND METHODS

The present investigation was conducted at the Agricultural Research Farm, Institute of Agricultural Sciences, Banaras Hindu University, Varanasi during *kharif* season of 2019 and 2020, in South-eastern part of Varanasi city of India (25°18'N latitude and 83°31'E longitude at an altitude of 128.93 meter above sea level). The experimental soil was Gangetic alluvial sandy clay loam. The experiment was laid out in a split-

plot design during both years with three replications. Crop establishment (as factor A) and Irrigation scheduling (as factor B) were allocated to the main plot, and nitrogen management to subplots. In main plot, two crop establishment methods (Factor A) *i.e.* CE<sub>1</sub>: Puddle Transplanted Rice (PTR) & CE<sub>2</sub>: Direct Seeded Rice (DSR) and three irrigation scheduling (Factor B) *i.e.* I<sub>1</sub>: Continuously submerged (CS) of 5 ± 2 cm depth, I<sub>2</sub>: Intermittent submergence (IS) of 5 ± 2 cm and irrigation after five days of disappearance of water from the soil surface & I<sub>3</sub>: Intermittent submergence of 5 ± 2 cm and irrigation after ten days of water disappearance from the soil surface were taken. In sub plot treatments four nitrogen management practices *i.e.* N<sub>1</sub>: recommended dose of nitrogen (RDN), N<sub>2</sub>: LCC threshold ≤ 4, N<sub>3</sub>: Soil plant analysis development (SPAD 30) & N<sub>4</sub>: Rice-Wheat Crop Manager Recommendation. A buffer of 1.0 m width surrounded each main plot, whereas the subplot was surrounded by 1.0 m width to protect the plots from accidental irrigation and water gain through seepage. The application of nutrients to the test crop was made based on recommendations for uniformly using phosphorous, potassium, and zinc (60-60-5 kg ha<sup>-1</sup>) for all the treatments. The sources used for nitrogen, phosphorous, potassium, and zinc were urea (46% N), single super phosphate (16% P<sub>2</sub>O<sub>5</sub>, 12% S, and 21% Ca), muriate of potash (60% K<sub>2</sub>O), and zinc sulphate monohydrate (33%), respectively. The total diammonium phosphate and potash dose were applied when sowing/transplanting as basal. Zinc was applied through two foliar sprays. Other agronomic practices were carried out as per the state recommendations. Nitrogen was applied as per treatments through leaf colour chart (LCC) threshold ≤ 4, SPAD 30, and Rice-Wheat crop manager recommendation, whereas in the recommended dose of nitrogen (120 kg ha<sup>-1</sup>), one-fourth dose was applied as basal, the half dose was applied at the maximum tillering stage, and the remaining one-fourth dose at the panicle initiation stage.

## 3. RESULTS

### 3.1 Growth Parameters

#### 3.1.1 Plant height

Direct-seeded rice recorded significant maximum plant height in comparison to puddled transplanted rice at 30, 60, and 90 days after sowing and at harvest stages. Marked variations

in plant height were recorded due to different water management practices at all the growth stages during both years. At 30 Days after sowing/Days after transplanting, among the different water management practices,  $I_2$  (Intermittent submergence of  $5 \pm 2$  cm and irrigation after 5 days of disappearance of water from the soil surface) recorded significantly higher plant height (37.86 and 39.77 cm) over  $I_3$  (Intermittent submergence of  $5 \pm 2$  cm and irrigation after 10 days of disappearance of water from the soil surface) and statistically at par with  $I_1$  (continuously submerged at 5 cm depth) (Table 1). A similar trend was observed at 60, 90 DAS/DAT and at the harvest stage during both the years of study.

Nitrogen management significantly affected plant height at all the growth stages. Use of the Rice-Wheat crop manager's recommendation, being statistically at par with and LCC threshold  $\leq 4$ , brought significantly higher plant height over the required dose of nitrogen and SPAD-30 based Nitrogen management during both years in all growth stages. The minimum plant height was observed with the treatment in which the recommended dose of nitrogen was applied for both years of investigation at all the growth stages.

### 3.1.2 Number of tillers

Among the crop establishment methods, direct-seeded rice recorded a maximum no. of tillers (35.17 and 35.60) compared to PTR (29.88 and 30.80) at 30 DAS/DAT during both years, respectively. With respect to no. of tillers, similar trends were observed at 60, 90 DAS/DAT and the harvest stage during the research period. Water management practices also influenced the number of tillers at all growth stages in both experimental years. At 30 DAS/DAT, the water management practice  $I_2$  (intermittent submergence of  $5 \pm 2$  cm and irrigation after 5 days of disappearance of water from the soil surface) resulted in a significantly higher number of tillers (35.21 and 35.98) compared to  $I_3$  (intermittent submergence of  $5 \pm 2$  cm and irrigation after 10 days of disappearance of water from the soil surface) and statistically at par with  $I_1$  (continuously submerged at 5 cm depth) in both years. At 60, 90 DAS/DAT, and at the harvest, similar trends were observed in both study years (Table 2). Further, among the different nitrogen management options, the highest number of tillers was recorded using the Rice-Wheat crop manager. Still, it was comparable to LCC threshold  $\leq 4$ , all bringing

significant improvement over RDN and SPAD 30-based nitrogen management. Moreover, RDN had a minimum no. of tillers at all the growth stages during each year of study.

### 3.1.3 Chlorophyll content (SPAD Value)

Chlorophyll content in the leaves (SPAD value) did not experience a significant change due to the effect of crop establishment and nitrogen management during both years. Irrigation scheduling significantly affected the chlorophyll content in leaves (SPAD value) at 60 DAS/DAT observation while non-significant at 30 and 90 DAS/DAT. Irrigation scheduling at Intermittent submergence of  $5 \pm 2$  cm and irrigation after 5 days of disappearance of water from the soil surface ( $I_2$ ) resulted in the highest chlorophyll content in leaves at 60 DAS/DAT (Table 3). The lowest chlorophyll content was recorded under Intermittent submergence of  $5 \pm 2$  cm and irrigation after 10 days of disappearance of water from the soil surface ( $I_3$ ).

### 3.2 Grain Yield

The Direct-seeded rice displayed superiority over puddled transplanted rice in terms of grain yield during both the years of study. DSR recorded a grain yield of 46.61 and 48.22 quintal  $ha^{-1}$  in the first and second year, respectively, while PTR with a lower grain of 42.54 and 43.83 quintal  $ha^{-1}$ , respectively. The effect of Irrigation scheduling was found significant on the grain yield of rice during both years. Water management with intermittent submergence of  $5 \pm 2$  cm and irrigation after 5 days of disappearance of water from the soil surface ( $I_2$ ) resulted in significantly higher grain yield over all other treatments at test. The highest grain yield (48.70 and 50.15 qt.  $ha^{-1}$  in the first and second year, respectively) was recorded under  $I_2$ , followed by  $I_1$  (45.01 and 46.46 qt.  $ha^{-1}$ , respectively) (Table 3). The lowest yield (40.02 and 41.47 qt.  $ha^{-1}$ , respectively) was observed under intermittent submergence of  $5 \pm 2$  cm and irrigation after 10 days of disappearance of water from the soil surface ( $I_3$ ). A significant influenced of nitrogen management was observed on the grain yield of rice during the experimentation period. Management of nitrogen through Rice-Wheat Crop Manager Recommendation ( $N_4$ ) resulted in significantly higher grain yield over the rest of the management treatments. The highest grain yield of 48.02 and 49.87 qt.  $ha^{-1}$  in the first and second year, respectively was recorded under  $N_4$ , which was 10.22, 3.57 (and 14.91% higher over  $N_3$ ,  $N_2$  and  $N_1$ , respectively).

**Table 1. Effect of crop establishment, irrigation schedule, and precision nitrogen management on plant height (cm) of rice at various crop growth stages**

Treatments	30 DAS/DAT		60 DAS/DAT		90 DAS/DAT		At harvest	
	2019	2020	2019	2020	2019	2020	2019	2020
<b>Main-plots treatment</b>								
<b>Crop establishment</b>								
CE <sub>1</sub> : Puddle Transplanted Rice (PTR)	33.31	35.11	67.70	70.05	99.81	103.55	97.80	99.68
CE <sub>2</sub> : Direct Seeded Rice (DSR)	37.06	38.90	73.21	76.48	108.48	112.71	105.33	108.42
SEm ±	0.93	0.91	1.67	1.63	1.97	1.92	1.93	1.86
LSD ( <i>P</i> =0.05)	2.84	2.79	4.96	4.88	6.21	6.06	6.09	5.94
<b>Irrigation scheduling</b>								
I <sub>1</sub> : Continuously submerged at 5 cm depth	35.78	37.56	71.91	74.60	106.16	110.06	103.46	105.98
I <sub>2</sub> : Intermittent submergence of 5 ± 2 cm and irrigation after 5 days of disappearance of water from the soil surface	37.86	39.77	75.33	78.46	111.44	115.79	108.72	111.40
I <sub>3</sub> : Intermittent submergence of 5 ± 2 cm and irrigation after 10 days of disappearance of water from the soil surface	31.91	33.69	64.13	66.73	94.82	98.53	92.51	94.78
SEm ±	0.84	0.78	1.59	1.53	1.81	1.77	1.77	1.73
LSD ( <i>P</i> =0.05)	2.66	2.36	4.81	4.72	5.20	5.20	5.07	4.98
<b>Sub-plots treatment</b>								
<b>Nitrogen Management</b>								
N <sub>1</sub> : RDN	33.16	34.73	65.91	68.21	97.23	102.51	95.71	97.14
N <sub>2</sub> : LCC threshold ≤ 4	36.15	38.12	73.17	76.10	108.13	110.71	104.59	108.04
N <sub>3</sub> : SPAD 30	34.04	35.69	67.82	70.10	99.96	105.58	97.15	99.87
N <sub>4</sub> : Rice-Wheat Crop Manager Recommendation	37.38	39.46	74.95	78.66	111.25	113.71	108.77	111.16
SEm ±	0.69	0.64	1.44	1.41	1.61	1.57	1.58	1.51
LSD ( <i>P</i> =0.05)	2.11	1.97	4.42	4.37	5.07	4.95	4.97	4.85

**Table 2. Effect of crop establishment, irrigation schedule, and precision nitrogen management on tiller count (per running meter) of rice at various crop growth stages**

Treatments	30 DAS/DAT		60 DAS/DAT		90 DAS/DAT		At harvest	
	2019	2020	2019	2020	2019	2020	2019	2020
<b>Main-plots treatment</b>								
<b>Crop establishment</b>								
CE <sub>1</sub> : Puddle Transplanted Rice (PTR)	29.88	30.80	66.36	67.27	63.70	64.58	61.09	61.96
CE <sub>2</sub> : Direct Seeded Rice (DSR)	35.17	35.60	75.13	78.09	73.25	76.14	71.14	72.77
SEm ±	0.67	0.63	1.32	1.21	1.28	1.20	1.17	1.11
LSD ( <i>P</i> =0.05)	2.13	1.98	4.15	3.82	4.02	3.88	3.68	3.46
<b>Irrigation scheduling</b>								
I <sub>1</sub> : Continuously submerged at 5 cm depth	33.15	33.79	71.96	73.94	69.65	71.58	67.25	68.52
I <sub>2</sub> : Intermittent submergence of 5 ± 2 cm and irrigation after 5 days of disappearance of water from the soil surface	35.21	35.98	76.21	78.40	73.77	75.91	71.27	72.59
I <sub>3</sub> : Intermittent submergence of 5 ± 2 cm and irrigation after 10 days of disappearance of water from the soil surface	29.22	29.83	64.07	65.71	62.01	63.60	59.81	60.97
SEm ±	0.64	0.59	1.19	1.10	1.15	1.07	1.06	1.01
LSD ( <i>P</i> =0.05)	1.82	1.70	3.41	3.15	3.30	3.08	3.21	3.08
<b>Sub-plots treatment</b>								
<b>Nitrogen Management</b>								
N <sub>1</sub> : RDN	30.38	30.68	66.39	68.23	62.80	63.29	60.04	60.43
N <sub>2</sub> : LCC threshold ≤ 4	33.34	34.23	72.40	74.34	71.41	74.05	69.27	70.98
N <sub>3</sub> : SPAD 30	31.63	32.26	68.91	70.78	65.67	66.52	62.95	63.60
N <sub>4</sub> : Rice-Wheat Crop Manager Recommendation	34.77	35.62	75.24	77.38	74.04	77.56	72.19	74.42
SEm ±	0.55	0.51	1.07	0.99	0.95	0.91	1.04	0.98
LSD ( <i>P</i> =0.05)	1.74	1.62	3.39	3.12	2.89	2.83	3.28	3.09

**Table 3. Effect of crop establishment, irrigation schedule, and precision nitrogen management on chlorophyll content in leaves (SPAD value) of rice at various crop growth stages**

Treatments	30 DAS/DAT		60 DAS/DAT		90 DAS/DAT		Grain yield (qt. ha <sup>-1</sup> )	
	2019	2020	2019	2020	2019	2020	2019	2020
<b>Main-plots treatment</b>								
<b>Crop establishment</b>								
CE <sub>1</sub> : Puddle Transplanted Rice (PTR)	30.99	31.85	35.40	36.00	30.80	31.01	42.54	43.83
CE <sub>2</sub> : Direct Seeded Rice (DSR)	31.51	32.39	36.09	36.24	31.35	31.60	46.61	48.22
SEm ±	0.55	0.53	0.58	0.56	0.43	0.41	58	56
LSD (P=0.05)	NS	NS	NS	NS	NS	NS	182	177
<b>Irrigation scheduling</b>								
I <sub>1</sub> : Continuously submerged at 5 cm depth	31.21	32.08	35.74	36.11	31.09	31.27	45.01	46.46
I <sub>2</sub> : Intermittent submergence of 5 ± 2 cm and irrigation after 5 days of disappearance of water from the soil surface	31.97	32.86	36.84	37.23	31.99	32.23	48.70	50.15
I <sub>3</sub> : Intermittent submergence of 5 ± 2 cm and irrigation after 10 days of disappearance of water from the soil surface	30.57	31.42	34.65	35.01	30.14	30.41	40.02	41.47
SEm ±	0.53	0.50	0.56	0.53	0.41	0.38	52	50
LSD (P=0.05)	NS	NS	NS	NS	NS	NS	148	143
<b>Sub-plots treatment</b>								
<b>Nitrogen Management</b>								
N <sub>1</sub> : RDN	31.07	31.94	35.41	35.77	30.61	30.87	40.86	41.76
N <sub>2</sub> : LCC threshold ≤ 4	31.33	32.21	35.85	35.98	31.21	31.43	46.32	47.53
N <sub>3</sub> : SPAD 30	31.18	32.05	35.60	35.96	30.98	31.27	43.11	44.95
N <sub>4</sub> : Rice-Wheat Crop Manager Recommendation	31.41	32.29	36.11	36.51	31.47	31.61	48.02	49.87
SEm ±	0.50	0.47	0.53	0.50	0.38	0.35	47	46
LSD (P=0.05)	NS	NS	NS	NS	NS	NS	149	144

#### 4. DISCUSSION

The final yield of a crop depends largely on the success of growth and development in individual plants, which is influenced by crop management practices such as establishment methods, irrigation and fertilization. Maintaining optimal soil-plant-atmosphere conditions through a combination of agronomic manipulations leads to the sustainable exploitation of crop yield potential within the local environment, specifically leveraging edaphic factors and genotypes to the crop's advantage [16].

DSR outperformed TPR for crop stand, tiller density, and leaf area development. Higher seedling vigour in DDSR resulted to early canopy cover, increased light interception, and higher WUE [17,18]. In TPR, root damage caused by nursery uprooting and transplanting trauma can inhibit early growth and vigour. Yadav et al. [19] found that DSR had greater growth than TPR, which was consistent with our findings. Alternate wetting and drying improved the number of productive tillers, while less water decreased the tillering density and number of productive tillers. The results were consistent with findings of previous studies which showed that AWD has an ability to increase productive tillers and reduces redundant growth by altering leaf angles, improving root health, shoot growth and LAI [20,21]. Optimal nutrient use during SSNM may lead to increased cell division, enlargement, photosynthesis, and protein synthesis, resulting in quantitative increases in plant growth characteristics such as tillering [22,23] found similar results.

The yield of paddy in direct seeding by rice-wheat seeder for all the varieties may be highest because of better root establishment, higher effective tillers and uniform plant geometry. These results were in close conformity with finding of Sharma et al. [24] and Gill et al. [25]. Higher yields under AWD might be attributable to improved grain filling and greater root proliferation to absorb more water and nutrients [21]. Furthermore, AWD raised effective tiller percentage and glucose translocation to grain, decreases spikelet sterility, and boosts grain weight [26]. Post-anthesis AWD improves grain filling rate because to increased enzyme activity, which increases grain production [27]. Improved nutrient delivery throughout rice's active growth and development stages might explain the increase in yield-contributing traits under SSNM-

based treatments [28,23] reported similar results [29,30].

#### 5. CONCLUSION

The traditional production practices of rice not only waste water, but it is also expensive, requiring a substantial manpower input, is energy-intensive, and is a difficult and time-consuming procedure, making rice cultivation less profitable. Effective water and nitrogen management practices are essential requirement for better rice growth and yield. These problems play an important cause for shifting towards direct seeded rice instead of present transplanted rice production system along with alternate wetting drying irrigation and site-specific nutrient management practices. Results showed that Direct Seeded Rice (DSR) and Intermittent submergence of  $5 \pm 2$  cm and irrigation after 5 days of disappearance of water among the main plot and nitrogen recommendation based on Rice-Wheat Crop Manager (RWCM) recorded higher plant height, number of tillers, chlorophyll content and grain yield of rice over other treatment. It can be concluded that DSR with intermittent submergence and precision nitrogen management can improve the growth and yield of rice under climatic condition of Eastern Uttar Pradesh.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Agricultural Statistics at a Glance, Government of India Ministry of Agriculture & Farmers Welfare, Department of Agriculture & Farmers Welfare, Economics & Statistics Division 2022
2. Pathak HP, Nayak AK, Jena M, Singh ON, Samal P, Sharma SG. Rice research for enhancing productivity, profitability and climate resilience. 2018 May 1;527
3. Chen S, Ge Q, Chu G, Xu C, Yan J, Zhang X, Wang D. Seasonal differences in the

- rice grain yield and nitrogen use efficiency response to seedling establishment methods in the Middle and Lower reaches of the Yangtze River in China. *Field Crops Research*. 2017 Apr 1;205:157-69.
4. Mahajan G, Chauhan BS, Timsina J, Singh PP, Singh K. Crop performance and water- and nitrogen-use efficiencies in dry-seeded rice in response to irrigation and fertilizer amounts in northwest India. *Field Crops Research*. 2012 Aug 12;134:59-70.
  5. Bhatt R, Kukal SS, Busari MA, Arora S, Yadav M. Sustainability issues on rice-wheat cropping system. *International Soil and Water Conservation Research*. 2016 Mar 1;4(1):64-74.
  6. Chauhan BS, Mahajan G, Sardana V, Timsina J, Jat ML. Productivity and sustainability of the rice-wheat cropping system in the Indo-Gangetic Plains of the Indian subcontinent: problems, opportunities, and strategies. *Advances in agronomy*. 2012 Jan 1;117:315-69.
  7. Mishra AK, Khanal AR, Pede VO. Is direct seeded rice a boon for economic performance? Empirical evidence from India. *Food Policy*. 2017 Dec 1;73:10-8.
  8. Mahajan G, Chauhan BS, Gill MS. Dry-seeded rice culture in Punjab State of India: lessons learned from farmers. *Field Crops Research*. 2013 Mar 20;144: 89-99.
  9. Saha S, Singh YV, Gaiind S, Kumar D. Water productivity and nutrient status of rice soil in response to cultivation techniques and nitrogen fertilization. *Paddy and water environment*. 2015 Oct;13:443-53.
  10. Brar AS, Buttar GS, Sharma R. Water and energy productivity of rice as influenced by duration of cultivars, dates of transplanting and irrigation regime in north-western India. *Paddy and water environment*. 2018 Oct;16:655-63.
  11. Yang J, Zhou Q, Zhang J. Moderate wetting and drying increases rice yield and reduces water use, grain arsenic level, and methane emission. *The Crop Journal*. 2017 Apr 1;5(2):151-8.
  12. Lampayan RM, Rejesus RM, Singleton GR, Bouman BA. Adoption and economics of alternate wetting and drying water management for irrigated lowland rice. *Field Crops Research*. 2015 Jan 1;170:95-108.
  13. Peng S, Buresh RJ, Huang J, Zhong X, Zou Y, Yang J, Wang G, Liu Y, Hu R, Tang Q, Cui K. Improving nitrogen fertilization in rice by sitespecific N management. A review. *Agronomy for sustainable development*. 2010 Sep;30:649-56.
  14. Zhao B, Ata-Ul-Karim ST, Liu Z, Ning D, Xiao J, Liu Z, Qin A, Nan J, Duan A. Development of a critical nitrogen dilution curve based on leaf dry matter for summer maize. *Field Crops Research*. 2017 Jul 1;208:60-8.
  15. Yao X, Ata-Ul-Karim ST, Zhu Y, Tian Y, Liu X, Cao W. Development of critical nitrogen dilution curve in rice based on leaf dry matter. *European Journal of Agronomy*. 2014 Apr 1;55:20-8.
  16. Bailey-Serres J, Parker JE, Ainsworth EA, Oldroyd GE, Schroeder JI. Genetic strategies for improving crop yields. *Nature*. 2019 Nov 7;575(7781):109-18.
  17. Farooq MK, Siddique KH, Rehman H, Aziz T, Lee DJ, Wahid A. Rice direct seeding: Experiences, challenges and opportunities. *Soil and Tillage Research*. 2011 Jan 1;111(2):87-98.
  18. Anwar P, Juraimi AS, Puteh A, Selamat A, Man A, Hakim A. Seeding method and rate influence on weed suppression in aerobic rice. *African Journal of Biotechnology*. 2011;10(68):15259-71.
  19. Sudhir-Yadav SY, Humphreys E, Kukal SS, Gill G, Rangarajan R. Effect of water management on dry seeded and puddled transplanted rice: Part 2: Water balance and water productivity. 2011.
  20. Norton GJ, Shafaei M, Travis AJ, Deacon CM, Danku J, Pond D, Cochrane N, Lockhart K, Salt D, Zhang H, Dodd IC. Impact of alternate wetting and drying on rice physiology, grain production, and grain quality. *Field Crops Research*. 2017 Apr 1;205:1-3.
  21. Pascual VJ, Wang YM. Utilizing rainfall and alternate wetting and drying irrigation for high water productivity in irrigated lowland paddy rice in southern Taiwan. *Plant Production Science*. 2017 Jan 2;20(1):24-35.
  22. Singh H, Singh UP, Singh SP, Singh Y. Effect of crop establishment and nutrient management on productivity and profitability of rice under rice-wheat system. *Int. J. Chem. Stud*. 2018;6(3):165-70.
  23. Shahi UP, Singh VK, Kumar A, Upadhyay PK, Rai PK. Site-specific nutrient management: Impact on productivity, nutrient uptake and economics of rice-

- wheat system. Indian J. Agric. Sci. 2022; 92:195-8.
24. Sharma S, Rout KK, Khanda CM, Tripathi R, Shahid M, Nayak A, Satpathy S, Banik NC, Iftikar W, Parida N, Kumar V. Field-specific nutrient management using Rice Crop Manager decision support tool in Odisha, India. Field Crops Research. 2019 Sep 1;241:107578.
25. Gill JS, Walia SS, Gill RS. Direct seeded rice: An alternative rice establishment technique in north-west India—A review. International Journal of Advanced Research. 2014;2(3):375-86.
26. Yang J, Zhang J. Crop management techniques to enhance harvest index in rice. Journal of experimental botany. 2010 Jul 1;61(12):3177-89.
27. Zhang H, Li H, Yuan L, Wang Z, Yang J, Zhang J. Post-anthesis alternate wetting and moderate soil drying enhances activities of key enzymes in sucrose-to-starch conversion in inferior spikelets of rice. Journal of Experimental Botany. 2012 Jan 1;63(1):215-27.
28. Singh VK, Dwivedi BS, Tiwari KN, Majumdar K, Rani M, Singh SK, Timsina J. Optimizing nutrient management strategies for rice–wheat system in the Indo-Gangetic Plains of India and adjacent region for higher productivity, nutrient use efficiency and profits. Field Crops Research. 2014; 164:30-44.
29. Girsang SS, Quilty JR, Correa Jr TQ, Sanchez PB, Buresh RJ. Rice yield and relationships to soil properties for production using overhead sprinkler irrigation without soil submergence. Geoderma. 2019 Oct 15;352:277-88.
30. Mohanty S, Swain CK, Sethi SK, Dalai PC, Bhattacharyya P, Kumar A, Tripathi R, Shahid M, Panda BB, Kumar U, Lal B. Crop establishment and nitrogen management affect greenhouse gas emission and biological activity in tropical rice production. Ecological Engineering. 2017 Jul 1;104:80-98.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/119319>