



# Unraveling Promising Genetic Variability Associated with Strong Culm and Yield Parameters in Inter Sub-Specific Cross Derived Recombinant Inbred Lines in Rice (*Oryza sativa* L.)

Mamidi Akshay <sup>a,b</sup>, Lavuri Krishna <sup>c</sup>, Jyothi Badri <sup>b\*</sup>,  
Kalyani M. Barbadikar <sup>b</sup> and D. Sanjeeva Rao <sup>b</sup>

<sup>a</sup> Department of Genetics and Plant Breeding, College of Agriculture, Professor Jayashankar Telangana State Agricultural University (PJTSAU), Rajendranagar, Hyderabad -500030, India.

<sup>b</sup> ICAR-Indian Institute of Rice Research (ICAR-IIRR), Rajendranagar, Hyderabad -500030, India.

<sup>c</sup> Institute of Rice Research (IRR), PJTSAU, Rajendranagar, Hyderabad -500030, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/ijpss/2024/v36i95019>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/119549>

**Original Research Article**

**Received: 12/05/2024**

**Accepted: 16/07/2024**

**Published: 20/09/2024**

\*Corresponding author: E-mail: [jyothirishik@gmail.com](mailto:jyothirishik@gmail.com);

**Cite as:** Akshay, Mamidi, Lavuri Krishna, Jyothi Badri, Kalyani M. Barbadikar, and D. Sanjeeva Rao. 2024. "Unraveling Promising Genetic Variability Associated With Strong Culm and Yield Parameters in Inter Sub-Specific Cross Derived Recombinant Inbred Lines in Rice (*Oryza Sativa* L.)". *International Journal of Plant & Soil Science* 36 (9):700-711. <https://doi.org/10.9734/ijpss/2024/v36i95019>.

## ABSTRACT

Strengthening the rice culm through genetic improvement has been an important target in breeding to abate the lodging problem in rice. The present investigation was attempted to study the variability, genetic parameters and trait association in recombinant inbred lines (RILs) derived from inter sub-specific cross of elite *indica* mega variety 'Swarna' and Tropical *japonica* strong culm accession 'IRGC 39111'. Analysis of Variance (ANOVA) displayed significant variation in RILs for all the traits studied. The investigation revealed high estimates of GCV and PCV along with high heritability and genetic advance as percent of mean (GAM) for all culm strength (CS) traits revealing the role of additive gene effect and selection of these traits shall be effective in rice improvement. Correlation studies revealed that the internode breaking weight (IBW) and bending moment at breaking (M) had a high significant positive correlation with all the CS traits, panicle weight and grain yield. Thus, the direct selection of positively associated traits can improve the lodging resistance in rice. The principal component analysis (PCA) showed that, seven principal components (PCs) out of 23 traits attributed approximately 92.9% of the total cumulative variability. The traits culm thickness (CT), Internode breaking weight (IBW), section modulus (SM), bending stress (BS), Bending moment at breaking (M) contributed maximum to the total diversity and the lines with higher trait values for the aforementioned traits could be utilized as donors to improve the culm strength and its attributing traits in future breeding programs. From this study, it is evident that tropical *japonica* lines can be a great source of genetic diversification for rice improvement.

**Keywords:** Rice; tropical *japonica*; inter sub-specific cross; PCV; GCV; heritability; GAM; correlation; PCA.

## 1. INTRODUCTION

"Rice (*Oryza sativa* L.) is the second most cultivated cereal crop in the world. It is estimated that 40% of the world's population consume rice and is the main stay food for people of Asia. To further enhance the yield of rice varieties amidst climate change and burgeoning population, it is necessary not only to improve yield component traits but also to strengthen the culms bearing heavy panicles. The present deceleration drift in production and yield is a cause of concern and has to be upturned to meet the growing demand" [1]. It is therefore necessary to harness the available resources and knowledge on rice and engage large genetic variability from outside because the available genetic diversity of the cultivated *indica* pool has already been utilized to greater extent. In the evolutionary process, two cultivated sub-species of rice, *indica* and *japonica* have become genetically isolated. This has stifled the genetic diversity within each sub-species [2]. *Indica* sub-species harboring the greatest genetic diversity, is grown across the world and thus has been the focus of most genetic research and utilization. However, the exploration of diversity panels has revealed that *indica* is relatively less diverse than the *japonica* [3]. *Japonica* rice consists of two subtypes: temperate and tropical. Tropical *japonica* (TrJ) initially considered as a third sub-species called *javanica*, represents an intermediary between *indica* and *japonica* and can hybridize with both

*indica* and *japonica*, overcoming the cross-incompatibility barrier between the two sub-species. Consequently, TrJ serves as a bridge for *indica-japonica* hybridization and a valuable source for increasing genetic base in rice [3]. Tropical *japonica* landraces were used in breeding program for new plant type (NPT) rice at IRRI (International rice research institute) to increase the yields of modern *indica* cultivars [4]. New plant type concept is gaining significant momentum in the context of doubling the farmer's income by enhancing the yield potential of the existing cultivars and focus is now on breeding for high yielding lines (Jyothi et al. 2018). In this context, it is imperative to increase culm strength that may provide scope for attaining targeted high yields in rice [5].

Moreover, the introduction of genes *Gn1a* and *APO1* for number of grains [6,7,8]; *DEP1* and *WFP* for panicle architecture [9,10]; *GS3*, *GW2*, and *qSW5* for grain size [11,12,13,4] has resulted in heavier panicles, but the culms supporting those panicles have not improved, and there is the concern of lodging again [14]. "Furthermore, erratic precipitation patterns, severe typhoons and alarming surge in the cyclonic turbulence in India and East Asia are expected to pose imminent threats to rice production. To cope with this situation, enhancing the mechanical strength of the culm is becoming inevitably important to develop the rice varieties having strong culm and lodging

resistance to bear heavy panicles for yield improvement. To understand the physiological and morphological basis of culm strength and improve it further, prior knowledge of the genetic control of mechanisms that regulate culm strength is a prerequisite" [15].

"The presence of adequate genetic variability is regarded as the fundamental pre-requisite to conduct any crop improvement programme. Hence, it is very crucial to understand the species' variability, the nature of character relationships and the role of various traits in increasing rice production through breeding programme" [16]. Indonesian tropical *japonica* landraces are characterized by large panicles, large leaves, strong culms, fewer tillers and vigorous root architecture, contributing to their adaptability and resilience in diverse environments [3]. Being an unexplored germplasm, Tropical *japonica* offers an abundant scope to dwell into and harness the desirable alleles for agronomically important traits like strong culm. To capture these desirable alleles, merging the genetic diversity across sub-species could bring in increased variability and therefore increased heterosis, for which tropical *japonica* would be useful.

"Sometimes phenotypic selection based on their performance may not be effective because these genotypes may perform poor in further segregating generations, so it is essential to select the genotypes based on heritability and genetic advance" [1]. "The genotypic coefficient of variation (GCV) measures the magnitude of genetic variability and reflects the heritable portion of variation. Genetic variability along with heritability estimates would provide the amount of genetic gain expected out of selection" [17]. "Therefore, assessment of variability is essential to formulate selection criteria for the improvement of culm strength and constructive augmentation of yield. The rice strong culm trait is influenced by the intricate interplay of many factors. To improve the complicated trait like strong culm in rice, it is crucial to understand how attributing traits interact with one another and helps to increase the grain yield. One of the requirements for directing selection is to know the degree of relation between strong culm and its contributing traits through correlation studies" [18]. "Since strong culm is a challenging complex trait that depends on several variables and the environment, to determine and reduce the number of characteristics for efficient selection, a method known as Principal Component Analysis

(PCA) can be effective. Principal component analysis (PCA), a multivariate technique, reduces data with a large number of correlated variables into a substantially smaller set of new variables through a linear combination of the variables that accounts for most of the variation present in the original variables" [19]. By keeping the aforesaid points in mind, the present investigation was undertaken to elucidate information on variability, heritability, and genetic advance in RILs derived from inter sub-specific cross to identify effective selection criteria for strong culm and yield attributes.

## 2. MATERIALS AND METHODS

A set of 353 RILs and six checks were evaluated during dry season (DS), 2022-23 at ICAR-IIRR, Hyderabad. The RILs were sown in the raised bed and transplanted into the main field 27 days after sowing with the spacing of 20 x 15 cm in Augmented randomized complete block design (ARCB). RILs were planted only once while the checks were replicated in all the seven blocks and a single seedling per hill was maintained. All the package of practices were followed for the good establishment of the crop. RILs were evaluated and observations were recorded on random four plants in each line for morphological traits viz., plant height (PH) (cm), internode length (IL) of fourth internode (cm), culm length (CL) (cm), outer diameter of major axis (ODMa)-b<sub>1</sub> (mm), outer diameter of minor axis (ODMi)-a<sub>1</sub> (mm), inner diameter of major axis (IDMa)-b<sub>2</sub> (mm), inner diameter of minor axis (IDMi)-a<sub>2</sub> (mm), culm diameter (CD) (mm), culm thickness (CT) (mm), section modulus (SM) (mm<sup>3</sup>), bending stress (BS), bending moment at breaking (M), Internode breaking weight (IBW) (g), pushing resistance (PR), culm with leaf sheath weight (CLSW)(g), tiller number (TN), panicle number (PN), panicle length (PL) (cm), panicle weight (PW) (g), Days to 50 percent flowering(DFF), total number of grains/panicle(GN), 1000 grain weight(TW) (g), grain yield (GY) (g). The genotypic coefficient of variation (GCV) and phenotypic coefficient of variation (PCV) were estimated according to the method Burton [17] and the estimates for variability were treated as per the categorization proposed by Sivasubramanian and Madhavamenon, [20]. The broad sense heritability (h<sup>2</sup>) and GAM were calculated as per Lush [21] and Johnson et al. [22]. The degree and direction of association between variables was estimated by using Karl Pearson's Coefficient of correlation (r) estimated from

variance and covariance of the variables and PCA was calculate using R ver 4.2.1 software.

The following traits were determined as per Ookawa et al. [23] and Badri et al. [15]

Culm diameter (CD) =  $(a_1+b_1)/2$   
 Culm thickness (CT) =  $[(a_1+b_1)/2] - [(a_2+b_2)/2]$   
 Section modulus (SM) =  $\pi/32 \times (a_1^3b_1-a_2^3b_2)/a_1$   
 Bending stress (BS) =  $(TR \div 40) \times (1000 \div TN)$   
 Bending moment at breaking (M) = Section modulus (SM) x Bending stress (BS)

Where; TR is the prostrate tester reading value (measure of pushing resistance) and TN is the tiller number. The prostrate tester (DIK-7400, Daiki Rika Kogyo Co. Ltd., Tokyo, Japan) was used to measure the pushing resistance of the culm [15].

### 3. RESULTS AND DISCUSSION

The identification of the key traits that can be used in crop improvement through various breeding techniques is extremely important. The detailed knowledge of variability, heritability, magnitude and direction of relationship between strong culm and its attributes along with PCA forms the foundation for effective selection of the traits in crop plants. In the present study, the tropical japonica line IRGC 39111, possessing a strong culm was used as a parent to develop RILs. The RILs elicited great phenotypic variability, particularly related to NPT phenology.

The Analysis of variance (ANOVA) displayed that the lines used in the study were significantly different for all the characters (Table 1). The variability analysis revealed that the PCV was

higher than GCV for all the traits in the study (Table 2) indicating the apparent variation is due to the environmental influence apart from genotypes. This result was in accordance with the findings of Akshay et al. [1]. The PCV and GCV were higher in the following traits viz., PR (27.5, 23.29), CT (42.45, 38.76); SM (56.22, 53.04); BS (35.58, 33.17), M (68.67, 67.89), IBW (32.71, 28.45) CLSW (28.82, 25.86), TN (31.08, 30.73), PN (31.03, 26.59), GN (21.73, 20.7), TW (26.36, 24.96) and similar findings were also reported by Reddy et al. [24], Bagudam et al. [5]. The observed variability among RILs probably be attributed to the inherent genetic differences and the environment in which they were grown. The findings are presented in Fig. 1.

The heritability for all the characters studied was high (>70%) except for ODMa, ODMi, GY which were medium and it ranged from 49.93 % to 97.74 % (Table 3). Heritability coupled with GAM was found to be more useful for selection (Johnson et al. 1955). In the present study, GAM ranged from 10.06 (GY) – 138.45 (M), which is a measure of culm strength. Most of the CS traits including PH, IL, CL, IDMa, IDMi, PR, BS, M, IBW and CLSW recorded high GAM and high heritability indicating that most likely the heritability is due to additive gene action and these traits were less influenced by the environment. Thus, can be easily be selected based on phenotype and can be effective. These results confirm that there is a possibility of direct selection for these traits, similar results were observed by Reddy et al. [24]. DFF and PL recorded high heritability (66.90 %) and moderate GAM (17.7) indicating the role of both additive and non-additive gene action (Fig. 2).




Fig. 1. Representation of Variability parameters for strong culm and yield components studied

**Table 1. Analysis of variance for strong culm and yield components in recombinant inbred lines of rice**

Source	Df	PH	IL	CL	ODMa	ODMi	IDMa	IDMi	TN	PR	CD	CT	SM
<b>Block (ignoring Treatments)</b>	6	1080.62**	34.58**	1055.09**	1.9**	2.23**	2.85**	3.15**	22.72*	104.96**	2.23**	0.65**	71.57**
<b>Treatment (eliminating Blocks)</b>	358	325.48**	28.42**	370.64**	1.19**	0.94**	0.78**	0.6**	14.02*	27.11**	0.98**	0.48**	57.19**
<b>Treatment: Check</b>	5	2383.28**	84.21**	982.8**	23.92**	21.7**	16.02**	8.54**	182.71**	340.5**	20.86**	3.49**	745**
<b>Treatment: Test and Test vs. Check</b>	353	296.33**	27.63**	361.97**	0.87**	0.65**	0.56**	0.49**	11.63*	22.68**	0.7**	0.44**	47.44**
<b>Residuals</b>	30	42.28	0.67	33.75	0.35	0.28	0.21	0.16	0.22	6.88	0.24	0.07	5.34
		<b>BS</b>	<b>M</b>	<b>IBW</b>	<b>CLSW</b>	<b>DFF</b>	<b>PN</b>	<b>PL</b>	<b>GN</b>	<b>PW</b>	<b>TW</b>	<b>GY</b>	
		456**	194770**	241952**	63.43**	289**	16.05*	35.1**	10370**	4.12**	110**	25.88**	
		341**	187328**	75993**	10.07**	76.19**	11.04*	7.77**	2126**	0.97**	33.34**	6.61**	
		3157**	1482436**	1379709**	104**	1762**	135**	46.11**	15518**	21.6**	127.44**	40.79**	
		301**	168984**	57526**	8.73**	52.3**	9.27*	7.23**	1936**	0.68**	32**	6.13*	
		37.69	3901.87	14743	1.76	4.91	1.93	2.19	193.67	0.1	3.37	3.09	

\*  $P \leq 0.05$ ; \*\*  $P \leq 0.01$

**Table 2. Estimates of variability, heritability and genetic advance for strong culm and yield contributing traits**

Trait	PV	GV	PCV	GCV	h(BS)	GAM
PH	256.71	214.43	12.91	11.8	83.53	22.25
IL	25.54	24.87	36.39	35.91	97.38	73.11
CL	235.49	201.74	15.5	14.35	85.67	27.4
ODMa	0.85	0.51	14.67	11.29	59.25	17.93
ODMi	0.66	0.39	16	12.19	57.99	19.14
IDMa	0.58	0.37	16.35	13.14	64.58	21.79
IDMi	0.5	0.34	19.77	16.31	68.04	27.75
TN	9.74	9.52	31.08	30.73	97.74	62.54
PR	24.36	17.48	27.5	23.29	71.74	40.7
CD	0.69	0.45	14.62	11.77	64.83	19.55
CT	0.45	0.37	42.45	38.76	83.36	73
SM	48.65	43.3	56.22	53.04	89.02	103.25
BS	288.82	251.14	35.58	33.17	86.95	63.82
M	171202.8	167300.9	68.67	67.89	97.72	138.45
IBW	60529.84	45785.87	32.71	28.45	75.64	51.04
CLSW	9.06	7.3	28.82	25.86	80.56	47.89
DFF	50.47	45.56	5.52	5.25	90.27	10.28
PN	7.26	5.33	31.03	26.59	73.41	51.26
PL	7.6	5.41	11.21	9.46	71.21	16.47
GN	2096.75	1903.08	21.73	20.7	90.76	40.68
PW	0.73	0.63	19.52	18.18	86.72	34.92
TW	32.62	29.25	26.36	24.96	89.68	48.76
GY	6.17	3.08	9.77	6.9	49.93	10.06

PV: Phenotypic variance; GV: Genotypic variance; PCV: Phenotypic coefficient of variance; GCV: Genotypic coefficient of variance; h(BS): Heritability in broad sense; GAM: Genetic advance as percent of mean, PH-Plant height, IL-Internode length, CL-Culm length, ODMa-Outer diameter of major axis, ODMi-Outer diameter of minor axis, IDMa- Inner diameter of Major axis, IDMi- Inner diameter of minor axis, TN-Tiller number, PR-Pushing resistance, CD- Culm diameter, CT-Culm thickness, SM- Section modulus, BS- Bending stress, M-Bending moment at breaking, IBW- Internode breaking weight, CLSW- Culm with leaf sheath weight, DFF-Days to 50% flowering, PN-Panicle number, PL- Panicle length, GN-No of filled grains per panicle, PW- Panicle weight, TW-Test weight, GY- grain yield.

**Table 3. Eigen value, contribution of variability for PCA in RILs**

Components	PC1	PC2	PC3	PC4	PC5	PC6	PC7
Eigen value	2.71	1.66	1.44	1.35	1.17	1.14	1.02
Percentage of variance	53.6	17.3	7.6	5.5	4.1	2.4	2.4
Cumulative percentage of variance	53.6	70.9	78.5	84	88.1	90.5	92.9
PH	0.17	0.38	0.23	-0.04	0.32	-0.21	0.03
IL	0.11	0.24	0.17	0.04	0.04	-0.21	0.00
CL	0.15	0.40	0.19	-0.01	0.31	-0.21	-0.03
ODMa	0.33	-0.17	-0.03	-0.05	0.12	0.06	-0.06
ODMi	0.33	-0.17	0.00	-0.02	0.10	0.08	0.01
IDMa	0.28	0.02	0.21	0.13	0.02	0.34	0.00
IDMi	0.24	0.03	0.19	0.18	0.04	0.47	0.00
TN	-0.11	-0.32	0.51	-0.12	-0.05	-0.07	0.02
TR	0.20	-0.01	0.20	0.13	-0.41	-0.07	0.37
CD	0.34	-0.17	-0.02	-0.05	0.12	0.07	-0.04
CT	0.20	-0.24	-0.25	-0.25	0.12	-0.40	-0.08
SM	0.30	-0.27	-0.06	-0.15	0.07	-0.17	-0.07
BS	0.20	0.24	-0.22	0.24	-0.36	-0.07	0.24
M	0.31	-0.11	-0.16	0.01	-0.16	-0.20	0.02
IBW	0.28	-0.05	0.05	0.01	-0.14	0.06	0.01
CLSW	0.18	0.11	0.10	0.04	-0.09	0.14	-0.52
DFF	0.02	0.15	0.12	0.02	-0.39	-0.18	-0.63
PN	-0.14	-0.33	0.46	-0.12	-0.02	-0.04	0.06
PL	0.07	0.19	0.17	-0.19	0.20	-0.04	0.08
GN	0.00	-0.14	0.16	0.51	0.03	-0.34	0.01
PW	0.07	0.13	0.24	-0.27	-0.37	-0.21	0.09
TW	0.05	0.19	-0.03	-0.61	-0.20	0.20	0.14
GY	0.10	0.04	0.04	0.07	0.06	-0.09	0.28

PH-Plant height, IL-Internode length, CL-Culm length, ODMa-Outer diameter of major axis, ODMi-Outer diameter of minor axis, IDMa-Inner diameter of Major axis, IDMi- Inner diameter of minor axis, TN-Tiller number, PR-Pushing resistance, CD- Culm diameter, CT-Culm thickness, SM- Section modulus, BS-Bending stress, M-Bending moment at breaking, IBW- Internode breaking weight, CLSW- Culm with leaf sheath weight, DFF-Days to 50% flowering, PN-Panicle number, PL-Panicle length, GN-No of filled grains per panicle, PW- Panicle weight, TW-Test weight, GY-grain yield

Pearson's correlation is the measure of strength of linear relationship between the characters [24]. The correlation analysis disclosed that the IBW and M had a high significant positive correlation with all the culm related traits, PW and GY (Fig. 3). Correlation studies allow a complete measure of the relationship between two traits. The positive correlation of strong culm with various traits was in concomitant with the reports of Ookawa et al. [23]; Zhang et al. [25]; Yang et al. [26]; Zhang et al. [27] and Badri et al. [15]. Such strong positive correlations among traits, heritability and genetic advance suggest that they are heritable and genetically controlled which could be simultaneously transferred to the desirable lines. The panicle length was negatively correlated with grain number. This may be due to longer panicle length but with a smaller number of filled grains. The distribution of traits was approximately continuous and indicated that the traits were controlled by a few minor genes and are typical quantitative traits.

Analysis of association among the strong culm attributing traits revealed that the IBW had the positive and highly significant correlation with panicle weight and grain yield. Hence selecting the plants with a higher IBW strongly improves the grain yield *via* panicle weight. Similarly, all the culm related traits were significantly and positively correlated with each other indicating that the selection may be positive direction for these traits. The above findings showed that strong selection for any one of the aforementioned attributes concurrently enhances the culm strength and grain yield. Panicle number per plant had a negative and significant correlation with the IBW, CD and CT. This clearly shows that increasing panicle number eventually reduces the strength of culm *via* partitioning of biomass to produce greater number of panicles by the plant. A critical analysis of the character association results revealed a favorable correlation between the grain yield per plant and IBW, CT, CD, CL [28]. As a result, choosing for

these characteristics may improve the yield components and culm strength.

Principal component (PC) analysis was carried out to identify the minimum number of components which can explain maximum variability out of total variability and also rank genotypes on the basis of PC scores Reddy et al. [24]. Among the 23 PCs, seven PCs had eigen values greater than 1.00 and accounted for

approximately 92.9% of the total cumulative variability. The scree plot (Fig. 4) showed the amount of variance described by each PC and showed that PC1 had the largest eigen value contributing highest amount of variability, at about 53.6%. Additionally, PC2, PC3, PC4, PC5, PC6, and PC7 showed a continuous decrease in variability, with comparable values of 17.3%, 7.6%, 5.5%, 4.1%, 2.4% and 2.4% respectively.




Fig. 2. Representation of Heritability and genetic advance as per cent of mean for strong culm and yield components





Fig. 3. Correlation matrix with significant values and trait distribution among 23 characters



**Fig. 4. Representation of Biplot diagram of PC1 and PC2 and Scree plot explaining variance components**

The first two primary PCs' biplot (Fig. 4) demonstrated how the features affected the divergence. The length of a trait's vector in PCA represents its contribution to the overall divergence; the longer the vector, the larger the contribution. All culm and yield related traits except for DFF, GN and GY exhibited maximum length and contributed maximum to the total diversity. These results were in conformity with

the findings of Reddy et al. [24], Gupte et al. [29], Chigira et al. [30], Tiwari et al. [31].

#### 4. CONCLUSION

In summary, high PCV and GCV, together with heritability combined with GAM, were reported by the traits PR, IL, CT, SM, BS, M, IBW, CLSW, TN, PN, GN, and TW. Therefore, while selecting

genotypes to increase culm strength, these attributes should be considered. Correlation analysis revealed a positive and substantial relationship between the GY and PW and the culm strength (IBW, M). In the PCA analysis, SM, ODMi, M, IBW, and CT had the longest vector lengths and contributed the most to overall diversity and the lines with high mean values for these traits might be utilized as donors to increase the culm strength and its attributing traits in a future breeding programs. In this study, we present the synergistic associations between culm strength and grain yield traits and tropical *japonica* germplasm can be a great source of genetic diversification for rice improvement. The variability and character associations identified in the present study demonstrated that introgression of desirable traits particularly thick culm with greater physical strength associated with grain yield traits from tropical *japonica* would broaden the genetic base of elite *indica* cultivars with enhanced yield potential. Our ongoing experiments aim to utilize the findings to decipher the complex intricated interactions involved in strong culm improvement and provide viable solution to lodging problem in rice.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

#### ACKNOWLEDGEMENT

The first author sincerely recognizes the guidance provided by Dr. Jyothi Badri, Senior scientist, IIRR and Dr. L. Krishna, Principal scientist, IRR, PJTSAU for providing all of the necessary facilities for our investigation.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Akshay M, Chandra BS, Devi KR, Hari Y. Genetic variability studies for yield and its attributes, quality and nutritional traits in rice (*Oryza sativa* L.). The Pharma Innovation Journal. 2022;11(5):167-172.
2. Eizenga, G.C., McClung, A.M. and Huggins, T.D. 2021. Registration of two *Oryza sativa* tropical japonica germplasm lines selected for panicle architecture and grain size traits. Journal of Plant Registrations. 15(3). 573-582.
3. Singh VJ, Bhowmick PK, Vinod KK, Krishnan SG, Nandakumar S, Kumar A, Kumar M, Shekhawat S, Dixit BK, Malik A, Ellur RK. Population structure of a worldwide collection of tropical japonica rice indicates limited geographic differentiation and shows promising genetic variability associated with new plant type. Genes. 2022;13(3):484.
4. Fujita D, Trijatmiko KR, Tagle AG, Sapasap MV, Koide Y, Sasaki K, Tsakirpaloglou N, Gannaban RB, Nishimura T, Yanagihara S, Fukuta Y. NAL1 allele from a rice landrace greatly increases yield in modern indica cultivars. Proceedings of the National Academy of Sciences. 2013;110(51):20431-20436.
5. Bagudam R, Eswari KB, Badri J, Devi GL, Jai Vidhya LRK, Bhavani P, Swathi G, Bharath M, Rao PR, Kumar JA, Fiyaz RA. Morphological and molecular characterization of new plant type core set for yield and culm strength traits in rice (*Oryza sativa* L.). Journal of Plant Biochemistry and Biotechnology. 2021;30: 233-242.
6. Ashikari M, Sakakibara H, Lin S, Yamamoto T, Takashi T, Nishimura A, Angeles ER, Qian Q, Kitano H, Matsuoka M. Cytokinin oxidase regulates rice grain reduction. Science. 2005;309(5735):741-745.
7. Ikeda-Kawakatsu K, Yasuno N, Oikawa T, Iida S, Nagato Y, Maekawa M, Kyozuka J. Expression level of ABERRANT PANICLE Organization determines rice inflorescence form through control of cell proliferation in the meristem. Plant physiology. 2009;150(2):736-747.
8. Terao T, Nagata K, Morino K, Hirose T. A gene controlling the number of primary rachis branches also controls the vascular bundle formation and hence is responsible to increase the harvest index and grain yield in rice. Theoretical and Applied Genetics. 2010;120:875-893.
9. Huang X, Qian Q, Liu Z, Sun H, He S, Luo D, Xia G, Chu C, Li J, Fu X. Natural variation at the DEP1 locus enhances grain yield in rice. Nature genetics. 2009;41(4):494-497.
10. Miura K, Ikeda M, Matsubara A, Song XJ, Ito M, Asano K, Matsuoka M, Kitano H,

- Ashikari M. OsSPL14 promotes panicle branching and higher grain productivity in rice. *Nature genetics*. 2010;42(6):545-549.
11. Fan C, Xing Y, Mao H, Lu T, Han B, Xu C, Li X, Zhang Q. GS3, a major QTL for grain length and weight and minor QTL for grain width and thickness in rice, encodes a putative transmembrane protein. *Theoretical and applied genetics*. 2006; 112:1164-1171.
  12. Song XJ, Huang W, Shi M, Zhu MZ, Lin HX. A QTL for rice grain width and weight encodes a previously unknown RING-type E3 ubiquitin ligase. *Nature genetics*. 2007;39(5):623-630.
  13. Shomura A, Izawa T, Ebana K, Ebitani T, Kanegae H, Konishi S, Yano M. Deletion in a gene associated with grain size increased yields during rice domestication. *Nature genetics*. 2008;40(8):1023-1028.
  14. Hirano K, Ordonio RL, Matsuoka M. Engineering the lodging resistance mechanism of post-Green Revolution rice to meet future demands. *Proceedings of the Japan Academy, Series B*. 2017;93(4):220-233.
  15. Badri J, Padmashree R, Anilkumar C, Mamidi A, Isetty SR, Swamy AVSR, Sundaram RM. Genome-wide association studies for a comprehensive understanding of the genetic architecture of culm strength and yield traits in rice. *Front. Plant Sci*. 2024;14:1298083.
  16. Khan MK, Pandey A, Hamurcu M, Hakki EE, Gezgin S. Role of molecular approaches in improving genetic variability of micronutrients and their utilization in breeding programs. In *Wheat and barley grain biofortification*. 2020;27-52.
  17. Burton GW. Quantitative inheritance in grasses. In *proceedings of 6<sup>th</sup> International Grassland Congress, Pennsylvania State College, USA*. 1952;277-283.
  18. Singh R, Yadav V, Mishra DN, Yadav A. Correlation and path analysis studies in rice (*Oryza sativa* L.). *Journal of Pharmacognosy and Phytochemistry*. 2018;7:2084-2090.
  19. Varanasi YVP, Isetty SR, Revadi P, Balakrishnan D, Hajira S, Prasad MS, Laha GS, Perraju P, Singh UM, Singh VK, Kumar A, Sundaram RM, Badri J. Molecular and Morphological Characterization of Introgression Lines with Resistance to Bacterial Leaf Blight and Blast in Rice. *Plants*. 2023;12(16): 3012.
  20. Sivasubramanian J, Madhavamenon. Genotypic and phenotypic variability in rice. *Madras Agric J*. 1973;12:15-16.
  21. Lush JL. Intra - sire correlation and regression of off spring on dams as a method of estimating heritability of characters. *Proc Am Soc Anim Prod*. 1940;33:293-301.
  22. Johnson HW, Robinson HF, Comstock RE. Estimates of genetic and environmental variability in soybean. *Agronomy Journal*. 1955;47:314-318
  23. Ookawa T, Hobo T, Yano M, Murata K, Ando T, Miura H, Asano K, Ochiai Y, Ikeda M, Nishitani, R, Ebitani T. New approach for rice improvement using a pleiotropic QTL gene for lodging resistance and yield. *Nature communications*. 2010;1(1): 132.
  24. Reddy VRP, Vemireddy LR, Srividhya A., Reddy KH, Siddiq EA. Genetic analysis of culm strength and its related traits in rice (*Oryza sativa* L.). *Applied biological research*. 2019;21(2).
  25. Zhang F, Jin Z, Ma G, Shang W, Liu H, Xu M. Relationship between lodging resistance and chemical contents in culms and sheaths of Japonica rice during grain filling. *Rice Science*. 2010;17:311-318.
  26. Yang H, Fang X, He H, Xie Z. Relationship of characteristics of culm construction to lodging resistance and yield of Fujian-bred super-rice cultivars. *Chinese Journal of Eco Agriculture*. 2012;20:909-913.
  27. Zhang W, Wu L, Wu X, Ding Y, Li G, Li J, Weng F, Liu Z, Tang S, Ding C, Wang S. Lodging resistance of Japonica rice (*Oryza sativa* L.). Morphological and anatomical traits due to top dressing nitrogen application rates. *Rice*. 2016;9:31.
  28. Yadav S, Singh UM, Naik SM, Venkateshwarlu C, Ramayya PJ, Raman KA, Sandhu N, Kumar A. Molecular mapping of QTLs associated with lodging resistance in dry directed-seeded rice (*Oryza sativa* L.). *Frontiers in Plant Science*. 2017;8:1431.
  29. Gupte VP, Manonmani S, Nivedha R, Suresh R, Kumar GS, Raveendran M. Genetic diversity studies and identification of donors for lodging resistance in rice (*Oryza sativa* L.). *Electronic Journal of Plant Breeding*. 2023;14(3):1158-1166.
  30. Chigira K, Yamasaki M, Adachi S, Nagano AJ, Ookawa T. Identification of novel quantitative trait loci for culm thickness of

- rice derived from strong-culm landrace in Japan, Omachi. Rice. 2023;16(1).
31. Tiwari DN, Tripathi SR, Tripathi MP, Khatri N, Bastola BR. Genetic variability and correlation coefficients of major traits in early maturing rice under rainfed lowland environments of Nepal. Adv Agric. 2019;1:1-9.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/119549>