



Improving Peanut (*Arachis hypogaea* L.) Productivity by Optimizing Nitrogen and Potassium Fertilization

E D Purbajanti ^{a*}, D Safitri ^a, A.F.Irawan ^a and S.Arafat ^a

^a Study Program of Agro-ecotechnology, Faculty of Animal and Agricultural Sciences, Diponegoro University, Indonesia.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ijpss/2024/v36i105102>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/125324>

Original Research Article

Received: 24/08/2024
Accepted: 27/10/2024
Published: 31/10/2024

ABSTRACT

Due to their high nutritious, peanuts (*Arachis hypogaea* L.) are a popular kind among consumers. For growth and productivity, peanut plants need vital nutrients including N, P, K, Ca, Mg and K. P is especially necessary for the development of blooms, pods, and seeds. A 2 × 4 factorial design was employed in the study, and it was repeated three times. Factor II consists of P fertilizers (SP36: 36 % P₂O₅), specifically 0, 50, 100, and 150 kg P₂O₅/ha. Factor I sources of N are NO₃⁻ (potassium nitrate) and NH₄⁺ (ammonium sulfate). The criteria that were observed were the amount of chlorophyll in the fifth leaf from the tip of the plant, NR activity, yield of the peanut pod, height of the peanut plant, and biomass of the peanut. The nitrogen source treatment has a major impact on ANR levels, yield, plant height, and chlorophyll. There is a greater interaction when using nitrate as

*Corresponding author: E-mail: endang.purbajanti@live.undip.ac.id;

a N fertilizer. Higher P₂O₅ NR yields, P up to 150 kg, Chlorophyll, height of peanut plants, peanut biomass, and yield of peanut pods are observed. Similarly, increased application of N fertilizer (ammonium, P up to 150 kg P₂O₅) results in higher NR yields. Height of the peanut plants, biomass, yield of the peanut pods, and chlorophyll.

Keywords: Peanut; nitrate; ammonium; phosphorus; peanut pod yield.

1. INTRODUCTION

Plants require phosphate minerals, which are macronutrients, in significant amounts for growth. Due to its direct involvement in nearly every biological function, phosphate is frequently referred to as the "key to life." Every living cell contains it, however seeds and growth points typically contain more of it [1]. The availability of nutrients, such as the element phosphate (P), influences peanut development and yield in addition to the physical state of the soil. Plants benefit greatly from the element phosphate, particularly in the areas of energy transfer, protein synthesis, coenzymes, metabolic acids, nucleic acids, and compounds [2].

The issue that frequently arises is the phosphate content of the soil. The availability of phosphate in Indonesian agricultural soil is poor, despite the soil's nearly whole high phosphate content. Due to their high nutritious content, peanuts (*Arachis hypogaea* L.) are a popular kind among consumers. According to Nurmi et al. [3], peanuts have the following nutritional value per 100 grams: 25 grams of protein, 21 grams of carbs, 48 grams of fat, 336 mg of phosphorus, 332 mg of potassium, 184 mg of magnesium, 62 mg of calcium, and trace amounts of iron, sodium, zinc, and vitamins B, E, and K. Peanuts with seed type 3 are those of the Katana 2 variety. The benefits of this variety include resistance to bacterial wilt disease, early maturity of 87 DAP, potential yield of 4.7 tons, and good paddy field adaptation [4]. Plants that produce peanuts For growth and production, soil needs vital minerals like N, P, and K, especially P for the development of flowers, pods, and seeds [5]. Peanut plants with a P nutritional deficit have stunted growth, pale green leaves, and low pod yields [6].

Both internal and exterior influences might have an impact on a plant's growth and development. Water, light, soil, and humidity are examples of external variables. Genes, anatomical structure, hormones, plant shape, and quantities of chlorophyll and other pigments are examples of internal influences. An essential metric for

assessing plant growth, photosynthetic capacity, and general health is chlorophyll. Purbajanti et al. [7] state that because leaf chlorophyll levels are correlated with plant conditions, it is possible to utilize them to estimate the amount of additional fertilizer that plants require. One technique to gauge plant growth and fertility, which may then be connected to forecasts of plant yield, is to measure the amount of chlorophyll. Phosphoric nutrients can be applied to peanut seeds to promote their growth and development [8]. Peanuts grow to lower heights, with little, pale green leaves and tiny pods, and produce very low yields due to phosphorus deficiency [9]. Phosphate increases the amount of fat in seeds and is essential for the process of cell formation. The weight of one hundred seeds demonstrates how the variety and capacity of plants for photosynthesis affect the quality of seeds [10]. In agronomy, the idea of essential N concentration is frequently utilized to diagnose N plant health and distinguish between circumstances involving suboptimal and supra optimal N supply. Several physiological processes in plants are regulated, which affects the link between N and biomass accumulation in plants. N intake, plant C assimilation and consequent growth rates, and the distribution of C and N among organs and plants are some of these activities. Plant uptake of N under Suboptimal Supply N is dependent on the distribution and availability of N-mineral soil as well as the distribution of roots. When there is an adequate supply of nitrogen, a plant's ability to absorb nitrogen is mostly determined by its internal regulation of growth [11]. In plants with low root elongation rates, mature leaves receive about 70% of the N absorbed; in plants with high root elongation rates, this percentage is just 35%. It was evident that N absorption and allocation affected the activity of root growth. Plants with low levels of root elongation allotted a modest quantity of N to their adult leaves, but plants with high levels of root elongation allocated much more N to their adult leaves. Given that low-level elongation roots may be adult leaves and are supplied by mobilization from other parts, it is now probable that N is required to promote adult leaf growth [12]. According to Shi-cheng et al. [13], most plants

grow similarly in both N sources at the low end of the N scale, or provisions. Nevertheless, among plants that have evolved in NH_4 -rich environments, there are a few noteworthy outliers to this general pattern. The purpose of this study was to determine how fertilizer combinations including N and P affected the growth and yield of peanuts.

2. MATERIALS AND METHODS

The study was conducted in the laboratory of Ecology and Plant Production, Faculty of Agriculture and Animal Husbandry, Diponegoro University, during the dry season (August to December) in 2022. The research station is located at $6^\circ 58' 0''$ SL $110^\circ 25' 0''$ EL, and has Oxisol soil with a clay texture, a soil pH of 6.5, 0.18% total N content, 0.15% total P, 0.24% total K, and 1.09% C. Katana2 peanut varieties were used, and the study was carried out using 24 experimental plots measuring $1.5 \times 1 \text{ m}^2$, with a spacing of $30 \times 20 \text{ cm}$ so that each plot contained 16 plants. The research used a 2×4 factorial design that was repeated three times.

Factor I: sources of N (NO_3^- (potassium nitrate) and NH_4^+ (ammonium sulfate))

Factor II: P fertilizers (SP36: 36 % P_2O_5), namely 0, 50, 100, and 150 kg $\text{P}_2\text{O}_5/\text{ha}$.

Each plant was fertilized with manure at 2 tons per hectare and K at 25 kg KCl per hectare. Micronutrients were administered in the form of approximately 5 ml of liquid fertilizer dissolved in 1 L of water and sprayed onto the plants 1 wk after planting. Fertilize P (SP 36) according to the dose, while N fertilizer required is 50 kg N per hectare. The chlorophyll content of the fifth leaf from the tip of the plant, NR activity, peanut pod yield, and peanut plant height and peanut biomass were determined. Chlorophyll content was measured in fresh leaf samples. Leaf samples (0.5 g) were homogenized with acetone (90% v/v), filtered, and made up to a final volume of 50 mL. Chlorophyll concentration was calculated based on the absorbance of the extract measured by a spectrophotometer 645 UV/Vis and 666 UV/Vis [14]. Analysis of NR activity was performed according to the methods used by Krywult and Bielec [15]. The third leaves of peanut plant shoots were harvested between 9 and 10 a.m. and used for further observation. The leaves were washed with distilled water and finely sliced, and 200 mg of the leaf sample was added to 5 ml NaH_2PO_4 and NaHPO_4 buffer

solutions at pH 7.5 in dark tubes that were covered and maintained for 24 h, after which the buffer solution was replaced with 5 mL of fresh buffer solution. Next, 0.1 mL of 5 M NaNO_3 was added to every dark tube. The time of addition of NaNO_3 was designated the 0-incubation point, and samples were incubated for 2 h. Meanwhile, another test tube was filled with 0.2 mL 1% sulfanilamide reagent, which was dissolved in 3N HCl and 0.2 mL 0.02% naphthylethylenediamide. Then 0.1 mL of the filtrate incubated for 2 h was placed in a test tube containing the sulfanilamide reagent, HCl, and a naphthylethylenediamide solution. The test tubes were agitated to mix the filtrate and accelerate the reaction, and allowed to stand for 15 min; this resulted in the reduction of NO_2 with dye reagents to bring up the pink color. Next, 2.5 ml distilled water was added to the test tube as a color diluent. The absorbance of this solution was measured at a wavelength of 540 nm.

2.1 Data Analysis

The data were analyzed using ANOVA. If the ANOVA revealed a difference, the analysis was followed by Duncan's multiple range test 5 % as described by Gomez and Gomez [16].

3. RESULTS AND DISCUSSION

3.1 Results

3.1.1 The effect of N form (NO_3^- and NH_4^+)

NR decreased by 16.20% for the use of NH_4 compared to NO_3 . The chlorophyll content of peanut leaves with N form NH_4 decreased 19.26% compared to the NO_3 form. Likewise, plant height decreased by 9.07% when using NH_4 compared to NO_3 . The use of ammonium form of nitrogen resulted in biomass decreasing by 110.77% and pod weight decreasing by 23.67%.

3.1.2 The effect phosphor dosages

The NR of peanuts with a P dose of 100 kg P_2O_5 increased by 23.00% and P with a P dose of 150 kg P_2O_5 increased by 26.82% compared to a P dose of 50 kg P_2O_5 . The results of the chlorophyll content increased by 6% when using a P dose of 100 kg P_2O_5 , also increased by 43.87% when using P fertilizer of 150 kg P_2O_5 compared to a P dose of 50 kg P_2O_5 . This pattern also occurred in plant height, biomass yield and pod weight. Plant height increased by 8.18% because P fertilizer

increased from 50 kg P₂O₅ to 100 kg P₂O₅, while it increased by 25.83% because P fertilizer increased from 50 kg P₂O₅ to 100 kg P₂O₅. Peanut biomass yield also increased by 32.03% due to the use of a P dose of 100 kg P₂O₅, and increased by 79.65% due to the use of a P dose of 150 kg P₂O₅ compared to a P dose of 50 kg P₂O₅.

3.1.3 The effect of Interaction

If you use N fertilizer in the form of nitrate, the higher the P fertilizer, up to 150 kg P₂O₅, the NR yield will be higher, likewise, if you use N fertilizer in the form of ammonium, the higher the P fertilizer, up to 150 kg P₂O₅, the higher the NR yield (Fig. 1).

Use of P fertilizer up to 150 kg P₂O₅ and nitrate fertilizer together resulted in an increase in chlorophyll content. Meanwhile, the application of a mixture of P and ammonium fertilizers up to

150 kg P₂O₅ resulted in an increase in the chlorophyll content (Fig. 2). In terms of plant height metrics, the use of P 100 kg P₂O₅ in conjunction with nitrate fertilizer and P fertilizer caused peanut plant height to grow; this combination did not differ significantly from that of P 150 kg P₂O₅ and nitrate fertilizer. In comparison to the combinations of ammonium fertilizer and P 50 kg P₂O₅ and P 100 kg P₂O₅, the combination of ammonium fertilizer and P 150 kg P₂O₅ produced the plants with the highest height (Fig. 3).

The usage of nitrate and P fertilizer up to 100 kg P₂O₅, which is not substantially different from the combination of nitrate and P up to 150 kg P₂O₅, has led to an increase in biomass characteristics. In contrast to the combinations of ammonium and P fertilizer up to 100 kg P₂O₅ and manure and P fertilizer up to 50 kg P₂O₅, the combination of ammonium and P fertilizer up to 150 kg P₂O₅ has the highest biomass output (Fig. 4).




Fig. 1. NR activity peanut plant




Fig. 2. Chlorophyll content in peanut plants due to interactions between nitrogen sources and Phosphorus fertilizer




Fig. 3. Peanut plant height due to interactions between nitrogen sources and Phosphorus fertilizer




Fig. 4. Weight of peanut plant biomass due to interactions between nitrogen sources and Phosphorus fertilizer




Fig. 5. Peanut pod yield due to interactions between nitrogen sources and Phosphorus fertilizer

In comparison to the combinations of nitrate fertilizer and P fertilizer up to 50 kg P₂O₅ and nitrate fertilizer and P fertilizer up to 100 kg P₂O₅, the combination of nitrate fertilizer and P fertilizer up to 150 kg P₂O₅ produced the highest pod weight result. When ammonium and P fertilizer are combined, the pod weight yield is often lower than when nitrate and P fertilizer are combined (Fig. 5).

4. DISCUSSION

Sampara et al. [17] reported that the availability of nutrients encourages the division of meristem cells in plants, which in turn leads to an increase in height. This suggests that the tendency for plant height to increase is influenced by these nutrients. Nitrogen, a macronutrient necessary for the synthesis of the amino acids that make up proteins, is necessary for plant growth. Furthermore, a number of genes' expression is regulated by the presence of nitrogen [18]. The primary enzyme in tissues that actively performs photosynthesis is called nitrate reductase (NR) [19]. This enzyme plays a crucial part in the initial steps of amino acid synthesis. Yanagisawa [20] states that nitrates in the leaves stimulate the activity of NR biosynthesis, which is dependent on the availability of nitrogen nutrients in the medium. Chlorophyll levels are another factor linked to nitrogen metabolism. NR will convert nitrate, a source of nitrogen, to ammonium (NH₄). The synthesis of glutamine from glutamate requires NH₄; the presence of 2-oxoglutarat increases glutamate, which is essential for the synthesis of several significant amino acids that are components of proteins, including chlorophyll [19]. Plant photosynthetic rate will increase with full solar intensity and large fertilizer dosages [7]. Nitrate reductase, which is utilized in the respiration process, is dependent on the availability of carbohydrates from photosynthesis. Such an ANR pattern on leaf strands is produced by the combination of treatments applied for 45 days in this study. This is due to regulation of activation of NR in the cytosol, among others, determined by numerous parameters, including light, CO₂ levels and availability of NADH / NAD [21]. The availability of carbon frames and the rates of photosynthesis are highly correlated with NR activity. Light can boost NR activity, and NO₃-inducing NR gene transcription occurs [22].

Chlorophyll reveals that peanut plants have higher nitrogen content, which is also used to make chlorophyll. Through the mechanisms of transcription and translation, nutrient N has a

variety of roles in the metabolic process, controlling the expression of multiple genes that produce proteins in high-level plants and algae while also preserving the stability of mRNA [23]. PEP-Case, nitrate reductase (NR), nitrite reductase (NiR), light harvesting complex protein (LHCP), proteins regulating vegetative growth, and chlorophyll a/b are among the proteins in discussion.

With the formulas $y = 0.9135x - 7.7415$ ($R^2 = 0.975$) for the nitrate form of N source and $y = 1.8323x - 91.568$ ($R^2 = 0.4742$) for the ammonium form of N source, the activity of nitrate reductase and peanut pod yield have a positive association. The study's findings demonstrated that phosphate fertilizer application significantly impacted the biomass of peanut plants. the pod weight for each plant in each plot. When coupled with ammonium and nitrate, 150 kg of P₂O₅ phosphate fertilizer had the best outcomes when compared to other lesser doses. Every stage of a plant's life is significantly impacted by the application of phosphate fertilizer.

This is in line with the findings of Pasaribu et al. [24], which claim that phosphate is essential for multiple processes in the plant body, including: (1) cell division, the synthesis of fat and albumin; (2) the development of flowers, fruit, and seeds; (3) plant maturity, which mitigates the effects of nitrogen; (4) stimulation of root development; (5) improvement of crop yield quality; and (6) resistance to pests and diseases. This demonstrates that by enhancing the chemical characteristics of the soil, P fertilizer can affect the generative phase of peanut plants.

5. CONCLUSION

The nitrogen source treatment has a major impact on ANR levels, yield, plant height, and chlorophyll. The interaction between the use of pupuk N bentuk nitrate as low as 150 kg P₂O₅, the results of NR becoming low, chlorophyll, the height of the peanut plant, peanut biomass, and the yield of the peanut pod. Additionally, the use of pupuk N bentuk ammonium was found to be as high as 150 kg P₂O₅, as well as high levels of chlorophyll, peanut plant height, peanut biomass, and peanut pod yield.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models

(ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript

ACKNOWLEDGEMENTS

Thank you to the Dean of the Faculty of Animal Husbandry and Agriculture for funding Research from PNPB funds.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Khan F, Siddique AB, Shabala S, Zhou M, Zhao C. Phosphorus plays key roles in regulating plants' physiological responses to abiotic stresses. *Plants (Basel)*. 2023 Aug;12(15):2861. DOI: 10.3390/plants12152861
2. Bindraban PS, Dimkpa CO, Pandey R. Exploring phosphorus fertilizers and fertilization strategies for improved human and environmental health. *Biol Fertil Soils*. 2020;56:299–317. DOI: 10.1007/s00374-019-01430-2
3. Nurmi A, Azis S, Mooduto SB. Growth and yield of peanut plants (*Arachis hypogaea* L.) with the application of organic fertilizer from husks charcoal and chicken manure. *Jurnal Produksi Tanaman*. 2023;11(10):786–792. DOI: 10.21776/ub.protan.2023.011.10.07
4. BSIP. Pertanian. 2022. Available: <http://banten.bsip.pertanian.go.id>
5. Zhu L, Zhen-wu N, Song-ming L, Hai-qiu Y, Li-yong X, Wei-wei M, Zheng Z, Shu-bo W. Millet/peanut intercropping at a moderate N rate increases crop productivity and N use efficiency, as well as economic benefits, under rain-fed conditions. *J Integr Agric*. 2023;22(3):738–751. DOI: 10.1016/j.jia.2022.08.078
6. Faza DA, Lukiwati DR, Karno. Growth and production of peanut (*Arachis hypogaea* L.) by vesicular-arbuscular mycorrhizal fungi inoculation and phosphorus fertilization. *J Agro Complex*. 2019;3(1):48–54. DOI: 10.14710/joac.3.1.48-54
7. Purbajanti ED, Soetrisno RD, Hanudin E, Budhi SPS. Photosynthesis and yields of grasses grown in saline condition. *J Indonesian Trop Anim Agric*. 2010;35(1):[no page numbers provided].
8. Arista D, Suryono, Sudadi. Effect of N, P and K fertilizer combinations to growth and yield of peanut on Alfisols dry land. *Agrosains*. 2015;17(2):49-52.
9. Rahman N, Suntoro, Sekya AT. Peanut growth and gynophore formation on boron and phosphor applications. *Sains Tanah J Soil Sci Agroclimatol*. 2019;16(1):57-66. DOI: 10.20961/stjssa.v16i1.25372
10. Suntoro, Syamsiyah J, Rahina W. Availability and uptake of Ca in peanuts in Alfisols soils treated with Kelud volcanic ash and manure. *Agrosains*. 2017;19(2):51-57.
11. Xia G, Wang Y, Hu J, Wang S, Zhang Y, Wu Q, Chi D. Effects of supplemental irrigation on water and nitrogen use, yield, and kernel quality of peanut under nitrogen-supplied conditions. *Agric Water Manag*. 2021;243:106518. DOI: 10.1016/j.agwat.2020.106518
12. Han F, Guo R, Hussain S, Guo S, Cai T, Zhang P, Jia Z, Naseer MA, Saqib M, Chen X, Ren X. Rotation of planting strips and reduction in nitrogen fertilizer application can reduce nitrogen loss and optimize its balance in maize-peanut intercropping. *Eur J Agron*. 2023;143:126707. DOI: 10.1016/j.eja.2022.126707
13. Shi-cheng Z, Ji-long L, Xin-peng X, Xiaomao L, Rosso LM, Shao-jun I, Ciampitti I, Ping H. Peanut yield, nutrient uptake and nutrient requirements in different regions of China. *J Integr Agric*. 2021;20(9):2502–2511. DOI: 10.1016/S2095-3119(20)63253-1
14. Rajalakshmi K, Banu N. Extraction and estimation of chlorophyll from medicinal plants. *Int J Sci Res*. 2015;4(11):209-212.
15. Krywult W, Bielec D. Method of measurement of nitrate reductase activity in field conditions. *J Ecol Eng*. 2013;14(1):7-11.
16. Gomez KA, Gomez AA. Statistical procedures for agricultural research. 2nd ed. New York: John Wiley and Sons; 1984. 680 p.
17. Sampara S, Syamsia S, Idhan A, Rosanna, Mado I. Growth and production of peanuts (*Arachis hypogaea* L.) at various doses of organic potassium and liquid organic fertilizer. *J Agroplanta*. 2023;12(2):181–191. DOI: 10.51978/agro.v12i2.695

18. Wang XB, Liu WX, Li ZG, Teng Y, Christie P, Luo YM. Effects of long-term fertilizer applications on peanut yield and quality and plant and soil heavy metal accumulation. *Pedosphere*. 2020;30(4): 555–562.
DOI: 10.1016/S1002-0160(17)60457-0
19. Purbajanti ED, Slamet W, Fuskhah E. Nitrate reductase, chlorophyll content and antioxidant in okra (*Abelmoschus esculentus* Moench) under organic fertilizer. *J Appl Horticult*. 2019;21(3):213-217.
20. Yanagisawa S. Transcription factors involved in controlling the expression of nitrate reductase genes in higher plants. *Plant Sci*. 2014;229:167-171.
DOI: 10.1016/j.plantsci.2014.09.006
21. Broadley M, Brown P, Cakmak I, Rengel Z, Zhao F. Chapter 7 - Function of nutrients: Micronutrients. In: Marschner's mineral nutrition of higher plants. 3rd ed. 2012. p. 191-248.
22. Berger A, Boscari A, Araújo NH, Maucourt M, Hanchi M, Bernillon S, Rolin D, Puppo A, Brouquisse R. Plant nitrate reductases regulate nitric oxide production and nitrogen-fixing metabolism during the *Medicago truncatula*–*Sinorhizobium meliloti* symbiosis. *Front Plant Sci*. 2020; 11:1313.
DOI: 10.3389/fpls.2020.01313
23. Martins T, Barros AN, Rosa E, Antunes L. Enhancing health benefits through chlorophylls and chlorophyll-rich agro-food: A comprehensive review. *Molecules*. 2023 Jul;28(14):5344.
DOI: 10.3390/molecules28145344
24. Pasaribu PK, Barus A, Mariati. Growth and production of peanuts (*Arachis hypogaea* L.) with cow manure and phosphate fertilizer application. *Jurnal Online Agroekoteknologi*. 2014;2(4): 1391-1395.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/125324>