



# Impact of Phosphorus Levels and PSB Application on Soil Physio-Chemical Properties in Mustard Fields under Gmelina-Based Agroforestry

Deeksha Gupta <sup>a\*</sup>, R. Bajpai <sup>a</sup>, S. B. Agrawal <sup>a</sup>,  
B. S. Dwivedi <sup>b</sup>, Komal Bhatt <sup>a</sup>, Poornima Malviya <sup>c</sup>  
and Rahul Mandloi <sup>d</sup>

<sup>a</sup> Department of Forestry, College of Agriculture, JNKVV, Jabalpur-482004, (Madhya Pradesh.), India.

<sup>b</sup> Department of Soil Science, College of Agriculture, JNKVV, Jabalpur-482004, (Madhya Pradesh), India.

<sup>c</sup> Department of Agronomy, College of Agriculture, JNKVV, Jabalpur-482004, (Madhya Pradesh), India.

<sup>d</sup> Department of Post Harvest Process and Food Engineering, College of Agricultural Engineering, JNKVV, Jabalpur-482004, (Madhya Pradesh), India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/ijpss/2024/v36i125206>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/128319>

Original Research Article

Received: 07/10/2024

Accepted: 16/12/2024

Published: 19/12/2024

\*Corresponding author: E-mail: [guptadeeksha008@gmail.com](mailto:guptadeeksha008@gmail.com);

**Cite as:** Gupta, Deeksha, R. Bajpai, S. B. Agrawal, B. S. Dwivedi, Komal Bhatt, Poornima Malviya, and Rahul Mandloi. 2024. "Impact of Phosphorus Levels and PSB Application on Soil Physio-Chemical Properties in Mustard Fields under Gmelina-Based Agroforestry". *International Journal of Plant & Soil Science* 36 (12):327-35. <https://doi.org/10.9734/ijpss/2024/v36i125206>.

## ABSTRACT

This study examines how soil qualities are affected by phosphorus (P) levels and phosphate solubilising bacteria (PSB) inoculation techniques in an agroforestry system based on *Gmelina arborea*, with a particular emphasis on mustard (*Brassica juncea* L.) growing. The experiment was conducted from November to April in two consecutive years 2022-2023 and 2023-2024 at JNKVV in Jabalpur, used a split-plot design with PSB application techniques (soil, seedling, and combination) as sub-treatments and phosphorus levels (30, 40, and 50 kg/ha) as the primary treatments. For soil analyses soil sample was collected from three different depth (0-15, 15-30 and 30-45 cm) and the soil type is black cotton soil, the pH was estimated as 6.32, the organic carbon content was 0.57%, and the phosphorus content was 16.84 kg/ha. The findings showed that PSB inoculation further increased phosphorus availability (pooled maximum: 19.16 kg/ha with combined soil and seedling inoculation), even though phosphorus levels considerably improved accessible phosphorus content (pooled maximum: 18.94 kg/ha at 50 kg P/ha). A slight difference in bulk density, approximately 0.02 g/cm<sup>2</sup>, was observed during the research trial period in the same way, water holding capacity showed a negligible increase of 0.05%. The findings underscore the potential of integrated P and PSB management to enhance soil nutrient availability, particularly phosphorus, within tropical agroforestry systems, promoting sustainable agriculture practices.

**Keywords:** Nutrient use efficiency; agro-ecological practices; sustainable agriculture; phosphorus solubilizing bacteria; soil nutrient dynamics.

## 1. INTRODUCTION

Agroforestry systems are increasingly recognized for their ability to improve soil fertility, biodiversity, and ecosystem sustainability. Among them, *Gmelina arborea*-based agroforestry is notable for enhancing soil health and crop productivity. Integrating mustard (*Brassica juncea* L.) as an intercrop benefit both oilseed production and soil fertility through nutrient cycling, but success depends on understanding soil nutrient dynamics, particularly phosphorus (P) and bio-fertilizers like phosphate solubilizing bacteria (PSB) (Nair, 1993).

Phosphorus is vital for plant growth, influencing root development and energy transfer, but its availability is limited in many tropical soils due to fixation in insoluble forms (Hinsinger, 2001). Conventional P fertilization often leads to inefficiencies, with much of the phosphorus becoming chemically bound and inaccessible to plants (Gyaneshwar et al., 2002). This inefficiency and environmental concerns like eutrophication highlight the need for sustainable alternatives, such as PSB, which solubilize insoluble phosphorus, enhancing its availability for plant uptake. The limitations of traditional phosphorus fertilization, coupled with environmental concerns like leaching and eutrophication, underscore the need for more sustainable nutrient management strategies (Reddy et al., 2002).

PSB's role in improving phosphorus bioavailability is crucial in agroforestry systems

where soil nutrient competition between trees and crops is common. PSB not only promotes phosphorus solubilization but also influences soil properties such as pH, organic matter content, and microbial activity, enhancing nutrient retention and cycling (Whitelaw, 2000; Vessey, 2003). The combination of phosphorus fertilization and PSB in *Gmelina* agroforestry systems can improve physio-chemical soil properties like pH, organic carbon, and cation exchange capacity (CEC), promoting sustainable agriculture by reducing chemical input reliance (Khan et al., 2007). Understanding these interactions is critical for optimizing nutrient management in agroforestry systems. In particular, *Gmelina arborea* known for its fast-growing nature and adaptability can improve soil structure and nutrient cycling (Montagnini & Nair, 2004).

In *Gmelina*-based systems, phosphorus management is critical for maintaining productivity, particularly in tropical soils where P mobility is restricted (Richardson et al., 2009). While chemical fertilizers are commonly used, their inefficiency and environmental impact (e.g., phosphorus runoff leading to eutrophication) suggest the need for integrating PSB into nutrient strategies to enhance P solubilization and plant growth (Schroder et al., 2011; Sharma et al., 2013).

PSB also enhances CEC, organic matter accumulation, and microbial activity, improving soil nutrient retention and reducing competition

between tree and crop roots (Brady & Weil, 2008; Singh & Reddy, 2011). Additionally, PSBs release enzymes that break down organic phosphorus compounds, contributing to nutrient cycling and benefiting mustard and *Gmelina* (Malik et al., 2013; Zaidi et al., 2009). PSB inoculation can also improve soil structure, texture, and water-holding capacity, promoting better root penetration and reducing erosion (Lal, 2004).

In conclusion, phosphorus fertilization combined with PSB inoculation enhances mustard cultivation while supporting the sustainability of *Gmelina* agroforestry systems. This integrated approach reduces chemical inputs, improves soil health, and aligns with global goals of sustainable agriculture and food security (Tilman et al., 2011).

This study aims to evaluate the effects of phosphorus and PSB on soil properties in mustard cultivation within *Gmelina* agroforestry systems, focusing on soil pH, organic carbon, CEC, and microbial activity to optimize nutrient management and enhance productivity in tropical agroforestry systems.

## 2. MATERIALS AND METHODS

**Study site:** The experiment was conducted in the *Gmelina arborea* based agroforestry system at the Research Farm, Department of Forestry, College of Agriculture, Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur, Madhya Pradesh. The experimental site is situated at an altitude of 391 meters above sea level. The location of the area is at a latitude of 23° 12' 50" north and a longitude of 79° 57' 56" east in the Kymore Plateau and Satpura Hill agro-climatic zones of Madhya Pradesh. The climate is defined by extremely hot and dry summers, with an average highest temperature of 46°C and extremely cold and dry winters, with an average lowest temperature of 4°C. Jabalpur receives an average annual precipitation of 1350 mm. The region is famous for its high relative humidity levels, which reach 80 to 90% during the rainy season, 60 to 75% during the summer and 20 to 23% during the winter. The experiment was conducted from November to April in two consecutive years 2022-2023 and 2023-2024.

The experiment employed a split plot design, with Phosphorus levels ( $P_{30} = 30$  kg/ha,  $P_{40} = 40$  kg/ha and  $P_{50} = 50$  kg/ha) as main plot treatment, PSB inoculation methods namely P1

= Soil application, P2 = Seedling application and P3 = Soil and seedling application as sub-plot treatments. The experiment consisted of using 12 distinct treatment combinations (P30P1, P30P2, P30P3, P30P4, P40P1, P40P2, P40P3, P40P4, P50P1, P50P2, P50P3, P50P4). The treatments were allocated randomly into three distinct replications.

Mustard was grown in the plots measuring 6.3 m x 6.6 m between the alleys of 8-year-old *Gmelina arborea* trees. The trees are planted with a uniform distance of 8 x 2.5 m. Recommended dose of fertiliser (80:40:40 N: P: K kg ha<sup>-1</sup>, respectively) was supplemented to the crop.

**Soil sampling and analysis:** To assess the initial soil condition, a soil sample was collected before planting the crop for the field experiment. This sample provided data on the soil's physico-chemical parameters of the site. Five different sampling sites were randomly selected within each area to obtain spatial diversity. The samples were taken with an auger from the root zone at depths of 0–15 cm, 15–30 cm, and 30–45 cm. Following the collection of soil samples, they were transported to the laboratory and subjected to various tests for various characteristics. The soil samples were pulverised into smaller particles and any unwanted material was eliminated in order to achieve homogeneity. After letting the soil settle for 30 minutes at a 1:2.5 soil to water ratio, the pH of the soil was measured using a glass electrode on a digital pH meter Singh and Reddy (2011). A conductivity meter (Subbiah and Asijah 1956) was used to measure the electrical conductivity of the soil sample at 25°C in a suspension of 1:2.5 dirt to water. The method created by Walkley and Black (1934) was used to determine the amount of organic carbon. The amount of nitrogen present in the soil was ascertained using the alkaline potassium permanganate method, as explained by Subbiah and Asija (1956). In order to extract the available phosphorous, the process outlined by Olsen et al. (1954) was undertaken. As explained by Jackson (1973), the amount of potassium that was available was evaluated by extracting it using a 1 N ammonium acetate solution at pH 7, and the potassium concentration was determined using a flame photometer. The hot-water solubility method created by Gupta (1967) was used to estimate the boron availability, and it was made even simpler by employing azomethine-H (Zaidi et al., 2009).

### 3. RESULTS

**Effect of phosphorus and PSB on bulk density of soil:** Bulk density analysis at various soil depths (0–15 cm, 15–30 cm, and 30–45 cm) showed no statistically significant differences caused by PSB inoculation techniques or phosphorus amounts (30, 40, and 50 kg/ha). In Table 2, across both years and pooled data, the results varied somewhat but stayed between 1.34 and 1.41 g/cm<sup>3</sup>. According to these findings, soil compaction and structure at the investigated depths are barely affected by PSB inoculation techniques and phosphorus treatment. The 1.41 g/cm<sup>3</sup> initial bulk density stayed mostly constant, confirming the treatments' negligible impact.

**Effect of phosphorus and PSB on water holding capacity of soil:** In Table 2, additionally, there were no appreciable variations in soil water-holding capacity as a result of different PSB inoculation techniques or phosphorus levels. The water holding capacity varied from 37.97% to 38.47% among years and pooled data, despite slight increases. These gains were not statistically significant, however the maximum capacity was noted below 50 kg P/ha. In a similar vein, although the results were not statistically significant, the combined soil + seedling inoculation strategy produced somewhat higher values. These results imply that, in the conditions under investigation, neither PSB inoculation nor phosphorus administration significantly increases soil moisture retention.

**Effect of phosphorus and PSB on available nitrogen in soil:** In Table 4 conclude that phosphorus levels and PSB inoculation techniques did not significantly alter the available nitrogen levels in post-harvest soils, which ranged from 237.8 kg/ha to 270.1 kg/ha. The combination soil + seedling inoculation approach and 50 kg P/ha produced the maximum nitrogen levels. In comparison to the control, other inoculation techniques, such as soil-only and seedling-only treatments, produced modest increases. Additionally, there was no significant relationship between PSB inoculation techniques and phosphorus levels.

**Effect of phosphorus and PSB on available potassium in soil:** In Table 4 it is shown that as with nitrogen, treatments had no discernible effect on the quantities of accessible potassium. The range of potassium content was 256.3 kg/ha to 288.6 kg/ha, with the highest levels found

under the soil + seedling inoculation approach and 50 kg P/ha. Although the changes were not statistically significant, potassium levels were consistently lower in the control therapy.

**Effect of phosphorus and PSB on available phosphorus in soil:** In Table 4 showed the both phosphorus levels and PSB inoculation techniques resulted in notable increases in phosphorus availability, in contrast to nitrogen and potassium. Under 50 kg P/ha, the highest phosphorus levels (18.94 kg/ha in pooled data) were recorded, greatly outperforming lower phosphorus levels. Similarly, phosphorus availability was greatly increased by the soil + seedling inoculation approach, with pooled levels up to 19.16 kg/ha. These findings show that increasing phosphorus treatment rates and using a combination of inoculation techniques can improve soil phosphorus availability.

**Effect of phosphorus and PSB on electrical conductivity of soil:** In Table 3 depicts the measurements of electrical conductivity (EC) stayed constant and did not significantly change across treatments. The values varied by depth and year, ranging from 0.14 to 0.19 ds/m<sup>3</sup>. EC values were somewhat higher when 50 kg P/ha and the soil + seedling inoculation approach were used, but these changes were not statistically significant. The consistency of EC across treatments indicates that the ionic balance and salinity of the soil are not substantially changed by PSB inoculation techniques or phosphorus levels.

**Effect of phosphorus and PSB on soil pH of soil:** In Table 3 showed the across treatments and depths (0–15 cm, 15–30 cm, and 30–45 cm), the pH values of the soil varied from 6.31 to 6.39. Both PSB inoculation and phosphorus administration did not significantly alter pH over years or pooled data. The original pH of 6.32 stayed rather constant, indicating the soil's ability to act as a buffer and its resistance to treatment-induced alterations.

**Effect of phosphorus and PSB on organic carbon of soil:** In Table 3, across phosphorus levels and PSB inoculation techniques, notable increases in organic carbon content were noted. All depths saw a considerable rise in organic carbon levels due to the combined soil + seedling inoculation approach and higher phosphorus administration (50 kg P/ha). The combined data showed that under the most successful treatment combinations, the organic

**Table 1. Mechanical and physio-chemical analysis of soil of the experimental field**

Particulars	Unit	Initial Value	Methods of analysis
<b>2. Physical constants</b>			
Bulk density	(g/cm <sup>3</sup> )	1.36	Core sample (Black et al., 1965)
<b>3. Chemical properties</b>			
Organic carbon	(%)	0.57	Rapid titration method (Walkley and Black, 1934)
Available N	(kg ha <sup>-1</sup> )	220.5	Alkaline potassium permanganate method (Subbiah and Asija, 1956)
Available P <sub>2</sub> O <sub>5</sub>	(kg ha <sup>-1</sup> )	16.84	0.5 M NaOCH <sub>3</sub> extractable (Olsen et al. 1954)
Available K <sub>2</sub> O	(kg ha <sup>-1</sup> )	243.7	1N neutral ammonium acetate method (Piper, 1966)
pH		6.32	Glass Electrode pH meter (Jackson, 1973)
Electrical conductivity	(dSm <sup>-1</sup> )	0.14	Systronics Electrical conductivity Meter (Jackson, 1973)

**Table 2. Physical Properties of Soil**

Treatments	Bulk density (g/cm <sup>3</sup> )									Water holding capacity (%)		
	0-15cm			15-30cm			30-45cm			Initial value: 37.56%		
Phosphorus levels (kg/ha)	1 <sup>st</sup> Year	2 <sup>nd</sup> Year	Pooled	1 <sup>st</sup> Year	2 <sup>nd</sup> Year	Pooled	1 <sup>st</sup> Year	2 <sup>nd</sup> Year	Pooled	1 <sup>st</sup> Year	2 <sup>nd</sup> Year	Pooled
Phosphorous 30	1.38	1.37	1.37	1.40	1.42	1.41	1.41	1.45	1.43	37.24	37.46	37.35
Phosphorous 40	1.36	1.32	1.34	1.38	1.40	1.39	1.39	1.43	1.41	38.08	37.56	37.82
Phosphorous 50	1.34	1.36	1.35	1.36	1.38	1.37	1.37	1.41	1.39	37.47	38.47	37.97
<b>SEm ±</b>	0.10	0.11	0.10	0.09	0.11	0.10	0.10	0.12	0.11	1.15	1.17	1.16
<b>C.D. at 0.05</b>	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
<b>Inoculation with PSB</b>												
Soil	1.36	1.35	1.35	1.38	1.40	1.39	1.39	1.43	1.41	38.30	37.97	38.13
Seedling	1.37	1.34	1.36	1.39	1.41	1.40	1.40	1.44	1.42	36.99	37.40	37.19
Soil + Seedling	1.36	1.35	1.36	1.38	1.40	1.39	1.39	1.43	1.41	38.10	38.47	38.29
Control	1.34	1.35	1.35	1.36	1.38	1.37	1.37	1.41	1.39	36.99	37.48	37.23
<b>SEm ±</b>	0.07	0.09	0.08	0.06	0.08	0.07	0.09	0.07	0.08	0.92	1.04	0.97
<b>C.D. at 0.05</b>	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS

**Table 3. Mechanical Properties of Soil**

Treatments	EC (ds/m)			SOIL pH			Organic Carbon (%)		
	Initial value: 0.14 ds/m			Initial value: 6.32			Initial value: 0.48		
Phosphorus levels (kg/ha)	1 <sup>st</sup> Year	2 <sup>nd</sup> Year	Pooled	1 <sup>st</sup> Year	2 <sup>nd</sup> Year	Pooled	1 <sup>st</sup> Year	2 <sup>nd</sup> Year	Pooled
Phosphorous 30	0.17	0.15	0.16	6.08	6.07	6.08	0.58	0.61	0.60
Phosphorous 40	0.14	0.12	0.13	6.35	6.34	6.35	0.61	0.65	0.63
Phosphorous 50	0.16	0.14	0.15	6.34	6.33	6.33	0.63	0.67	0.65
<b>SEm ±</b>	0.02	0.01	0.02	0.33	0.25	0.31	0.009	0.010	0.008
<b>C.D. at 0.05</b>	NS	NS	NS	NS	NS	NS	0.036	0.039	0.034
<b>Inoculation with PSB</b>									
Soil	0.15	0.13	0.14	6.27	6.26	6.26	0.62	0.66	0.64
Seedling	0.16	0.14	0.15	6.32	6.31	6.32	0.60	0.64	0.62
Soil + Seedling	0.16	0.14	0.15	6.17	6.16	6.16	0.67	0.69	0.68
Control	0.15	0.13	0.14	6.26	6.25	6.26	0.53	0.57	0.55
<b>SEm ±</b>	0.03	0.04	0.03	0.22	0.20	0.21	0.011	0.010	0.010
<b>C.D. at 0.05</b>	NS	NS	NS	NS	NS	NS	0.031	0.031	0.031

**Table 4. Chemical Properties of Soil**

Treatments	Available Nitrogen (kg/ha)			Available Phosphorus (kg/ha)			Available Potassium (kg/ha)		
	1 <sup>st</sup> Year	2 <sup>nd</sup> Year	Pooled	1 <sup>st</sup> Year	2 <sup>nd</sup> Year	Pooled	1 <sup>st</sup> Year	2 <sup>nd</sup> Year	Pooled
Phosphorous 30	246.7	249.7	248.2	17.75	18.00	17.87	265.7	268.5	267.1
Phosphorous 40	258.0	261.0	259.5	18.13	18.37	18.25	277.0	279.0	278.0
Phosphorous 50	267.4	270.4	268.9	18.82	19.06	18.94	286.4	288.4	287.4
<b>SEm ±</b>	13.41	14.46	13.81	0.20	0.18	0.19	13.41	11.68	12.24
<b>C.D. at 0.05</b>	NS	NS	NS	0.77	0.70	0.73	NS	NS	NS
<b>Inoculation with PSB</b>									
Soil	258.7	261.5	260.3	18.18	18.42	18.30	277.6	279.7	278.7
Seedling	265.5	268.7	267.4	18.31	18.55	18.43	284.7	287.8	286.3
Soil + Seedling	268.7	271.6	270.2	19.04	19.28	19.16	287.8	289.7	288.7
Control	236.3	239.3	237.8	17.41	17.65	17.53	255.33	257.3	256.3
<b>SEm ±</b>	18.28	19.65	18.94	0.34	0.36	0.35	18.27	23.07	20.50
<b>C.D. at 0.05</b>	NS	NS	NS	1.01	1.07	1.04	NS	NS	NS

carbon content increased from an initial value of 0.48% to as high as 0.69%. Improved organic matter turnover, improved root development, and increased microbial activity are the causes of these changes.

#### 4. DISCUSSION

Several significant findings were obtained from the examination of soil characteristics under different phosphorus concentrations (30, 40, and 50 kg/ha) and phosphate-solubilizing bacteria (PSB) inoculation techniques. There were no statistically significant variations in bulk density or water-holding capacity between years, pooled data, or soil depths (0–15 cm, 15–30 cm, and 30–45 cm). Between 1.34 and 1.41 g/cm<sup>3</sup>, bulk density stayed constant, suggesting that there were few impacts on compaction or soil structure. Similar to this, water-holding capacity varied between 37.97% and 38.47%, with somewhat higher values noted under combined soil + seedling inoculation and 50 kg P/ha. Both results are consistent with research by Choudhary (2023) and Ullah et al., (2024), who found that PSB inoculation and phosphate application had no discernible effects on bulk density or soil moisture retention. This highlights the limited impact of both treatments on physical soil metrics under comparable circumstances.

However, there were notable impacts on the amount of organic carbon and the availability of phosphorus. Under 50 kg P/ha, phosphorus availability peaked at 18.94 kg/ha; levels were further increased to 19.16 kg/ha by combining soil and seedling inoculation. Higher phosphorus levels and PSB inoculation, perhaps as a result of enhanced microbial activity and root growth, caused the organic carbon content to dramatically improve from 0.48% to 0.69%. These results are supported by Bindu et al., (2024) and Ríos-Ruiz et al., (2024), who point out that PSB and other integrated nutrient management significantly raise organic carbon and phosphorus availability, improving soil fertility. Even if their effects on physical characteristics like bulk density and water retention are still negligible, this data highlights the contribution that PSB inoculation and combined phosphorus application provide to improving soil nutrient dynamics.

#### 5. CONCLUSION

The study emphasises how diverse soil qualities are affected by phosphorus levels and PSB

inoculation techniques. Significant gains were observed in phosphorus availability and organic carbon content, but no changes were observed in bulk density, water holding capacity, accessible nitrogen, potassium, electrical conductivity, or pH. The most successful method for increasing soil fertility by raising phosphorus availability and organic matter content was the combination of soil + seedling inoculation and higher phosphorus levels (50 kg P/ha). These findings provide valuable insights for sustainable soil management practices, emphasizing the potential benefits of integrating optimized phosphorus applications with PSB inoculation to improve soil health and productivity.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

- Bindu, D., Mamatha, B., Gayathri, B., Anand, S. R., Subbarayappa, C. T., & Rohitha, D. S. (2024). The Influence of Integrated Nutrient Management on Major Nutrient Status and Physico-chemical Properties of Soil in Quinoa (*Chenopodium quinoa* Willd.). *Asian Journal of Soil Science and Plant Nutrition*, 10(3), 467-474.
- Black CA. (1965). *Methods of Soil Analysis*: CA Black, Editor-in-chief. DD Evans [and Others] Associate Editors. RC Dinauer, Managing Editor. *American Society of Agronomy*.
- Brady, N. C., & Weil, R. R. (2008). *The nature and properties of soils* (14th ed.). *Pearson Prentice Hall*.
- Choudhary, S. (2023). *Effect of Rock Phosphate Tailing and Phosphate Solubilizing Bacteria on Phosphorus Pools and Productivity of Wheat (Triticum aestivum L.)* (Doctoral dissertation, MPUAT, Udaipur).
- Gupta UC. (1967). A simplified method for determining hot-water soluble boron in podzol soils. *Soil Science*, 103, 424-28.

- Gyaneshwar, P., Kumar, G. N., Parekh, L. J., & Poole, P. S. (2002). Role of soil microorganisms in improving P nutrition of plants. *Plant and Soil*, 245(1), 83-93.
- Hinsinger, P. (2001). Bioavailability of soil inorganic P in the rhizosphere as affected by root-induced chemical changes: A review. *Plant and Soil*, 237(2), 173-195.
- Hu, Z., Wei, S., Li, W., Wu, T., Ullah, S., & Yang, M. (2024). Effect of Inoculation with Rhizosphere Phosphate-Solubilizing Bacteria on the Growth and Physiological Characteristics of *Parashorea chinensis*. *Forests*, 15(11), 1932.
- John MK, Chuah HH, Nduefeld JH. (1975). Application of improved azomethine-H method for determination of Boron in Soils and Plants. *Soil Science*, 8, 559-68.
- Khan, M. S., Zaidi, A., & Wani, P. A. (2007). Role of phosphate-solubilizing microorganisms in sustainable agriculture—A review. *Agronomy for Sustainable Development*, 27(1), 29-43.
- Lal, R. (2004). Soil carbon sequestration impacts on global climate change and food security. *Science*, 304(5677), 1623-1627.
- Malik, M. A., Zia, M. H., & Hussain, F. (2013). Phosphorus management in agroforestry systems: A review. *International Journal of Agriculture and Biology*, 15(2), 231-240.
- Montagnini, F., & Nair, P. K. R. (2004). Carbon sequestration: An underexploited environmental benefit of agroforestry systems. *Agroforestry Systems*, 61(1), 281-295.
- Nair, P. K. R. (1993). An introduction to agroforestry. *Kluwer Academic Publishers*.
- Olsen SR, Watanabe FS, Cosper HR, Larson WE, Nelson LB. (1954). Residual phosphorus availability in long-time rotations on calcareous soils. *Soil Science*, 78(2), 141-152.
- Piper CS. (1966). Soil and Plant Analysis. *Hans Publishers, Bombay, India*.
- Reddy, K. C., Zunaid, R., & Jat, R. S. (2002). Phosphorus management in pulse-based cropping systems. *Journal of Agronomy and Crop Science*, 188(1), 1-9.
- Richardson, A. E., Hocking, P. J., Simpson, R. J., & George, T. S. (2009). Plant mechanisms to optimise access to soil phosphorus. *Crop and Pasture Science*, 60(2), 124-143.
- Ríos-Ruiz, W. F., Castro-Tuanama, R., Valdez-Núñez, R. A., Torres-Bernal, L., Jave-Concepción, H. G., Daza-Pérez, A. C., & Archentti-Reátegui, F. (2024). Co-Inoculation of Phosphate-Solubilizing Bacteria and Rhizobia Increases Phosphorus Availability and Promotes the Development of Forage Legumes. *Agronomy*, 14(11), 2493.
- Rodríguez, H., & Fraga, R. (1999). Phosphate solubilizing bacteria and their role in plant growth promotion. *Biotechnology Advances*, 17(4-5), 319-339.
- Schroder, J. J., Smit, A. L., Douwes, M., & Neeteson, J. J. (2011). Phosphorus fertilization in Dutch agriculture: Trends and environmental impacts. *Soil Use and Management*, 27(3), 341-344.
- Sharma, S. B., Sayyed, R. Z., Trivedi, M. H., & Gobi, T. A. (2013). Phosphate solubilizing microbes: Sustainable approach for managing phosphorus deficiency in agricultural soils. *SpringerPlus*, 2(1), 1-14.
- Singh, B. P., & Reddy, K. S. (2011). Influence of microbial inoculants on soil properties and crop productivity in agroforestry systems. *Indian Journal of Agroforestry*, 13(1), 53-60.
- Subbiah BV, & Asijah GL. (1956). A rapid procedure for determination of available nitrogen in soil. *Current Sciences*, 25, 259-260.
- Tilman, D., Cassman, K. G., Matson, P. A., Naylor, R., & Polasky, S. (2011). Agricultural sustainability and intensive production practices. *Nature*, 418(6898), 671-677.
- Vessey, J. K. (2003). Plant growth promoting rhizobacteria as biofertilizers. *Plant and Soil*, 255(2), 571-586.
- Walkley and Black CA. (1934). An examination of the different methods for determining soil organic matter and a proposed modification of the chromic acid titration method. *Soil Science*, 37, 29-33.
- Whitelaw, M. A. (2000). Growth promotion of plants inoculated with phosphate-solubilizing fungi. *Advances in Agronomy*, 69, 99-151.

Zaidi, A., Khan, M., Ahemad, M., & Oves, M. (2009). Plant growth promotion by phosphate solubilizing bacteria. *Acta Microbiologica et Immunologica Hungarica*, 56(3), 263-284.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/128319>