



# Seasonal Incidence of Shoot and Fruit Borer [*Earias vittella* (Fab.)] on Okra [*Abelmoschus esculentus* (L.)] in Relation to Abiotic Factors

Aayush Kumar Sharma <sup>a</sup>, Lavlesh Kumar <sup>a</sup>,  
Sitaram Seervi <sup>b++\*</sup> and Meena Sanjay <sup>a</sup>

<sup>a</sup> Department of Agricultural Zoology and Entomology, RBS College Bichpuri, Agra (DR. B. R. Ambedkar University Agra) U.P., India.

<sup>b</sup> Department of Agricultural Entomology, College of Agriculture, Madhav University, Pindwara, Sirohi, Rajasthan, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/ijpss/2025/v37i15281>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/130464>

Original Research Article

Received: 23/11/2024

Accepted: 25/01/2025

Published: 27/01/2025

## ABSTRACT

The present investigation was carried out during *kharif* season 2023 at Agricultural Research Farm of RBS College Bichpuri, Agra (Uttar Pradesh). During the investigation, it has been revealed that the infestation of the shoot and fruit borer, *Earias vittella* (Fab.) on okra during the *kharif* season commenced from shoots 31<sup>th</sup> SMW (3.20%) and fruit borer 35<sup>th</sup> SMW (3.20%). The infestation

<sup>++</sup> Assistant Professor;

\*Corresponding author: E-mail: [sitaramseervi88@gmail.com](mailto:sitaramseervi88@gmail.com);

**Cite as:** Sharma, Aayush Kumar, Lavlesh Kumar, Sitaram Seervi, and Meena Sanjay. 2025. "Seasonal Incidence of Shoot and Fruit Borer [*Earias Vittella* (Fab.)] on Okra [*Abelmoschus Esculentus* (L.)] in Relation to Abiotic Factors". *International Journal of Plant & Soil Science* 37 (1):393-99. <https://doi.org/10.9734/ijpss/2025/v37i15281>.

ranged from shoot (3.20 - 29.70%) and fruit borer (3.20 - 36.42%), with the peak occurring in the 36<sup>th</sup> (29.70) and 39<sup>th</sup> (36.42) SMW and continuing until harvesting. The percentage of shoot and fruit borer infestation has a positive significant correlation with maximum temperature, minimum temperature, evening relative humidity and wind velocity. However, morning relative humidity, rainfall and sunshine hour has a negative significant correlation during *kharif* season 2023.

**Keywords:** Okra; shoot and fruit borer; abiotic factor; correlation; infestation.

## 1. INTRODUCTION

Lady finger, also known as okra, *Abelmoschus esculentus* (L.) is a popular vegetable in tropical regions like India, Nigeria, Pakistan, Cameroon, Iraq and Ghana. Despite being primarily farmed outside Europe and North America, it is known for its high folic acid, vitamin A, carbohydrates, phosphorus, and magnesium content. Okra is referred to by various regional names globally, including lady's finger in England, gumbo in the United States, guino-gumbo in Spain, guibero in Portugal, and bhindi in India. It is also grown in countries like India, Turkey, Iran, Western Africa, Yugoslavia, Bangladesh, Afghanistan, Burma, Japan, Malaysia, Brazil, Ghana, Ethiopia, and the Southern United States.

Okra plays an important role in the human diet and is a good source of protein, carbohydrates, vitamins, calcium, potassium, enzymes, and total minerals which are often lacking in the diet of developing countries (Saifullah et al., 2009 & Gopalan et al., 2007). "The composition of okra pods per 100 gm edible portion is water 88.6 g, energy 144.00 kJ (36 kcal), protein 2.10 g, carbohydrate 8.20 g, fat 0.20 g, fiber 1.70 g, Ca 84.00 mg, P 90.00 mg, Fe 1.20 mg,  $\beta$ -carotene 185.00  $\mu$ g, riboflavin 0.08mg, thiamin 0.04 mg, niacin 0.60 mg, ascorbic acid 47.00 mg. The fruits of okra have reawakened beneficial interest in bringing this crop into commercial production. Its medicinal value has also been reported in curing ulcers and relief from haemorrhoids. It has also been found to be medically beneficial as a plasma replacement or blood volume expander and also useful in genitourinary disorders, spermatorrhea and chronic dysentery" (Gemedede et al., 2015 & Basnet et al., 2023). "Among all pests, shoot and fruit borer (*Earias insulana* and *E. vittella*) the most destructive pests of okra, as young larva bores into tender shoots in early vegetative growth of plants and grown-up larvae damage fruits resulting in serious loss in yield. *E. vittella* is an oligophagous pest and its main hosts are okra and cotton" (Rajendran et al., 2018). It is also found feeding on a large number of cultivated and wild malvaceous plants species.

The pest is active almost the year-round and prefers high humidity and high temperature for its growth and development. The affected fruits are rendered unfit for human consumption, as well as for procurement of seed. The borer has been reported to cause 24.6 to 26.0 per cent damage to okra shoots (Pareek et al. & Zala et al., 1999) and 40 to 100 per cent loss to fruits (Pareek et al., 2003, Shinde et al., 2007 & Rai et al., 2014). "The pest is active almost year-round and prefers high humidity and high temperatures for its growth and development. In India, an estimated loss of 69.00 per cent in marketable yield was due to the attack of this insect on okra alone" (Ray et al., 2019, Anonymous 2021).

The incidence of *E. vittella* shows violent fluctuations due to changes in climatic conditions. Information on the impact of abiotic factors on the incidence of *E. vittella* is very useful for the management of this pest in western semiarid zone of Uttar Pradesh. The seasonal incidence study will help to determine the relation between the weather factors and the population of this pest, this research study also helps okra to understanding pest dynamics, developing effective management strategies, and reducing crop losses. The study helps identify peak infestation periods, informing targeted pest management interventions and contributing to sustainable agriculture practices. Keeping in view the importance of okra and the incidence of present work was carried out to know the seasonal incidence of pest.

## 2. MATERIALS AND METHODS

The field experiment was carried out during the *kharif* season 2023-24 at Agricultural Research Farm of Raja Balwant Singh College Bichpuri, Agra (U.P). The Research Farm is situated about 11 km to the west of Agra on Agra Bharatpur Road at latitude of 27°2' N and longitude of 77°9' E with an elevation of 163.4 m above the mean sea level. Ankur-41 cultivar a commonly grown and recommended in this area was sown on the first week of July 2023 with a plots size 4.5×1.5 m<sup>2</sup> and keeping row to row and plant to plant

**Table 1. Seasonal incidences of shoot and fruit borer (*Earias vittella*) on okra in relation to abiotic factor during *kharif* 2023**

S. NO.	Date of Observation	SMW	Per cent shoot infestation	Per cent fruit infestation	Temperature (oC)		Relative Humidity (%)		Rainfall (mm)	Sun shine hr/day	Wind velocity (km/hr)
					Maximum	Minimum	Morning	Evening			
1	16 Jul 23	29 <sup>th</sup>	0	0	36	26.2	76	49	36	7	10
2	23 Jul 23	30 <sup>th</sup>	0	0	38.1	26.4	73	48	16	8	12
3	30 Jul 23	31 <sup>st</sup>	3.2	0	35.3	25.2	71	47	15	7	14
4	6 Aug 23	32 <sup>nd</sup>	9.2	0	33.6	24.7	78	48	26	7	15
5	13 Aug 23	33 <sup>rd</sup>	14.3	0	37.3	26.1	71	46	10	7	21
6	20 Aug 23	34 <sup>th</sup>	18.4	0	37.3	26.1	69	50	06	6	14
7	27 Aug 23	35 <sup>th</sup>	15.6	3.2	36.8	26.2	58	32	0	7	20
8	3 Sep 23	36 <sup>th</sup>	29.7	7.9	38.5	25.6	63	42	0	0.7	17
9	10 Sep 23	37 <sup>th</sup>	25.6	22.54	36.1	25.6	67	50	04	4	9
10	17 Sep 23	38 <sup>th</sup>	16.5	27.45	33.6	24.4	75	49	09	7	10
11	24 Sep 23	39 <sup>th</sup>	14.8	36.42	36.4	25.3	64	39	01	5	9
12	1 Oct 23	40 <sup>th</sup>	15.0	34.3	37.8	23.1	72	44	0	2	6
13	8 Oct 23	41 <sup>st</sup>	10.6	30.4	37.3	22.1	48	28	0	1.5	9
14	15 Oct 23	42 <sup>nd</sup>	7.7	32.4	34.7	21.1	52	27	02	0	9
15	22 Oct 23	43 <sup>rd</sup>	5.4	31.36	32.7	18.1	54	24	0	2	8
Mean			12.4	15.06	36.1	24.41	66.06	41.53	8.34	4.74	12.2




Fig. 1. Seasonal incidences of shoot and fruit borer (*Earias vittella*) on okra in relation to abiotic factor during kharif 2023

**Table 2. Correlation analysis between seasonal incidence of shoot and fruit borer and abiotic factors during *kharif* 2023**

<b>Abiotic factors</b>	<b>Correlation (r) Per cent shoot infestation</b>	<b>Correlation (r) Per cent fruit infestation</b>
Maximum temperature (°C)	0.319*	-0.242*
Minimum temperature (°C)	0.222*	-0.674*
Morning Relative Humidity (%)	-0.091*	-0.513*
Evening Relative Humidity (%)	0.136*	-0.532*
Rainfall (mm)	-0.559*	-0.611*
Sun shine hr/day	-0.321*	-0.667*
Wind velocity (km/hr)	0.213*	-0.748*

distance of 45×30 cm, two seeds sown in one place at 2-3 cm depth on first week of July 2023 respectively. The crop was left for the natural infestation. The data of shoot and fruit borer, *Earias spp.* were recorded on the five randomly selected and tagged plant throughout the crop period at weekly intervals by a visual count of the plant in which the top portion was damaged, starting after two weeks of sowing to last picking of the fruits.

The per cent shoot infestation was calculated by counting the total number of shoots and the number of damaged shoots. The per cent infestation of fruits on a number basis was calculated by counting the infested and healthy fruits separately from selected tagged plants at each picking till the last picking of fruits. The data recorded on shoot and fruit damage and meteorological parameters were used for statistical analysis (Panse, 1967). The simple correlation was computed between shoot and fruit damage in relation to abiotic factors, viz.; maximum and minimum temperature, relative humidity and rainfall. The following formulae were used for calculating the correlation co-efficient is given below (Gupta, 1996).

$$\text{Per cent shoot damage} = \frac{\text{Total number of damage shoots}}{\text{Total number of shoots}} \times 100$$

$$\text{Per cent fruit damage} = \frac{\text{Total number of damage fruit}}{\text{Total number of fruit}} \times 100$$

$$\text{Correlation coefficient} = \frac{\text{cov}(xy)}{\sqrt{\text{var}(x)\text{var}(y)}}$$

Where,

r = correlation coefficient.

Cov (xy) = covariance of the characters X and Y.

Var (x) and (y) = variance of the characters of X and Y, respectively.

### 3. RESULTS AND DISCUSSION

The data on the infestation of shoot and fruit borer *Earias vittella* (Fab.) on okra crop during the *khariif* season is presented in (Table 1 and Fig. 1). The infestation ranged from 3.2 per cent (31<sup>st</sup> SMW) to 29.70 per cent (36<sup>th</sup> SMW). The incidence of shoot borer was observed in the 5<sup>th</sup> week after and it was continued up to harvesting. The peak incidence of shoot borer was recorded at 29.07 per cent shoot infestation at 36<sup>th</sup> SMW. The population was decreasing. The incidence of shoot and fruit borer ranged from 3.2 per cent to

36.42 per cent during the 38<sup>th</sup> to 42<sup>nd</sup> SMW. The incidence of the fruit borer on the fruits started from 38<sup>th</sup> SMW coinciding with the setting of fruits. During pest incidence increased and recorded a peak of 36.42 per cent at 39<sup>th</sup> SMW.

The peak incidence of shoot borer was recorded at 29.07 per cent shoot infestation at 36<sup>th</sup> SMW. Thereafter the population was decreasing. The above results corroborated the findings of earlier researchers, Kantipudi et al., (2017) who observed the highest per cent shoot infestation in the second week of September. Similarly, Gangwar and Singh (2014) reported that *Earias vittella* infestation started from the last week of August and continued till the last week of December *i.e.*, this pest was found infesting the crop throughout the crop season.

The result indicated that maximum temperature positive correlation with shoot infestation (r = 0.319), negative correlation with fruit infestation (r = -0.242). Minimum temperature positive correlation with shoot infestation (r = 0.222), negative correlation with fruit infestation (r = -0.674). Morning relative humidity negative correlation with both shoot infestation (r = -0.091) and fruit infestation (r = -0.513). Evening relative humidity positive correlation with shoot infestation (r = 0.136), negative correlation with fruit infestation (r = -0.532). Rainfall negative correlation with both shoot infestation (r = -0.559) and fruit infestation (r = -0.611). Sunshine hours per day negative correlation with both shoot infestation (r = -0.321) and fruit infestation (r = -0.667). Wind velocity positive correlation with shoot infestation (r = 0.213), negative correlation with fruit infestation (r = -0.748) (Table 2).

### 4. CONCLUSION

The study reveals that the *Earias vitella* infestation on okra crop during the *khariif* season 2023 with a peak infestation in 36<sup>th</sup> and 39<sup>th</sup> SMW. The seasonal incidence of shoot and fruit borer, *Earias vittella* (Fab.) on okra maximum and lowest infestation observed in various meteorological weeks, correlation studies indicated that there was a significant and non-significant influence of different weather factors and field infestation.

### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image

generators have been used during writing or editing of this manuscript.

## ACKNOWLEDGEMENTS

The authors are thankful to Head, Department of Agricultural Zoology and Entomology, RBS College Bichpuri, Agra Uttar Pradesh, for providing necessary facilities and support for carrying out the study successfully.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

- Anonymous. (2021). *Agricultural statistics at a glance*. Ministry of Agriculture and Farmers' Welfare, GOI.159
- Basnet, S., Lamichane, P., Meheta, A., & Rajbanshi, Y. (2023). A review on biochemical, nutritional, and medicinal properties of okra. *International Congresses of Turkish Science and Technology Publishing*, 46-53.
- Gemedede, H. F., Ratta, N., Haki, G. D., Woldegiorgis, A. Z., & Beyene, F. (2015). Nutritional quality and health benefits of okra (*Abelmoschus esculentus*): A review. *Journal of Food Process Technology*, 6(458).
- Gopalan, C., Sastri, S. B. V., & Balasubramanian, S. (2007). *Nutritive value of Indian foods*. National Institute of Nutrition (NIN), ICMR, India.
- Gupta, S. C. (1996). Correlation. In *Fundamentals of statistics* (pp. 510-587). Himalaya Publishing House.
- Panase, V. G., & Sukhatme, P. V. (1967). *Statistical methods for agricultural workers*. ICAR.
- Pareek, B. L., & Bhargava, M. C. (2003). Estimation of avoidable losses in vegetable crops caused by borers under semi-arid conditions of Rajasthan. *Insect Environment*, 9, 59-60.
- Pareek, B. L., Sharma, G. R., & Bhatnagar, K. N. (1986). Seasonal incidence of major insect pests of okra in semi-arid region of Rajasthan. *Annals of Arid Zone*, 25(3), 222-224.
- Rai, A. B., Halder, J., & Kodandaram, M. H. (2007). Emerging insect pest problems in vegetable crops and their management in India: An appraisal. *Pest Management in Horticultural Ecosystems*, 20(2), 113-122.
- Ray, A., Kerketta, A., Awasthi, A. K., Tomar, R. K. S., Chandan, M., & Chandravanshi, D. (2019). Impact of weather aspects on the prevalence and correlation of okra shoot and fruit borer (*Earias vittella* Fab.) during the summer season. *Journal of Entomology and Zoology Studies*, 7(5), 773-776.
- Saifullah, M., & Rabbani, M. G. (2009). Evaluation and characterization of okra (*Abelmoschus esculentus* L. Moench.) genotypes. *SAARC Journal of Agriculture*, 7, 92-99.
- Shinde, B. D., Sarkate, M. B., Memade, P. W., & Sable, Y. R. (2007). Bio-efficacy of botanical, microbial, and synthetic insecticides against okra fruit borer. *Pestology*, 31(3), 19-22.
- Zala, S. P., Patel, J. R., & Patel, N. C. (1999). Impact of weather on magnitude of *Earias vitella* infesting okra. *Indian Journal of Entomology*, 61(4), 351-355.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:  
The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/130464>