



Effectiveness of Talc Formulation (TNAUPc001) Nematode Egg Parasitic Fungus, *Pochonia chlamydosporia* on the Management of Root-knot Nematode, *Meloidogyne incognita* in Brinjal (*Solanum melongena*)

K. Senthamizh ^{a*}, V. Vijayageetha ^a, Sibi Sebastian ^a,
E. Jamuna ^a, K. Sakthivel ^b and N. Swarnakumari ^c

^a Krishi Vigyan Kendra, Tindivanam, Tamil Nadu, India.

^b Anbil Dharmalingam Agricultural College and Research Institute, Trichy, Tamil Nadu, India.

^c Horticultural College and Research Institute, Trichy, Tamil Nadu, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ijpss/2025/v37i15283>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/130341>

Original Research Article

Received: 21/11/2024

Accepted: 23/01/2025

Published: 28/01/2025

ABSTRACT

A pot culture experiment was conducted to evaluate the efficacy of talc formulation of nematode egg parasitic fungus, *Pochonia chlamydosporia* in different method of application viz., seed treatment, nursery application in growth media and soil application at the time of transplanting in

*Corresponding author: E-mail: senthamizhk@tnau.ac.in;

Cite as: Senthamizh, K., V. Vijayageetha, Sibi Sebastian, E. Jamuna, K. Sakthivel, and N. Swarnakumari. 2025. "Effectiveness of Talc Formulation (TNAUPc001) Nematode Egg Parasitic Fungus, *Pochonia Chlamydosporia* on the Management of Root-Knot Nematode, *Meloidogyne Incognita* in Brinjal (*Solanum Melongena*)". *International Journal of Plant & Soil Science* 37 (1):411-17. <https://doi.org/10.9734/ijpss/2025/v37i15283>.

brinjal for the management of *Meloidogyne incognita* at Vegetable Research Station, Palur, Tamil Nadu, India during 2020-2021. These experiments clearly indicated the effectiveness of a bio-agent *P. chlamydosporia*, TNAU (Pc001) talc formulation were more when applied in combined application viz., seed (8g/kg seed) + nurse (10/kg growth media) + soil application method (5kg/ha). The brinjal yield was significantly high in vegetable yield (880 g/plant) in brinjal (Var PLR 2), Which is 55.3 per cent increase over untreated control and recorded with lowest root-knot nematode, *M. incognita* population (81.66/250 cc soil and 8.33/ 5 g of root) which was more than 60 per cent reduction over control. The colony forming units in the soil also more (75×10^4) in the above treatments. Through these investigations we could standardize a strategy for the sustainable management of nematode by integration of seed treatment @ 8g/kg, nursery application @10g/kg growth media and soil application @5kg/ha in the main field in brinjal.

Keywords: Brinjal; *Meloidogyne incognita*; root knot nematode; *Pochonia chlamydosporia*; bio-agents.

1. INTRODUCTION

Brinjal is one of the important vegetable crop in India. Most of the vegetable crops are affected by the parasitic nematodes in all stages of the crop development. Nematode attack on the root system makes the seedlings weak and also vulnerable to the infection by secondary pathogens (soil born fungus & bacteria, Abad et al. 2003). Nematode damage also break down the resistance against pathogenic fungi and bacteria (Agrios, 2005). Further nematode infested nursery seedlings facilitate the spread of the nematodes in the main fields making the problem more difficult to manage in a larger area (Abad et al. 2023).

Meloidogyne incognita is a significant plant parasitic nematode that affects vegetables (Sasser 1952). It is a polyphagous nematode pest and has more than 3000 host plants all over the world (Noling, 1999). Frequent application of chemical nematicides for the control of soil borne plant pests, leads to the pollution of the ecosystem and thereby disturbing the ecological balance. Alternative techniques based on biological methods are needed to solve the problem. The nematode egg parasitic fungus, *Pochonia chlamydosporia* is one of the biocontrol agent against plant parasitic nematodes. The fungus is also a facultative parasite of nematode and mollusk eggs, and a hyperparasite of other fungi (Uddin et al. 2019). Plant-fungus-nematode interaction, an interaction that needs to be addressed to ensure the efficient use of *P. chlamydosporia* as a bio-pesticide as part of an integrated pest management approach.

2. MATERIALS AND METHODS

Department of Nematology, Tamil Nadu Agricultural University, Coimbatore has

developed one *P. chlamydosporia* bio product as commercial formulation viz., TNAU (Pc001) talc formulation with the spore load of spore load 10×10^8 . In the earlier studies we optimized the dose for seed treatment, nursery application in growth media and soil application during transplanting, In this research we studied the different methods of applications viz., seed treatment, nursery application and soil application during transplanting individually and combination of different methods of application were tried and the data were collected.

A pot culture experiment was taken with *P. chlamydosporia* TNAU (Pc001) talc formulation with the following treatments Viz., T1 – P.c (TNAU Pc001) talc formulation as seed treatment @ 8 g/kg seed, T2 - P.c (TNAU Pc001) talc formulation as nursery application @ 10 g/kg growth media, T3 - P.c (TNAU Pc001) talc formulation at the time of transplanting @5 kg/ha, T4 – T1 + T2; T5 – T1 + T3; T6 – T2 + T3; T7 – T1 + T2 + T3; T8 – T a (TV 1) seed treatment @ 4 g/kg + Soil application @ 2.5 kg/ha, T9 – Untreated control. Each treatment replicated three times. Completely Randomized Design were used for this trial. The brinjal var. PLR2 were taken for this study. Plants were maintained properly. Yield were recorded for each picking, Plants were uprooted at the end of the experiment. Observation on various plant growth parameters length (shoot and root) in centimeter, weight (fresh) in grams, root-knot index were recorded. The root population were observed by following root staining method (Byrd et al., 1983) The plant tissues were stained in 0.1% acid fuchsin in lacto phenol at 80°C for 2-3 minutes. Then after gentle wash, roots were kept in clear lacto phenol for 24 hours the the roots were observed through Sterio zoom microscope for counting the number of galls, number of egg masses per plant and number of eggs per egg

mass and the parasitized egg was calculated through compound microscope. For root gall index evaluating standard scale has been used as described by Taylor and Sasser (1978). After removing the plant from each pot, the soil was mixed well and 250cc soil from each pot was processed by Cobb's sieving and decanting technique (Cobb, 1918) followed by modified Baermann's funnel technique for the estimation of nematode population in soil (Christie and Perry, 1951). After termination the one gram of soil samples were collected from each replications and the Colony Forming Units (CFU) were calculated by serial dilution method (Waksman, 1922). The fungal colonies were identified and confirmed with character recorded in number. All the data viz., nematode population, Plant yield, growth parameters were processed by analysis of variance (ANOVA).

3. RESULTS AND DISCUSSION

All the treatments were significantly differ from each other, The treatment contains combination of three method of application viz., P.c (TNAU

Pc001) talc formulation as seed treatment @ 8 g/kg seed + nursery application @ 10 g/kg growth media + 5 kg/ha at the time of transplanting (T7), recorded lowest nematode population 81.66/250 cc soil and 8.33/ 5 g of root (>60 per cent reduction over control) and high vegetable yield (880 g/plant) in brinjal (Var PLR 2), Which is 55.3 per cent increase over control (Table 1). This was followed by PC (TNAU Pc001) talc formulation as nursery application @ 10 g/kg growth media +soil application @ 5kg/ha at the time of transplanting combination (T6) recorded 128.33 and 12.33 nematodes in 250 cc soil and in 5 g of root respectively and obtained the yield of 841.66 g/plant (Table 2). In this studies in general the TNAU Pc001 treated potted plants the growth of the brinjal plant and yield was significantly higher than the untreated control when compared to the untreated pots. Pravez et al. (2024) reported that *P. chlamydosporia* emerged as a plant growth promoter and biocontrol agent. *P. chlamydosporia* was effective in promoting plant growth and reducing RKN populations in soil & root in various crops (Ghahremani et al. 2019 & Nagesh and Janakiram (2004).

Table 1. Effect of different method of application of *P. chlamydosporia* TNAU Pc001 talc formulation on the yield and growth parameters of Brinjal PLR 2

Treatment details	Plant growth characters						
	Shoot length (cm)	Root Length (cm)	Shoot Weight (g)	Root Weight (g)	No of Fruits/ plant	Yield/ Plant (g)	Yield (t/ha)
T1 – P.c (TNAU Pc001) talc formulation as seed treatment @ 8 g/kg seed	42.66	12.66	63.00	12.33	15.33	616.6 (+8.8)	17.12
T2 - P.c (TNAU Pc001) talc formulation as nursery application @ 10 g/kg growth media	43.66	13.83	76.66	12.50	15.33	620.00 (+9.4)	17.22
T3 - P.c (TNAU Pc001) talc formulation at transplanting@5kg /ha	52.66	16.16	149.33	14.66	16.66	711.66 (+25.6)	19.76
T4 – T1+T2	47.33	15.16	117.33	14.66	15.33	633.33 (+11.6)	17.57
T5 – T1 + T3	62.00	17.16	175.00	22.66	20.66	816.66 (+44.1)	22.68
T6 – T2 + T3	66.66	17.50	198.33	24.00	21.33	841.66 (+48.5)	23.37
T7 – T1 + T2 + T3	82.33	18.00	249.66	25.50	24.00	880.00 (+55.3)	24.44
T8 – T a (TV 1) seed treatment @4 g/kg + Soil application @ 5 kg/ha	63.33	16.66	171.66	20.00	16.00	723.33 (+27.6)	20.08
T9 – Untreated control	26.66	9.00	21.33	4.00	14.66	566.66	15.73
CD (0.05%)	5.19	0.99	15.51	1.70	1.83	34.35	1.42

Note: Value in the parenthesis indicates percent reduction/increase over untreated control

Table 2. Effect of different method of application of *P. chlamydosporia* TNAU Pc001 talc formulation on root-knot nematode population in Brinjal PLR 2

Treatment details	Nematode population				No. of eggs parasitized	CFU of bio agent
	Nematode population /250 cc soil	Nematode RKN population index / 5 g of root	Eggs/ mass			
T1 – P.c (TNAU Pc001) talc formulation as seed treatment @ 8 g/kg seed	234.67 (-10.2)	22.00 (-13.14)	5	313.33 (-9.4)	12.00	32.66
T2 - P.c (TNAU Pc001) talc formulation as nursery application @ 10 g/kg growth media	228.33 (-12.62)	21.00 (-17.0)	5	303.00 (-12.4)	12.33	33.00
T3 - P.c (TNAU Pc001) talc formulation at transplanting @10 kg/ha	195.00 (-25.41)	17.00 (-32.88)	3	265.33 (-23.3)	13.66	58.00
T4 – T1+T2	211.66 (-19.00)	18.66 (-26.33)	4	240.00 (-30.6)	13.66	38.33
T5 – T1 + T3	161.67 (-38.13)	14.33 (-43.42)	3	158.33 (-54.33)	18.33	68.00
T6 – T2 + T3	128.33 (50.9)	12.33 (-51.32)	3	151.66 (-56.2)	26.00	68.00
T7 – T1 + T2 + T3	81.66 (-68.75)	8.33 (-67.11)	2	134.33 (-61.1)	37.00	73.00
T8 – T a (TV 1) seed treatment @ 4 g/kg + Soil application @5 kg/ha (Standard Check)	196.66 (-25.14)	17.33 (-31.58)	4	255.66 (26.10)	6.66	85.66
T9 – Untreated control	261.33	25.33	5	346.00	0	0
CD (0.05%)	41.29	2.114		15.98	4.45	6.29

Note: value in the parenthesis are percent increase / decreases over control
*value in the parenthesis are percent reduction over standard check



Fig. 1a



Fig. 1b



Fig. 1c

Fig. 1(A-C). Pot culture trial

Significantly higher number of *M. incognita* eggs in the egg mass were parasitised by *P. chlamydosporia* and higher Colony Forming Units (75×10^4) were recorded in the combined treatment of *P. chlamydosporia*, which was comparatively less than the standard check, *T. asperillum* application @ 5kg/ha which was recorded as 85×10^3 . *P. chlamydosporia* is a multi trophic microorganism because it parasitizes nematodes, colonizes plant roots and has saprophytic ability. These characteristics are desirable in a biological control agent because they allow the fungus to survive in the soil even in the absence of the nematode (Stirling, 2014).

Lopez-Lloeca et al., 2009 also stated that the *P. chlamydosporia* fungus can remain saprophytic in soil in the absence of both plant and nematode hosts. In the rhizosphere, the fungus can colonize the roots of host plants, and several *Pochonia* species have even been reported to show endophytic behavior in some Gramineae and Solanaceae species a growth habit that may result in benefits to the host plant defense against soil-borne pathogens Maciá Vicente et al. 2009. Crops and plant species that can support more than 200 colony forming units (CFU)/cm² of root are considered to be good hosts of the fungus. Good hosts for *P. chlamydosporia* include beans, cabbage,

crotalaria, pigeon pea, potato, pumpkin, and tomato. Chili, sweet potato, cowpea, rye, tobacco, and cotton are moderate hosts (100–200 CFU/cm² of root), whereas poor hosts (<100 CFU/cm² of root), include aubergine, okra, soybean, sorghum, and wheat Bourne et al, 1996).

4. CONCLUSION

The result confirmed that the nematode eggparasitic fungus *P. chlamydosporia* as talc based TNAU Pc001 bio product in combined application of seed treatment @ 8g/kg, nursery application @10g/kg growth media and soil application @ 5kg/ha during transplanting in main field in brinjal recorded with higher yield.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Abad, P., Favery, B., Rosso, M. N., & Castagnone-Sereno, P. (2003). Root-knot nematode parasitism and host response: Molecular basis of a sophisticated interaction. *Molecular Plant Pathology*, 4(3), 217–224.
- Agrios, G. (2005). *Plant pathology* (5th ed., pp. 26-27, 398-401). Elsevier Academic Press.
- Bourne, J. M., Kerry, B. R., & Leij, F. (1996). The importance of the host plant on the interaction between root-knot nematodes (*Meloidogyne* spp.) and the nematophagous fungus *Verticillium chlamydosporium* Goddard. *Biocontrol Science and Technology*, 6(4), 539–548.
- Byrd, D. W., Kirkpatrick, T., & Barker, K. R. (1983). An improved technique for clearing and staining plant tissue for detection of nematodes. *Journal of Nematology*, 14(2), 142–143.
- Christie, J. R., & Perry, V. G. (1951). Removing nematodes from soil. *Proceedings of the Helminthological Society of Washington*, 18, 106–108.
- Cobb, N. A. (1918). Estimating the nematode population of soil. *US Department of Agriculture, Bureau of Plant Industry, Agricultural Technical Circular*, 1, 1–48.
- Ghahremani, Z., Escudero, N., Saus, E., Gabaldón, T., & Sorribas, F. J. (2019). *Pochonia chlamydosporia* induces plant-dependent systemic resistance to *Meloidogyne incognita*. *Frontiers in Plant Science*, 10, 945. <https://doi.org/10.3389/fpls.2019.00945>
- Lopez-Llorca, L. V., Olivares-Bernabeu, C., Salinas, J., Jansson, H. B., & Kolattukudy, P. E. (2009). Pre-penetration events in fungal parasitism of nematode eggs. *Mycological Research*, 106(4), 499–505.
- Maciá-Vicente, J. G., Rosso, L. C., Ciancio, A., Jansson, H. B., & Lopez-Llorca, L. V. (2009). Colonization of barley roots by endophytic *Fusarium equiseti* and *Pochonia chlamydosporia*: Effects on plant growth and disease. *Annals of Applied Biology*, 155(3), 391–401.
- Nagesh, K., & Janakiram, T. (2004). Root knot nematode problem in polyhouse roses and its management using dazomet, neem cake, and *Pochonia chlamydosporia* (*Verticillium chlamydosporium*). *Journal of Ornamental Horticulture New Series*, 7(2), 147–152.
- Noling, J. W. (1999). Nematode management in tomatoes, peppers, and eggplant. *Florida Cooperative Extension Service ENY No. 028*. University of Florida.
- Prevez, R., Rajkumar, C., Chandiramani, D., Waghmare, R., & Surekha, R. (2024). Management of root-knot nematode *Meloidogyne incognita* infesting turmeric (*Curcuma longa*) under coconut (*Cocos nucifera*) cropping system. *Indian Journal of Agricultural Sciences*, 94, 422–426.
- Sasser, J. N. (1952). Identification of root-knot nematodes (*Meloidogyne* spp.) by host reaction. *Plant Disease Reporter*, 36, 84–86.
- Stirling, G. (2014). *Biological control of plant parasitic nematodes in sustainable agriculture* (2nd ed., Vol. 24). CABI Publishing.
- Taylor, A., & Sasser, J. (1978). *Biology, identification and control of root-knot nematodes*. North Carolina State University Graphics.
- Uddin, M. N., Saifullah, S., Ahmad, M., Khan, W., & Khan, B. M. (2019). Evaluation of *Pochonia chlamydosporia* (Goddard) isolates for suppression of *Meloidogyne*

incognita, root-knot nematode of tomato. Waksman, S. A. (1922). A method for counting the number of fungi in the soil. *Journal of Bacteriology*, 7, 339–341.

The Journal of Agricultural Sciences, 11, 70–81.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/130341>