



Studies on Analysis of Correlation Coefficient and Path Coefficient for Certain Quantitative Traits in Fieldpea (*Pisum sativum* L. var. *arvense*)

Umesh ^{a*}, Alok Kumar Singh ^b, Kanhaiya Lal ^a,
Sachhidanand Tripathi ^a, Abhishek Kumar ^a,
Vishal Singh ^a, Anurag Pandey ^a, Abhay Singh ^a
and Deepak Yadav ^c

^a Department of Genetics and Plant Breeding, Faculty of Agriculture, Kamla Nehru Institute of Physical and Social Sciences, Sultanpur, 228118, U.P., India.

^b Kamla Nehru Institute of Physical and Social Sciences, Sultanpur, 228118, U.P., India.

^c Chandra Shekhar Azad University of Agriculture and Technology, Kanpur, 208002, U.P., India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ijpss/2025/v37i25309>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://pr.sdiarticle5.com/review-history/131101>

Original Research Article

Received: 03/12/2024

Accepted: 05/02/2025

Published: 08/02/2025

ABSTRACT

The experiment was conducted with twenty-two (22) genotypes of fieldpea including four (4) checks in Randomized Block Design with three replications at the experimental farm of department of Genetics and Plant Breeding, Faculty of Agriculture, Kamla Nehru Institute of Physical and Social

*Corresponding author: E-mail: umeshkashyap00365@gmail.com;

Cite as: Umesh, Alok Kumar Singh, Kanhaiya Lal, Sachhidanand Tripathi, Abhishek Kumar, Vishal Singh, Anurag Pandey, Abhay Singh, and Deepak Yadav. 2025. "Studies on Analysis of Correlation Coefficient and Path Coefficient for Certain Quantitative Traits in Fieldpea (*Pisum Sativum* L. Var. *Arvense*)". *International Journal of Plant & Soil Science* 37 (2):106-113. <https://doi.org/10.9734/ijpss/2025/v37i25309>.

Sciences Sultanpur 228118 (U.P.) India, during Rabi, season 2022-2023. Phenotypic data were recorded for nine characters viz. Days to 50% flowering, Plant height, Branches per plant, Number of pods per plant, Days to maturity, Harvest index, 100-seed weight, Biological yield per plant and Grain yield per plant. It was expressed in general Correlation coefficient, provide an indication of the type and extent of the relationship between yield and yield contributing components by measuring the magnitude of linear relationships between two variables and path coefficient analysis partitions the observed correlation coefficient into direct and indirect effects on grain yield. The maximum value of highly significant positive genotypic correlation was observed for days to 50% flowering with days to maturity (0.88), biological yield per plant with grain yield per plant (0.84), number of branches per plant with 100-seed weight (0.84) respectively. The high direct positive genotypic effect on grain yield per plant were exerted by days to 50 % flowering (0.78) followed by biological yield per plant (0.40). The very high indirect positive genotypic effects on grain yield per plant by harvest index via biological yield per plant (1.47). Correlation coefficient gives an indication of the type and degree of the relationship between yield and yield contributing traits by measuring the magnitude of linear relationship between two variables, that's why it leads us in selection of yield and other desirable characteristics of plant in crop improvement program and path analysis has emerged as an important and widely used technique for understanding the direction of contribution of traits on grain yield. It contributes either directly or indirectly to the total grain yield of the plant and this technique provides information to breeders for selection of yield contributing traits for improvement of crop in plant breeding.

Keywords: Fieldpea genotypes; correlation; path analysis; direct effect; indirect effect.

1. INTRODUCTION

“Fieldpea (*Pisum sativum* L. var. *arvense*) is a mediterranean origin, the Near East and Ethiopia are considered as its secondary centre of origin” plant famous as pulse crop worldwide (Al-Jibouri et al., 1958). It is one of the most important pulse crop in India and 3rd most important pulse crop at global level, after Kidney bean and Bengal gram and 3rd most popular Rabi pulse of India after Bengal gram and lentil” India holds 4th rank in area (10.53 %) and 5th rank in production (6.96 %)” (Anonymous, 2022; Lal et al., 2011; Sharma & Sharma, 2012). “In India it is cultivated over an area of 7.45 lakh ha with a production of about 9.10 lakh tonnes during the period of 2016-17 to 2020-21 (Azam et al., 2024; Bijalwan et al., 2018). “Uttar Pradesh holds first rank as fieldpea growing state. It alone shares about (46%) of fieldpea production of India. Besides, Uttar Pradesh, Madhya Pradesh, Rajasthan, Jharkhand and Assam are the important fieldpea growing states” (Bashir et al., 2017; Blixt, 1970).

Fieldpea belongs to Leguminosae (Fabaceae) family having 2n=14 number of chromosomes. It has tap root system, herbaceous and weak stem, compound leaf with tendrils and stipules are well developed for photosynthesis, inflorescence is solitary terminal and axial type with zygomorphic bisexual flowers and fruit type is legume (Dewey & Lu, 1959; FAO, 2021).

“Fieldpea is a rich source of protein (20-23%), amino acids and carbohydrates. It is consumed alone and mixed with other vegetables but it is mainly utilized as a pulse and vegetables. It keeps special position in crop rotation with cereals because their cultivation breaks cereal disease cycles, facilitates weed control and improves soil condition and fertility by fixing atmospheric nitrogen into the soil” (Dhama et al., 2010; Gautam et al., 2017; Parveen et al., 2025).

“Field pea has high genetic yielding potential but due to various climatic factors the overall production is not good therefore it is important to identify such kind of genotypes those are able to provide maximum yield under various ecological conditions. Estimation of correlation coefficient between the yield and yield contributing traits are necessary to understand the direction of selection. Prediction of relationship between yield and yield contributing traits are indispensable to understand the direction of selection” (Kumar et al., 2015; Lal et al., 2024). “The measurements which estimate the relation between factors are called as correlation coefficient”. It quantifies the relation between various plant traits and decides the constituent parameters by them yield could be improved through selection. Yield of the crop can be improved by indirect selection of highly heritable traits, which are related with the yield (Singh, 1983; Kumar et al., 2024a).




Fig. 1. Field study at Chandpur, Sultanpur, Uttar Pradesh, India

“Path coefficient analysis is a crucial means for division of correlation coefficient into direct and indirect effects of independent variables on dependent variable. It also has been used for those characters which have vast impact on yield and correlated traits for potential use in crop improvement programmes” (Panwar et al., 2023).

2. MATERIALS AND METHODS

The experiment under relevance was conducted during Rabi 2022-23 at experimental field of, Department of Genetics and Plant Breeding, Faculty of Agriculture, Kamla Nehru Institute of Physical and Social Sciences Sultanpur 228118 (U.P.) India. The 22 collections of germplasm comprising of indigenous genotypes, varieties and advance breeding lines constituted the experimental materials for the experiment. The genotypes were obtained from the pulse section, Department of Genetics and Plant Breeding, Acharya Narendra Deva University of Agriculture and Technology, Kumarganj, Ayodhya, 224229 (U.P.) India. We randomly selected five plants from each plot for recording of data on all the nine characters. Average of the data from selected plants of each plot in respect of different characters were used for various statistical analysis. The data were recorded for the following characters. Days to 50% flowering, Plant height, Branches per plant, Number of pods per plant, Days to maturity, Harvest index,

100-seed weight, Biological yield per plant and Grain yield per plant.

3. STATISTICAL ANALYSIS

Replication wise mean data of 22 entries for 9 characters was used for statistical and biometrical analysis for the following parameters.

- Analysis of variance for Randomized Block Design was done as per the formula given by Panse and Sukhatme (1967).
- Estimation of correlation coefficient as per suggested by Al-ji-bouri et al. (1958).
- Estimation of path coefficient as per suggested by Dewey and Lu (1959).

4. RESULTS AND DISCUSSION

The analysis of variance for the design of experiment involved 22 fieldpea genotypes including 4 checks with 3 replications for the 9 characters in Randomized Block Design. The mean squares resulting from replications, treatments, and error for all characters are shown in (Table 1). The variation due to treatments were found to be highly significant for all the characters, whereas, the variation due to replications were found non-significant for all characters. Jaiswal et al., 2015 also reported highly significant variation in Fieldpea for the studied characters.

Table 1. Analysis of variance of randomized block design for different characters in Fieldpea genotypes

S.V.	d f	Days of 50% flowering	Days to maturity	Plant height (cm)	Number of Pods/plant	Number of branches/ plant	100- Seed weight (g)	Biological yield/plant(g)	Seed yield (g)	Harvest index (%)
Treatments	21	48.245**	26.7193**	2033.83**	51.120**	0.59976**	22.7272**	139.34**	59.023**	124.697**
Replication	2	5.015	1.4091	65.83*	129.523	2.10335	0.4482	576.35	95.083	18.883
ERROR	42	2.888	4.0916	18.75	44.058	0.44715	1.9033	77.64	21.264	52.513

Table 2. Analysis of simple genotypic correlation coefficient

Characters	Days to 50% flowering	Days to maturity	Plant height (cm)	Number of Branches/plant	Number of Pods / Plant	100-Seed weight	Biological Yield/ plant	Harvest Index (%)	Seed Yield/ plant (g)
Days to 50% flowering	1 **	0.8844**	0.1449 NS	-0.4238 *	-0.7244 **	-0.3342 NS	-0.4089 NS	-0.6625 **	-0.5027 *
Days to maturity		1 **	0.3615 NS	-0.3581 NS	0.1127 NS	-0.5584 **	-0.1947 NS	-0.9555 **	-0.4792 *
Plant height(cm)			1 **	-0.3322 NS	0.3966 NS	0.0011 NS	0.3308 NS	-0.4889 *	-0.0099 NS
Branches/ plant				1 **	-1.5011 **	0.8275 **	0.4168 NS	0.3565 NS	0.3736 NS
Pods/plant					1 **	0.203 NS	0.4941 *	-0.2112 NS	0.3964 NS
100- seed weight (g)						1 **	0.5568 **	0.6231 **	0.5222 *
Biological yield/ plant(g)							1 **	0.2754 NS	0.8425 **
Harvest index (%)								1 **	0.7442 **
Seed Yield/plant (g)									1 **

Table 3. Analysis of path coefficient at genotypic level

Characters	Days to 50% flowering	Days to maturity	Plant height (cm)	No. of branches / Plant	No. of Pods / plant	100-seed weight (g)	Biological yield / plant(g)	Harvest index (%)	Seed yield / plant(g)
Days to 50% flowering	0.78433	-1.22693	-0.06023	0.17275	0.08089	0.05964	-0.60348	0.29037	- 0.5027
Days to maturity	0.69370	-1.38722	-0.15021	0.14599	-0.01258	0.09964	-0.28726	0.41874	- 0.4792
Plant height (cm)	0.11369	-0.50144	-0.41554	0.13543	-0.04428	-0.00020	0.48821	0.21426	- 0.0099
No. of branches /plant	-0.33239	0.49683	0.13806	-0.40763	0.16761	-0.14766	-0.15623	0.61504	0.3736
Pods / plant	-0.56820	-0.15631	-0.16480	0.61189	-0.11166	-0.03623	0.72919	0.09254	0.3964
100-seed weight (g)	-0.26213	0.77458	-0.00046	-0.33732	-0.02267	-0.17844	0.82169	-0.27308	0.5222
Biological yield / plant(g)	-0.51966	1.32542	0.20316	-0.14531	0.02358	-0.11119	0.40646	-0.43826	0.8425
Harvest index (%)	-0.32074	0.27002	-0.13747	-0.16988	-0.05517	-0.09936	1.47577	-0.12071	0.7442

The estimates of Correlation coefficient and Path coefficient are presented in Table 2. Correlation coefficient estimates the actual relation of certain quantitative traits either significant or non-significant for grain yield. It also suggests that quantitative characters how's related to each other either positive or negative which give us direction of selection of characters in breeding programme. The correlation coefficient estimation done as per concept of Al-ji-bouri et al. (1958).

Grain yield is a complex character because it is a consequence of interaction between number of plant characters. Various yield components interact directly or indirectly with each other to produce the genetic architecture of grain yield in Fieldpea and other crops. In other words, choosing plants based on their yield alone would not matter much, unless selection in conjunction with various characteristics than that condition of the plant. Therefore, identification of the important yield contributing traits and their association with grain yield with each other is essential for developing an efficient breeding strategy in development of high yielding varieties. Correlation coefficient provide an indication of the type and degree of the relationship between yield and yield contributing traits by measuring the direction of linear relationship between two variables.

The days to 50% flowering showed highly significant positive correlations with days to maturity, biological yield per plant with grain yield per plant, branches per plant with 100-seed weight, harvest index with grain yield per plant, 100-seed weight with harvest index, 100-seed weight with biological yield per plant respectively. Significant positive correlation was found for 100-seed weight with grain yield per plant and pods per plant with biological yield per plant respectively. Characters that shown positive and significant correlation with other characters provide the information about contribution of characters in grain yield per plant in breeding programme.

“Path coefficient analysis partitions the observed correlation coefficient into direct and indirect effects on grain yield. Analysis of direction, provides a good understanding of character interactions to formulate an effective selection strategy. In contrast to a simple correlation, the route coefficient emphasizes the causes and their relative importance, while the latter tests merely the reciprocal relationship, which lacks

causation. Sewal Wright (1921) introduced the idea of a path coefficient analysis and Dewey and Lu (1959) first used the path coefficient analysis technique in selection of plants for crop improvement. To access the relative value of the different yield contributing characteristics, path coefficient analysis is an important and widely used technique for understanding the direction of contribution of traits on grain yield. It contributes either directly or indirectly to the grain yield of a plant.

“The high direct effect on grain yield per plant were exerted by days to 50 % flowering followed by biological yield per plant. The very high indirect positive effects on grain yield per plant by harvest index via biological yield per plant, followed by biological yield per plant via days to maturity. High indirect positive effect on grain yield 100-grain weight via biological yield per plant. The moderate, low and very low estimates of direct and indirect effects recorded in case of remaining characters indicated that their direct & indirect contribution to grain yield was too low to be considered of any consequence. All these characters had showed direct and indirect positive effect on grain yield give us direction for selection of characters in breeding programme therefore estimates of path coefficient between characters is very much important in breeding programme.

5. CONCLUSION

Positive and highly significant association of grain yield per plant was found with biological yield per plant and harvest index. Positive significant association with 100- grain weight while negative non – significant with plant height. Negative significant association with days to 50% flowering and days to maturity. Positive and non-significant association with branches per plant and number of pods per plant. On the basis of result, some traits considered as yield contributing traits which usher us towards selection of best yield contributing traits in our breeding programme. Plant breeder can utilize such technique for the genetic improvement of crops by selection of good yield contributing traits.

“Days to 50 % flowering exhibited highly positive direct contribution followed by biological yield per plant (g) to grain yield per plant while pods per plant, 100-grain weight, and harvest index via biological yield per plant exhibited highly indirect effect for grain yield per plant. Those characters

identified above as direct and indirect components of yield give us right direction in the formulation of effective selection strategy in field pea for developing high yielding varieties of fieldpea in breeding program.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Authors hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Al-Jibouri, H. A., Miller, P. A., & Robinson, H. F. (1958). Genotypic and environmental variances and co-variances in an upland cotton cross of interspecific origin. *Agronomy Journal*, 50, 633–636.
- Anonymous. (2022). *Annual Report 2021-22*. Government of India, Ministry of Agriculture & Farmers Welfare, Department of Agriculture and Farmers Welfare Directorate of Pulses Development.
- Azam, M. G., Sarker, U., Hossain, M. A., Mahabubul Alam, A. K. M., Islam, M. R., Hossain, N., & Alamri, S. (2024). Phenotypic diversity in qualitative and quantitative traits for selection of high yield potential fieldpea genotypes. *Scientific Reports*, 14(1), 18561.
- Bashir, I., Ishtiaq, S., Fiaz, S., & Sajjad, M. (2017). Association of yield attributing traits in pea (*Pisum sativum* L.) germplasm. *Banat's Journal of Biotechnology*, 8(15), 43–49.
- Bijalwan, P., Raturi, A., & Mishra, A. C. (2018). Character association and path analysis studies in garden pea (*Pisum sativum* L.) for yield and yield attributes. *International Journal of Current Microbiology and Applied Sciences*, 7(3), 3491–3495.
- Blixt, S. (1970). The PeaP. 181-221. In R. C. King (Ed.), *Handbook of Genetics*. Plenum Press.
- Dewey, D. R., & Lu, K. H. (1959). A correlation and path coefficient analysis of components of crested wheat grass seed production. *Agronomy Journal*, 51, 515–518.
- Dhama, S. K., Tyagi, N. K., & Singh, P. B. (2010). Interrelationship and path analysis for seed yield and its component characters under eight environments in pea (*Pisum sativum* L.). *Legume Research*, 33(2), 87–94.
- FAO. (2021). *World Food and Agriculture-Statistical Yearbook 2021*. Rome. Available: <https://doi.org/10.4060/cb4477en>
- Gautam, K. K., Syamal, M. M., Singh, A. K., & Gupta, N. (2017). Variability, character association, and path coefficient analysis of green pod yield and its related traits in pea (*Pisum sativum* L.). *Legume Research*, 40(5), 818–822.
- Kumar, G. P., N., S., Sekhar, J. C., & Chary, D. S. (2024). Assessment of genetic variability, heritability, and genetic advance in maize genotypes (*Zea mays* L.). *Journal of Experimental Agriculture International*, 46(3), 146–155.
- Kumar, R., Kumar, M., Dogra, M., & N., B. (2015). Variability and character association studies in garden pea (*Pisum sativum* var. hortense L.) during the winter season at M Id Hills of Himachal Pradesh. *Legume Research*, 38(2), 164–168.
- Lal, G. M., Meena, M. L., Chandra, K., & Singh, C. M. (2011). Assessment of genetic variability and interrelation between yield and its contributing components in field pea (*Pisum sativum* L.). *Environmental Ecology*, 29, 1235–1239.
- Lal, K., Kumar, R., Shrivastav, S. P., Kumar, A., & Singh, Y. (2024). Genetic variability, character association, and path analysis of seed yield and its contributing traits in field pea (*Pisum sativum* L. var. arvense). *International Journal of Current Microbiology and Applied Sciences*, 7(6), 1815–1820.
- Panase, V. G., & Sukhatme, P. V. (1985). *Statistical methods for agricultural workers* (361 p.). ICAR, New Delhi.
- Panwar, S., Thakur, S., Mushtaq, M., & Kumar, A. (2023). Estimation of correlation coefficient and path analysis in field pea (*Pisum sativum* L.). *International Journal of Environment and Climate Change*, 13(11), 1871–1877.
- Parveen, N., Umer, S., Tan, C., Jabbar, A., Kanwal, B., Haider, I., ... & Iqbal, R. (2025). Multivariate and association analyses of various seed yield contributing

- traits divulge genetic diversity among horticultural traits in garden pea. *Plant Molecular Biology Reporter*, 1–9.
- Sharma, B. B., & Sharma, V. K. (2012). Character association and path analysis studies for yield and Singh, B. D. (1983). *Plant breeding: Principles and methods* (pp. 123–132). Kalyani Publishers, New Delhi.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://pr.sdiarticle5.com/review-history/131101>