



Study on Genetic Diversity and Character Associations in Traditional and Indigenous Rice (*Oryza sativa* L.) Genotypes Under Rainfed

Beninu B S^{a++*}, Jenny Priya Ekka^{a#}, Arpan Sherring^{a†},
Vaidurya Pratap Sahi^{b‡}, Gera Roopa Lavanya^{a^}
and Shalini Masih^{c##}

^a Department of Genetics and Plant Breeding, SHUATS Prayagraj, India.

^b VIAET, SHUATS Prayagraj, India.

^c Department of Mathematics and Statistics, SHUATS Prayagraj, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ijpss/2025/v37i55425>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://pr.sdiarticle5.com/review-history/134628>

Original Research Article

Received: 22/02/2025

Accepted: 23/04/2025

Published: 28/04/2025

⁺⁺ MSc.;

[#] Advisor & Assistant Professor;

[†] Co-Advisor & Head and Associate Professor;

[‡] Chairperson & Professor and Dean;

[^] Member & Professor;

^{##} Member & Assistant Professor;

*Corresponding author: E-mail: bsbeninu@gmail.com;

Cite as: B S, Beninu, Jenny Priya Ekka, Arpan Sherring, Vaidurya Pratap Sahi, Gera Roopa Lavanya, and Shalini Masih. 2025. "Study on Genetic Diversity and Character Associations in Traditional and Indigenous Rice (*Oryza Sativa* L.) Genotypes Under Rainfed". *International Journal of Plant & Soil Science* 37 (5):22-27. <https://doi.org/10.9734/ijpss/2025/v37i55425>.

ABSTRACT

The nature and magnitude of genetic divergence were estimated in thirty rice landraces using Mahalanobis D^2 statistics considering twenty quantitative characters. ANOVA exhibits the presence of the considerable amount of variability among the genotypes. A high genotypic coefficient of variation (GCV) was observed for number of secondary branches. High phenotypic coefficient of variation (PCV) in biological yield per plant. High heritability coupled with high genetic advance as percentage was shown by number of grains per panicle. A considerable amount of diversity was revealed by Mahalanobis D^2 analysis. The 30 landraces were grouped into 9 clusters. In the present study the highest inter-cluster distance observed between cluster V and cluster VII (3913.55) followed by cluster I and cluster VII (3144.26) suggesting that the genotypes in these clusters may be used as parent for future hybridization programmes. The maximum intra cluster distance of shown by cluster IV (544.43) followed by cluster III (512.44) which showed maximum divergence. Traits like; days to heading, number of secondary branches and grain length were major contributors to genetic divergence. Parents showing more divergence can be crossed, which could produce high heterotic hybrids.

Keywords: Rice landraces; genetic divergence; GCV; PCV; D^2 analysis; heritability and genetic advance.

1. INTRODUCTION

Rice (*Oryza sativa* L.) is a self-pollinated, short-day, annual, monocotyledonous crop of family Poaceae (Graminae) with chromosome number $2n = 24$. Rice is considered an energising and reviving food (Rasane et al., 2015). It is the most staple and cereal component that sustains the two-third of the world population (Sen et al. 2020). The protein-fat proportion in food reduces with the degree of increase in polishing (Nawaz et al. 2018).

“Rice landraces have been developed through artificial selection imposed by farmers during the long-term domestication process. Although the global rice diversity is well characterized, few studies have conducted an in-depth analysis of genetic diversity on a local scale. India being the centre of rice diversity, has a wealth of traditional landrace varieties. Important issues concerning crop genetic diversity and its relationship with local cultures can only be answered by rigorously studying rice germplasm collected from geographically isolated regions” (Roy et al., 2014). “The landraces are valuable as they possess a huge treasure of genetic material which may prove valuable in future crop development and improvement

Programs” (Sinha and Mishra., 2013). Landraces possess significant genetic potential for enhancing desirable traits, making them valuable for agricultural improvement. Unlike highyielding varieties that have limited variability due to long-term selection, landraces maintained by farmers exhibit tremendous genetic diversity.

“For a better breeding program, the major thrust area lies in identifying desirable parents for hybridization for which knowledge about germplasm diversity and genetic relationship among breeding materials serves as valuable aid” (Kumar et al., 2017). “Genetic divergence is an efficient tool for an effective choice of parents for the hybridization programme. The parents involved in the development of varieties should be divergent. The germplasm provides immense scope for wide variability” (Banumathy et al., 2010). “Cross combinations involving divergent parents provide an opportunity for bringing together gene constellations of diverse nature and also promising segregant derivatives which result from to complementary interaction of divergent genes in parents are expected” (Kumar et al., 2017). “Therefore, the characterization of genetic diversity on existing landraces of rice reveals important traits of interest that can be utilized in rice improvement programmes” (Guru et al., 2017).

2. MATERIALS AND METHODS

The present study comprised 30 rice landraces laid in a randomized block design (RBD) with three replications conducted at the Field Experiment Centre of Department of Genetics and Plant Breeding, Allahabad School of Agriculture, U.P., during kharif, 2022. Recommended package of practices was followed accordingly. Observations were recorded for ten randomly selected plants from each replication for Plant Height (cm), Number of Tillers Per Plant, Number of Productive Tillers Per Plant, Flag Leaf Length (cm), Flag Leaf

Width (cm), Flag Leaf Area, Panicle Length (cm), Number of primary branches, Number of secondary branches, Number of Spikelets Per Panicle and Number of Grains Per Panicle. Grain Length (mm), Grain Breadth (mm), Test Weight (g) and Days to Heading, Days to Fifty percent Flowering, Days to maturity were recorded on plot wise. The data was subjected to Mahalanobis D^2 statistics to measure genetic divergence as suggested by Rao (1952).

3. RESULTS AND DISCUSSION

Analysis of variance showed significant differences for all the characters studied among thirty genotypes. A range of high GCV and PCV were recorded for biological yield per plant, grain yield per plant, and number of secondary branches. High heritability was expressed by days to heading, followed by days to fifty percent flowering, days to maturity, grain length, number of secondary branches, number of grains per panicle, number of primary branches, test weight, flag leaf length, grain breadth, flag leaf area, number of spikelets per panicle, grain yield per plant, plant height, number of tillers per plant, biological yield per plant. High genetic advance as percentage mean was recorded for a number of secondary branches, grain yield per plant,

biological yield per plant, number of grains per panicle, test weight, number of tillers per plant, number of spikelets per panicle, flag leaf area, number of primary branches, grain length, number of productive tillers per plant, days to heading, flag leaf length, days to fifty percent flowering, harvest index, days to maturity, plant height, grain breadth, and flag leaf width. The perusal of results is depicted in Table 1.

Based on D^2 values, 30 rice landraces were grouped into into 9 clusters following Tocher's method. The cluster I, II, III, IV and V were poly genotypic with 11, 5, 3, and 2 genotypes respectively (Table 2). Whereas cluster VI, VII, VIII, and IX were mono genotypic (1 genotypes each). The values of average intra and inter cluster distance (D^2) revealed that the maximum intra cluster distance of showed by cluster IV (544.43) followed by cluster III (512.44), cluster II (488.58), cluster I (451.04) and cluster V (438.61). There is no inter cluster distance recorded for Clusters VI, VII, VIII, and IX.

The highest inter cluster distance was observed between cluster V and cluster VII (3913.55), followed by cluster I and cluster VII (3476.76), cluster VII and cluster VIII (3144.26), cluster II and cluster V (2720.65), cluster IV and

Table 1. GCV, PCV, h^2 (bs) % and Gen. Adv as % of mean

Sl. No	Characters	GCV	PCV	h^2 (bs) %	Gen. Adv as % of Mean
1	Days to heading	17.52	17.58	99.30	35.96
2	Days to Fifty percent Flowering	16.39	16.46	99.20	33.63
3	Days to maturity	14.32	14.40	98.90	29.34
4	Plant Height (cm)	16.70	19.69	71.95	29.18
5	Number of Tillers Per Plant	29.54	34.98	71.34	51.41
6	Number of Productive Tillers Per Plant	25.68	36.73	48.88	36.99
7	Flag Leaf Length (cm)	18.61	20	84.81	35.30
8	Flag Leaf Width (cm)	14.11	15.01	88.42	27.34
9	Flag Leaf Area	25.06	27.01	86.08	47.89
10	Panicle length (cm)	8.29	14.35	33.43	9.88
11	Number of primary branches	20.48	21.40	91.63	40.39
12	Number of secondary branches	38.00	38.27	98.57	77.71
13	Number of Spikelets Per Panicle	26.11	28.55	83.64	49.19
14	Number of Grains Per Panicle	27.08	27.97	93.72	54.01
15	Grain Length (mm)	18.47	18.59	98.75	37.81
16	Grain Breadth (mm)	14.81	15.84	87.41	28.53
17	Test Weight (g)	26.91	28.15	91.40	53.01
18	Harvest Index	19.75	26.15	57.06	30.73
19	Biological Yield Per Plant (g)	34.91	41.75	69.91	60.13
20	Grain Yield Per Plant (g)	34.01	39.04	75.88	61.03

GCV = Genotypic coefficient of variation PCV = Phenotypic coefficient of variation h^2 (bs) % = Heritability in broad sense

Gen. Adv as % of Mean = Genetic advance as percentage mean

Table 2. Clustering of 30 genotypes by tocher method

SI. NO	Cluster	No. of Genotypes	Genotypes
1	I	11	Kalamallifool, Poongar, Njavara, Mahamaya, Zinco rice MS, Triveni, C.G zinc Rice 1, Bhadi, Dular, C G MAdhuraj, and Basmathi 370
2	II	5	TarnBhogsel 1, Parimala sale, Green rice, Kala jeera and, Kliro
3	III	5	Red rice, F R 13 A, Numali, Black Rice, and Kogli sale
4	IV	3	C G Devbhog, Tyaa pin, and IGKV Maheshwari
5	V	2	Shyamala and Purnima
6	VI	1	Assam black
7	VII	1	Anamika
8	VIII	1	Madhu scale
9	IX	1	Sheetal

Table 3. Average intra and inter cluster D² values of rice landraces

	Cluster 1	Cluster 2	Cluster 3	Cluster 4	Cluster 5	Cluster 6	Cluster 7	Cluster 8	Cluster 9
Cluster I	451.04	1954.13	1315.77	999.46	818.99	853.03	3476.76	1093.74	1394.00
Cluster II		488.58	856.15	1534.75	2720.65	1411.96	1617.72	1882.83	858.65
Cluster III			512.44	1355.53	1463.06	831.57	1539.73	1153.05	722.34
Cluster IV				544.43	1745.19	1173.45	2242.04	2263.77	1514.27
Cluster V					438.61	1024.09	3913.55	1056.62	2068.86
Cluster VI						0.00	2172.61	748.71	729.43
Cluster VII							0.00	3144.26	1453.63
Cluster VIII								0.00	727.21
Cluster IX									0.00

cluster VIII (2263.77), cluster IV and cluster VII (2242.04), cluster VI and cluster VII (2172.61), and cluster V and cluster IX (2068.86). The lowest inter cluster distance was observed between cluster III and cluster IX (722.34) followed by cluster VIII and cluster IX (727.21), cluster VI and cluster IX (729.43), and cluster VI and cluster VIII (748.71). The average intra and inter-cluster D² values are depicted in Table 3.

It is evident from divergence analysis that the highest inter-cluster distance was between cluster V and VII, followed by cluster I and VII. Indicates wide diversity between them. Genotypes from these clusters could be used as parents in the hybridization programme for the identification of best recombinants in further

segregating generations. Best recombinants may be selected for the development of improved rice cultivars. The minimum inter-cluster distance was found between cluster III and IX followed by cluster VIII and IX indicating that the genotypes belonging to these clusters were genetically similar and selection of parents from these clusters should be avoided to maintain the relatively broad genetic base.

The percentage contribution of 20 biometrical traits of 30 rice landraces towards genetic divergence was estimated and given in Table 4. The character first flowering (19.5%) had maximum contribution towards genetic divergence followed by number of secondary branches(13.9%), grain length (13.8%), days to

Table 4. Per cent contribution of 20 biometrical traits of 30 rice landraces towards genetic divergence

Charachers	Contribution (%)	Times Ranked First
Days to heading	19.5	85
Days to Fifty percent Flowering	7.6	33
Days to maturity	11.5	50
Plant Height (cm)	0.7	3
Number of Tillers Per Plant	0.6	3
Number of Productive Tillers Per Plant	0.3	1
Flag Leaf Length (cm)	2.2	10
Flag Leaf Width (cm)	3.4	15
Flag Leaf Area	1.9	8
Panicle length (cm)	0.4	2
Number of primary branches	2.2	10
Number of secondary branches	13.9	60
Number of Spikelets Per Panicle	11.2	49
Number of Grains Per Panicle	4.6	20
Grain Length (mm)	13.8	60
Grain Breadth (mm)	1.3	6
Test Weight (g)	2.7	12
Harvest Index	1.1	5
Biological Yield Per Plant (g)	0.2	1
Grain Yield Per Plant (g)	0.9	4

maturity (11.5%), spikelets per panicle (11.2%), days to fifty percent flowering (7.6%), number of grains per panicle (4.6%), leaf width (3.4%), test weight (2.7%), leaf length (2.2%), number of primary branches (2.2%), leaf area (1.9%), seed width (1.3%), harvest index (1.1%), yield per plant (0.9%), plant height (0.7%), number of tillers per plant (0.6%), panicle length (0.4%), productive tillers per plant (0.3%) and biological yield per plant (0.3%).

Crosses between divergent parents usually produce greater heterotic effect than closely related ones. The results revealed that the contribution of trait days to heading was maximum for genetic divergence, followed by number of secondary branches, grain length, days to maturity, spikelets per panicle, and days to fifty percent flowering. Higher inter-cluster distance was observed between cluster V and cluster VII followed by cluster V and VII.

Higher intra cluster distance was observed between genotypes of cluster IV

4. CONCLUSION

The investigation reported significant morphological variation, particularly in seed coat color and seed shape, among 30 rice landraces. Analysis of variance revealed significant differences in all studied traits, with Klirio standing

out as superior for grain yield per plant. Traits such as biological yield per plant, grain yield per plant, secondary branches, productive tillers, spikelets per panicle, test weight, grains per panicle, primary branches, and leaf area exhibited high genetic and phenotypic variability. These traits were considered essential for selection, with PCV surpassing GCV, suggesting less environmental influence. High heritability and genetic advance were observed for most traits, indicating their suitability for direct selection, except for productive tillers, harvest index, and panicle length. Selection based on positive correlations between biological yield, tiller numbers, productive tillers, harvest index, grain length, and fertile grains per panicle could lead to high yield and quality offspring. Path analysis highlighted the importance of biological yield, harvest index, grains per panicle, and productive tillers for grain yield enhancement. The results revealed that the contribution of trait days to heading was the maximum for genetic divergence, followed by number of secondary branches, grain length, days to maturity, spikelets per panicle, and days to fifty percent flowering. Higher inter-cluster distance was observed between cluster V and cluster VII followed by cluster V and VII. Higher intra-cluster distance was observed between genotypes of cluster IV. Parents showing more divergence can be crossed, which could produce high heterotic hybrids.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Banumathy, S., Manimaran, R., Sheeba, A., Manivannan, N., Ramya, B., Kumar, D., & Ramasubramanian, G. V. (2010). Genetic diversity analysis of rice germplasm lines for yield attributing traits. *Electronic Journal of Plant Breeding*, 1(4), 500–504.
- Carcea, M. (2021). Value of wholegrain rice in a healthy human nutrition. *Agriculture (Switzerland)*, 11(8), 720.
- Guru, T., Padma, V., Reddy, D. V. V., Rao, P. R., Sanjeeva Rao, D., Ramesh, T., & Radhakrishna, K. V. (2017). Genetic diversity analysis for yield attributing traits in rice genotypes. *Research on Crops*, 18(2), 311–315.
- Kumar, B., Kour, A., & Laxami, J. (2017). Genetic diversity for agro-morphological traits in rice germplasm. *Electronic Journal of Plant Breeding*, 8(4), 1216–1222.
- Mohidem, N. A., Hashim, N., Shamsudin, R., & Che Man, H. (2022). Rice for food security: Revisiting its production, diversity, rice milling process and nutrient content. *Agriculture*, 12(6), 741.
- Nawaz, M. A. (2018). *Processing and quality of glutinous rice* (Master's thesis, The University of Queensland Australia).
- Rasane, P., Jha, A., Sabikhi, L., Kumar, A., & Unnikrishnan, V. S. (2015). Nutritional advantages of oats and opportunities for its processing as value added foods—a review. *Journal of Food Science and Technology*, 52, 662–675.
- Roy, S., Banerjee, A., Pattanayak, A., Roy, S. S., Rathi, R. S., Misra, A. K., & Bansal, K. C. (2014). Chakhao (delicious) rice landraces (*Oryza sativa* L.) of North-east India: Collection, conservation and characterization of genetic diversity. *Plant Genetic Resources*, 12(3), 264–272.
- Sen, S., Chakraborty, R., & Kalita, P. (2020). Rice—not just a staple food: A comprehensive review on its phytochemicals and therapeutic potential. *Trends in Food Science & Technology*, 97, 265–285.
- Sinha, A. K., & Mishra, P. K. (2013). Agro-morphological characterization and morphology based genetic diversity analysis of landraces of rice variety (*Oryza sativa* L.) of Bankura district of West Bengal. *International Journal of Current Research*, 5(10), 2764–2769.
- Uraipong, C. (2016). *Investigation into the biological functions of rice bran protein hydrolysates* (Doctoral dissertation, UNSW Sydney).

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://pr.sdiarticle5.com/review-history/134628>