



Agronomic Evaluation of Artificial Termite Mound Waste for Enhancing Rainfed Rice Production in Central-Western Côte d'Ivoire

YEO Lacina ^{a*}, TOURE Ambéyin ^a, KOFFI Antoine ^b
and SORO Dogniméton ^a

^a UFR Agroforesterie, Université Jean Lorougnon, Laboratoire Amélioration de la Production Agricole, BP 150 Daloa, Côte d'Ivoire.

^b Université Peleforo Gon Coulibaly, UFR des Sciences Biologiques, B. P. 1328, Korhogo, Côte d'Ivoire.

Authors' contributions

This work was carried out in collaboration among all authors. Author YL designed the study, performed the statistical analysis, wrote the protocol, and wrote the first draft of the manuscript. Authors TA and KA managed the analyses of the study. Author SD managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ijpss/2025/v37i95746>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://pr.sdiarticle5.com/review-history/144109>

Original Research Article

Received: 08/07/2025

Accepted: 16/09/2025

Published: 19/09/2025

ABSTRACT

Aims: To contribute to improving the productivity of rainfed rice by amending the soil with a waste product.

Study Design: The experiment was conducted using a randomized complete block design (RCBD) with three replicates, each comprising ten treatments.

*Corresponding author: E-mail: tchebelass@yahoo.fr;

Place and Duration of Study: The field trial was conducted in 2018–2019 in Daloa, Côte d'Ivoire.

Methodology: A sandy soil was amended with the waste product “Termisol.” Six doses of Termisol (29.1 t ha⁻¹ to 175 t ha⁻¹) and four controls (zero fertilizer, NPK fertilizer (0.15 t ha⁻¹), manure (7 t ha⁻¹), and natural termite mound soil (116.5 t ha⁻¹) were tested. Observations focused on leaf area, plant height, biomass production, and yield.

Results: The results showed that in the first cycle, the dose of 116.5 t ha⁻¹ produced the largest leaves (56.11±10.48 cm²), the highest plant biomass (21.32±7.75 g), while the dose of 175 t ha⁻¹ produced the tallest plants (67.43±1.95 cm) and the dose of 87.30 t ha⁻¹ produced the highest fresh yield (5.71±1.26 t ha⁻¹). The 175 t ha⁻¹ dose produced the widest leaves (48.24±1.95 cm²), the tallest plants (47.60 ±4.04 cm), the largest biomass (11.50 ± 1.60 g), and the highest yield (4.43 ± 0.15 t ha⁻¹) in the second cycle.

Conclusion: The use of the waste product “Termisol” as a soil amendment to improve soil fertility could be a viable method for reducing production costs and increasing rice yields. However, this study needs to be expanded to refine the economically viable application rates for widespread use.

Keywords: Amendment; waste product; “Termisol”; rice; Daloa; Côte d'Ivoire.

1. INTRODUCTION

Food crop farming in Africa is characterized by low productivity due to rapid soil nutrient depletion and the combined effects of poor agricultural land management, varying climatic conditions, and population growth (Bationo et al., 2004; Tumawu et al., 2025). In sub-Saharan Africa, agricultural residues are generally collected after harvest by farmers for domestic use or burned in bush fires to create new pastureland. These actions promote the irreversible export of soil nutrients, leading to a disruption in the renewal of soil organic matter. Furthermore, fallow land is becoming less and less common due to strong pressure on arable land caused by growing needs resulting from population growth (Thomas et al., 2011). Thus, the dominant role played by agriculture in the economy of Côte d'Ivoire forces farmers to resort to modern agricultural techniques such as varietal selection and the rational use of pesticides and chemical fertilizers to improve yields (Barrett & Sheahan, 2014). While chemical fertilizers have a rapid beneficial effect on crop productivity, justifying their use, their misuse leads to soil degradation through loss of organic matter and acidification. Furthermore, the cost of fertilizers makes them almost inaccessible to small farmers (Gala et al., 2007; Useni et al., 2012), while the risks of environmental pollution and water eutrophication pose a threat to biodiversity and endanger the lives of populations. As a result, alternative methods of soil fertility management are being sought.

In addition, various studies report on certain soil macroinvertebrates such as termites and

earthworms and the use of agricultural residues to improve soil properties (Guei et al., 2020). Thus, the work of Dosso et al. (2017) highlighted the contribution of termites to restoring the fertility of degraded soils by examining their contribution of nutrients through the on-site decomposition of residues from a few plant species, while the work of Soro et al. (2021) demonstrated the fertilizing capacity of waste products from artificial termite mounds and their potential use in rainfed rice cultivation. These results open up an opportunity to initiate the formation by termites of a waste product that retains the characteristics of the natural termite mound in a semi-controlled environment. The overall objective of this study is to contribute to improving rainfed rice productivity by amending the soil with the waste product “Termisol” from the digestion of agricultural residues by termites in an artificial termite mound. The specific objective is to evaluate the fertilizing capacity of the waste product in rainfed rice cultivation.

2. MATERIALS AND METHODS

2.1 Location of the Study Site

The work was carried out in Daloa and Campement Zongo. The first site was used for the production of the termite waste product “Termisol” and the second for agronomic testing (Fig. 1).

2.2 Material

2.2.1 Biological material

The biological material used in this study consists of rice plants grown from seeds of the




Fig. 1. Location of the production site and the evaluation test of the termite waste product "Termisol"

IDSA 10 variety. This is a rainfed rice variety with a 105-day cycle and an average yield of 2.5 t ha⁻¹. It also has good drought tolerance (CNRA, 2016).

2.2.2 Technical equipment

The technical equipment used includes an electronic scale for weighing, a carpenter's folding ruler for measuring the height of rice plants, a pair of scissors used for sampling plants, tracing paper for evaluating leaf area, an oven, and aluminum foil for drying samples.

2.2.3 Soil, fertilizers, and soil amendments

The termite waste product "Termisol" is evaluated for its fertilizing capacity in rainfed rice cultivation in the presence of three controls:

- Poultry manure (organic fertilizer) is characterized by nitrogen (N), phosphorus (P), potassium (K), calcium (Ca), and magnesium (Mg) contents of 1.40%, 0.45%, 1.72%, 2.44%, and 0.32%, respectively. As for trace elements, the copper (Cu), iron (Fe), manganese (Mn), and zinc (Zn) contents were 5.03 mg kg⁻¹, 10.11 mg kg⁻¹, 5.09 mg kg⁻¹, and 7.19 mg kg⁻¹, respectively.

- NPK (15-15-15) and urea (46% N);

Termite mound soil characterized by a silty sandy texture, low in organic matter (1.16%), slightly acidic with a CEC of 7.2 cmol kg⁻¹. It is a desaturated soil with a saturation rate (V) of 34.44% and a base sum equal to 2.48 cmol kg⁻¹.

Termisol was obtained through the degradation of corn residues and the rise of mineral constituents by termites in an artificial termite mound. It is characterized by a sandy loam texture, rich in organic matter (5.86%), slightly acidic with a CEC of 9.92 cmol kg⁻¹. It is a moderately desaturated soil with a saturation rate (V) of 74.14% and a base sum of 7.4 cmol kg⁻¹ (Soro et al., 2021).

2.3 Methods

2.3.1 Agronomic evaluation of the termite waste product "Termisol"

2.3.1.1 Experimental design

The experimental design is completely randomized blocks (CRB) with three replicates of ten (10) treatments each (Fig. 2). The treatments consist of six (6) doses of "Termisol" and four controls.




Fig. 2. Experimental design applied in the field

The doses of “Termisol” applied are calculated based on the mass of soil contained in one square meter at a working depth of 22 cm of the plot to be treated (Soro et al., 2021).

The doses of Termisol used were therefore:

- 29.1 t ha⁻¹, or 1% Termisol (T1);
- 58.30 t ha⁻¹, or 2% Termisol (T2);
- 87.30 t ha⁻¹, or 3% Termisol (T3);
- 116.50 t ha⁻¹, or 4% Termisol (T4);
- 145.50 t ha⁻¹, or 5% Termisol (T5);
- 175 t ha⁻¹, or 6% Termisol (T6).

The control treatments are:

- ZeroF: absolute control without fertilizer (T0);
- NPK: application of 0.15 t ha⁻¹ of NPK (15-15-15): mineral control (T7);
- TermNat: application of 116.50 t ha⁻¹ of natural termite mound soil (T8) taken from the trial site (Boga, 2007);
- Manure: application of chicken manure at a rate of 7 t ha⁻¹: organic control (T9).

The treatments were applied once as basal dressing in cycle 1. During the vegetative phase, top dressing with urea (46% N) at a rate of 0.07 t ha⁻¹ was applied 40 days after sowing (das) on all plots during both cycles.

The basic plot is rectangular, measuring 1.25 m long and 1 m wide.

2.3.1.2 Evaluation of leaf area

Leaf area was evaluated at 64 days after sowing (das) by reproducing 4 leaves per treatment on tracing paper. The traced image of the leaf was then cut out and its mass evaluated. This mass was used to estimate the leaf area, knowing the mass per unit area of the tracing paper. The leaf areas were then deduced by correspondence from the calculated masses (Lepengue et al., 2012). Leaf area is expressed in square centimeters (cm²).

2.3.1.3 Evaluation of plant height

Plant height was measured using a carpenter's folding ruler at 75 days after sowing (das). This measurement was taken from the base of the plant at the point where the panicle was attached to the panicle leaf of the most vigorous plant in the plot. Height was expressed in cm.

2.3.1.4 Determination of biomass and root size

Destructive sampling was carried out on two plants per treatment at 75 das. The selected plants were carefully pulled up, the roots were rinsed with water, dried with blotting paper, and their lengths measured with a folding carpenter's ruler. Each fresh plant was then weighed and its root cut at the collar. The fresh roots and fresh aerial parts were weighed separately and placed in an oven to dry at 70°C for 48 hours before dry

weighing. Lengths were expressed in cm and masses in g.

2.3.1.5 Evaluation of paddy yield

The rice is harvested by bunch at 105 das. The panicles from the same bunch are then weighed together. Three panicles per plot are sent to the laboratory to be dried in an oven at 65°C for 24 hours.

The dry paddy yield is calculated at 12% moisture content, using a conversion table to convert fresh yield to dry yield:

$$\text{Yield} = (\text{harvested mass (kg)}) / (\text{area (m}^2\text{)}) \times k,$$

$$\text{where } k = (100-H)/(100-h)$$

Yield = yield in kg m⁻²;

H = moisture content at harvest (%);

h = moisture content of dry grains (12%);

k = conversion factor from wet grain to dry grain.

2.3.2 Data processing

The data were processed using XLSTAT software. The normality of the sample distributions was verified using the Shapiro- Wilk test and the homogeneity of variances was verified using Levene's test. If the variable from which the sample originates follows a normal distribution and the variances are homogeneous, an ANOVA is performed; otherwise, the

nonparametric Kruskal-Wallis test is applied. The analysis of variance was supplemented by the Newman-Keuls test, which allowed the means to be ranked using the smallest significant difference (*P*values) at a 5% probability threshold. A principal component analysis (PCA) was then performed using R software.

3. RESULTS AND DISCUSSION

3.1 Effect of “Termisol” on the growth and Yield of Rainfed rice Depending on Treatment

3.1.1 Effect of “Termisol” dose on rice plant height

No significant differences were observed between treatments in terms of rice plant height (Haut), regardless of the crop cycle (Table 1). However, the average plant height varied overall between 31.33±2.47 cm (1% Termisol, cycle 2) and 67.43±1.95 cm (6% Termisol, cycle 1). Between these two extreme values, plants in the 3% and 5% Termisol treatments reached heights of 63.80±1.50 cm and 64.56±0.62 cm, respectively, in cycle 1. The plants in the ZeroF and TermNat treatments were slightly taller than those in the 1% Termisol treatment, reaching heights of 32.53±3.62 cm and 34.07±8.52 cm, respectively, in cycle 2. Overall, the plants were taller in cycle 1.

Table 1. Variation in the number of leaves, leaf area, and height of rice during the two cycles

Treatments	Growth parameters			
	Surfol (cm ²)		Haut (cm)	
	Cycle 1	Cycle 2	Cycle 1	Cycle 2
ZeroF (without fertilizer)	43.05±1.94 ^a	25±0.96 ^c	58.46±5.50 ^a	32.53±3.62 ^a
1%Termisol (29.1 t ha ⁻¹)	43.14±3.22 ^a	29.35±3.49 ^{bc}	61.03±1.57 ^a	31.33±2.47 ^a
2%Termisol (58.30 t ha ⁻¹)	50.74±17.76 ^a	35.09±3.07 ^{abc}	61.11±1.46 ^a	35.08±5.42 ^a
3%Termisol (87.30 t ha ⁻¹)	53.24±9.36 ^a	29.16±4.88 ^{bc}	63.80±1.50 ^a	39.57±10.39 ^a
4%Termisol (116.50 t ha ⁻¹)	56.11±10.48 ^a	33.98±11.62 ^{abc}	63.62±1.62 ^a	37.73±4.98 ^a
5%Termisol (145.50 t ha ⁻¹)	53.61±14.07 ^a	43.61±3.63 ^{ab}	64.56±0.62 ^a	38.49±4.77 ^a
6%Termisol (175 t ha ⁻¹)	54.16±9.05 ^a	48.24±4.78 ^a	67.43±1.95 ^a	47.60±4.04 ^a
NPK (0.15 t ha ⁻¹)	44.35±2.85 ^a	33.88±2.37 ^{abc}	62.70±5.20 ^a	38.57±4.48 ^a
TermNat (116.50 t ha ⁻¹)	46.85±10.35 ^a	24.81±0.32 ^c	59.05±6.77 ^a	34.07±8.52 ^a
Manure (7 t ha ⁻¹)	43.98±8.95 ^a	30.46±11.60 ^{bc}	62.44±5.99 ^a	43.53±2.80 ^a
CV	0.1949	0.2603	0.0652	0.1749
<i>P</i>	.6217	.0017	.2566	.0559
Effect	ns	**	ns	ns

Averages followed by the same letter in the same column are not significantly different at the 5% threshold for the Newman-Keuls test; **highly significant difference *P* = .001; ns: Non-significant difference *P* > .05; CV: Coefficient of variation; TermNat: Natural termite mound; ZeroF: Control without fertilizer; Surfol: Leaf area; Height: Height 75 days after sowing (das).

3.1.2 Effect of the dose of “Termisol” on leaf area

The leaf surface area (Surfol) of rice plants varied significantly in cycle 2 ($P = .001$) between treatments (Table 1). It ranged from $24.81 \pm 0.32 \text{ cm}^2$ in the TermNat treatment in cycle 2 to $56.11 \pm 10.48 \text{ cm}^2$ in the 4% Termisol treatment (cycle 1). The 5% Termisol and 6% Termisol treatments had leaf areas of $53.61 \pm 14.07 \text{ cm}^2$ and $54.16 \pm 9.05 \text{ cm}^2$ respectively in cycle 1, below that of the 4% Termisol treatment in the same cycle. The ZeroF and 3% Termisol treatments had leaf areas slightly larger than that of the TermNat treatment, which was the smallest in cycle 2.

3.1.3 Effect of dose on dry biomass production in rice

The fertilizer dose had a highly significant effect ($P = .001$) on dry matter production in cycle 2 (Table 2).

Above-ground biomass (MsTF) was higher in the 6% Termisol ($18.06 \pm 5.34 \text{ g}$), 2% Termisol ($19.45 \pm 10.78 \text{ g}$), and 4% Termisol ($19.84 \pm 7.26 \text{ g}$) treatments in cycle 1. The lowest amounts of dry matter were recorded in the ZeroF, 1% Termisol, and TermNat treatments, with values of $2.17 \pm 0.43 \text{ g}$, $4.19 \pm 0.96 \text{ g}$, and $5.15 \pm 0.98 \text{ g}$, respectively, in cycle 2. Above-ground dry matter was more abundant in cycle 1.

Root biomass (MsRac) varied between $0.44 \pm 0.03 \text{ g}$ (ZeroF, cycle 2) and $1.43 \pm 0.46 \text{ g}$ (4% Termisol, cycle 1). Between these two

values, the 6% Termisol (cycle 2) and 2% Termisol (cycle 1) treatments achieved dry matter quantities of $1.37 \pm 0.24 \text{ g}$ and $1.41 \pm 0.59 \text{ g}$. The TermNat and 1% Termisol treatments showed low biomass values of $0.65 \pm 0.15 \text{ g}$ and $0.68 \pm 0.13 \text{ g}$, respectively, in cycle 2, which were higher than those of the ZeroF treatment in the same cycle, but with a highly significant difference between these three treatments. The total biomass of the plants (MsPE) evolved similarly to the biomass of the aerial part (MsTF). With the exception of the 6% Termisol and NPK treatments, the proportion of root dry matter (MsRac) was higher in cycle 1.

3.1.4 Effect of the dose of Termisol on 1000-grain weight, grain moisture, and yield

The addition of Termisol had a significant effect ($P < .001$) on 1000-grain weight (PMG), relative grain moisture at harvest (HR(%)), and fresh yield at harvest (Rendfrais) in the second cycle (Table 3). The PMG varied overall between $17.84 \pm 1.58 \text{ g}$ (ZeroF, cycle 2) and $26.88 \pm 3.19 \text{ g}$ (4% Termisol, cycle 2). Thus, the highest PMGs were obtained with the 4%, 5%, and 2% Termisol treatments, with $26.88 \pm 3.19 \text{ g}$ (cycle 2); $26.22 \pm 5.31 \text{ g}$ (cycle 2) and $25.74 \pm 0.76 \text{ g}$ (cycle 1), respectively. The moisture content of the grain at harvest varied significantly ($P < .001$) between treatments in cycle 2 only. The ZeroF treatment (cycle 2) had the highest grain moisture content at harvest, $23.45 \pm 0.92\%$, and the 4% Termisol treatment had the lowest moisture content in cycle 1, $15.31 \pm 1.04\%$.

Table 2. Variation in rice biomass during two crop cycles according to fertilizer dose

Treatments	Growth parameters					
	MsPE (g)		MsTF (g)		MsRac (g)	
	Cycle 1	Cycle 2	Cycle 1	Cycle 2	Cycle 1	Cycle 2
ZeroF	14.69 ± 2.25^a	2.63 ± 0.41^b	13.45 ± 2.17^a	2.17 ± 0.43^e	1.18 ± 0.08^a	0.44 ± 0.03^b
1%Termisol	14.7 ± 2.90^a	4.89 ± 1.08^{ab}	13.39 ± 2.83^a	4.19 ± 0.96^{de}	1.24 ± 0.07^a	0.68 ± 0.13^b
2%Termisol	20.94 ± 11.40^a	6.18 ± 1.64^{ab}	19.45 ± 10.78^a	5.44 ± 1.52^{bcde}	1.41 ± 0.59^a	0.75 ± 0.12^b
3%Termisol	16.02 ± 0.83^a	9.01 ± 2.52^{ab}	14.72 ± 0.78^a	8.06 ± 2.33^{abc}	1.25 ± 0.08^a	0.94 ± 0.19^{ab}
4%Termisol	21.32 ± 7.75^a	8.97 ± 3.12^{ab}	19.84 ± 7.26^a	7.99 ± 2.95^{abc}	1.43 ± 0.46^a	0.99 ± 0.19^{ab}
5%Termisol	15.55 ± 2.98^a	9.89 ± 3.21^a	14.33 ± 2.80^a	8.91 ± 3.07^{ab}	1.18 ± 0.22^a	0.98 ± 0.15^{ab}
6%Termisol	19.34 ± 5.45^a	11.50 ± 1.60^a	18.06 ± 5.34^a	10.15 ± 1.47^a	1.26 ± 0.18^a	1.37 ± 0.24^a
NPK	14.52 ± 0.87^a	7.86 ± 2.41^{ab}	13.62 ± 0.66^a	6.92 ± 2.21^{abcd}	0.88 ± 0.22^a	0.92 ± 0.26^{ab}
TermNat	14.07 ± 1.39^a	5.81 ± 1.16^{ab}	13.14 ± 1.33^a	5.15 ± 0.98^{cde}	0.95 ± 0.20^a	0.65 ± 0.15^b
Manure	14.89 ± 2.14^a	7.39 ± 3.84^{ab}	13.80 ± 2.03^a	6.55 ± 3.51^{abcd}	1.05 ± 0.21^a	0.84 ± 0.32^b
CV	0.2937	0.4224	0.3004	0.4385	0.2430	0.3336
P	.5098	.0069	.5144	.0084	.3553	.0017
Effect	ns	**	ns	**	ns	**

Averages followed by the same letter in the same column are not significantly different at the 5% threshold for the Newman-Keuls test; * significant difference $P = .05$; **: highly significant difference $P = .001$; ns: non-significant difference $P > .05$; CV: Coefficient of variation;

Table 3. Average change in production parameters during the two cycles

Treatments	Production parameters					
	PMG (g)		HR(%)		Rendfrais (t ha ⁻¹)	
	Cycle 1	Cycle 2	Cycle 1	Cycle 2	Cycle 1	Cycle 2
ZéroF	24.29±2.92 ^a	17.84±1.58 ^c	16.39±0.41 ^a	23.45±0.92 ^a	3.77±0.41 ^a	2.29±0.45 ^f
1%Termisol	24.76±0.72 ^a	21.05±3.10 ^c	16.05±0.68 ^a	22.05±1.28 ^a	5.16±0.14 ^a	2.96±0.41 ^{def}
2%Termisol	25.74±0.76 ^a	19.08±1.39 ^c	15.92±0.59 ^a	21.8±1.37 ^a	5.64±1.42 ^a	2.95±0.54 ^{def}
3%Termisol	25.57±2.64 ^a	21.48±0.51b ^c	16.18±1.01 ^a	22.33±0.49 ^a	5.71±1.26 ^a	3.15±0.63 ^{cde}
4%Termisol	24.60±2.08 ^a	26.88±3.19 ^a	15.31±1.04 ^a	18.11±1.84 ^b	4.76±0.38 ^a	4.08±0.54 ^{ab}
5%Termisol	24.28±3.54 ^a	26.22±5.31 ^a	15.66±1.31 ^a	19.28±1.80 ^b	4.67±0.29 ^a	3.96±0.45 ^{abc}
6%Termisol	22.94±2.45 ^a	25.56±2.09 ^{ab}	16.48±0.53 ^a	19.28±1.08 ^b	4.86±0.93 ^a	4.43±0.15 ^a
NPK	25.85±0.73 ^a	19.71±1.16 ^c	16.87±1.30 ^a	23.28±1.27 ^a	4.07±0.63 ^a	3.48±0.29 ^{bcd}
TermNat	24.32±1.07 ^a	18.33±2.27 ^c	16.98±0.24 ^a	22.98±1.56 ^a	5.08±0.62 ^a	2.53±0.70 ^{ef}
Manure	25.24±2.22 ^a	20.70±0.95 ^c	15.33±0.77 ^a	22.18±0.88 ^a	3.90±0.45 ^a	3.97±0.38 ^{abc}
CV	0.0785	0.1748	0.0555	0.0971	0.1917	23.21
P	.8542	.0006	.2744	.0004	.0696	.0008
Effect	ns	***	ns	***	ns	***

Averages followed by the same letter in the same column are not significantly different at the 5% threshold for the Newman-Keuls test; ** $P < 0.0001$




Fig. 3. Yield curves for paddy rice in $t\ ha^{-1}$ as a function of the dose in cycles 1 and 2




Fig. 4. PCA combining growth and yield parameters for the IDSA10 rice variety
Dens: plant density; LgRac: root length; TalTot: total number of tillers per plant; Talfert: number of fertile tillers per plant.

The fresh yield at harvest of rice was strongly influenced ($P < .001$) by treatment during cycle 2 only. The highest fresh yields were obtained with the 3%, 2%, and 1% Termisol treatments, with $5.71 \pm 1.26\ t\ ha^{-1}$; $5.64 \pm 1.42\ t\ ha^{-1}$; and 5.16 ± 0.14

$t\ ha^{-1}$ respectively in cycle 1. The ZeroF, TermNat and 2% Termisol treatments had the lowest fresh yields, with values of $2.29 \pm 0.45\ t\ ha^{-1}$, $2.53 \pm 0.70\ t\ ha^{-1}$ and $2.95 \pm 0.54\ t\ ha^{-1}$, respectively, in cycle 2.

Paddy yields varied according to the dose of Termisol in cycle 1 and cycle 2 (Fig. 3). In cycle 1, fresh yields increased gradually with the doses of Termisol, from $3.77 \pm 0.36 \text{ t ha}^{-1}$ (ZeroF) to $5.71 \pm 1.26 \text{ t ha}^{-1}$ (3% Termisol), while dry yields increased from $3.58 \pm 0.32 \text{ t ha}^{-1}$ (ZeroF) to $5.44 \pm 1.18 \text{ t ha}^{-1}$ (3% Termisol). Beyond these values, the yield decreased despite the increase in the Termisol dose. For the second cycle carried out in 2019, yields were lower than those obtained in 2018. For this first cycle, the best yield was obtained on the plot treated with 3% Termisol.

Fresh and dry yields in cycle 2 fluctuated, with the best yields obtained on plots treated with 4% Termisol or more for both fresh and dry yields.

3.2 Relationship Between Growth and Production Parameters in Rice

The projection in factorial plane 1-2 associates all growth and production parameters with axis 1 (Fig. 4 A).

The factorial map allows two productivity groups to be defined that are opposed on axis 1 (Fig. 4 B). Group 1, characterized by low production levels, includes the treatments Termitière naturelle (TermNat), ZéroF, 1%Termisol, NPK, and Fiente, forming the Fiente - NPK group. Group 2, comprising treatments based on Termisol at doses of 2 to 6%, constitutes the Termisol group, characterized by better vegetative growth and higher rice productivity.

3.3 Discussion

The results of the study showed that soil amendment with Termisol improved rice growth and production parameters. Termisol promoted good height growth, larger leaf areas, and greater plant biomass. It also promoted good yields with reduced grain moisture at harvest. Thus, despite a slightly lower thousand-grain weight compared to NPK fertilization, the waste products from artificial termite mounds achieved the best yields (Kpangba et al., 2020).

The effect of Termisol on the agronomic parameters of rice demonstrates its good fertility potential. Plots treated with Termisol have a structural advantage in that they remain within the same pH range favorable to rice (Zingore et al., 2014). In addition, the organic matter in Termisol and the presence of clay + silt contributed to the development of physical and

chemical conditions favorable to good soil growth and productivity (Drouet, 2010). Organic matter both nourishes plants by releasing adsorbed mineral elements and stores these elements. It forms the basis of soil fertility and plays a role in soil structure (Some et al., 2015). These results confirm those of Boga (2007) and Soro et al. (2021), who observed improved growth parameters for maize and rice in the former case and for rainfed rice in the latter. Mokossesse et al. (2009) showed that termite mound soil significantly influenced the growth of *Sorghum sp* and *Crotalaria ochroleuca* in pots, while Kpangba et al. (2020) observed that termite mound soil significantly influenced the growth parameters of black nightshade. Lepage and Duponnois (2006) also showed that *Cubitermes* termite powder used in gardening improves tomato fruit production by 150 to 300%. However, according to these authors, this influence of *Cubitermes* termite mound soil is due to the high nitrogen-rich fulvic acid content of the nest.

The results of this study show that *Macrotermes* termite mound soil, a natural termite mound at the test site, has little influence on the agronomic parameters of rice. The results obtained are similar to the observations made by Boga (2007) on corn plants. This low influence is thought to be due to the low colloidal content in natural termite mound soil. Indeed, termite mound soil is highly sandy and lacks chemical fertility potential. However, according to Lin et al., (2024), in addition to this low mineral content, the phosphorus contained in *Macrotermes* termite mound material is difficult to mineralize and therefore unavailable to plants. The results obtained contradict those of Beugré et al. (2020), who showed that *Macrotermes bellicosus* termite mound soil was rich in minerals and organic matter, but linked its low influence on the agronomic parameters of cowpea to its high acidity.

The addition of 4% and 6% Termisol had the most significant effects on biomass and rice plant height. The large amount of biomass and large plant size obtained with these treatments compared to the other treatments could be explained by the good organomineral level of these doses of Termisol. Indeed, these doses improved the nitrogen and phosphorus content of the soil. These results corroborate those of Kpangba et al. (2020), who obtained significant effects on the vegetative growth and biomass yield of black nightshade (*Solanum nigrum L.*)

plants with the addition of 2 kg, 4 kg, and 6 kg of termite mound soil. Rice plants require significant amounts of nutrients, particularly nitrogen, potassium, and phosphorus, as well as calcium and magnesium (Du et al., 2022; Rakotoson et al. 2023). The content of these elements in the soil determines the growth and productivity of rice, which explains why rice grows well with these treatments. Wang et al. (2024) and Akhtar et al., (2024) reported that the mineral nitrogen absorbed by the plant was used for amino acid synthesis and played an important role in the growth and development of the vegetative system. Termisol probably improved the effectiveness of the urea applied due to its contribution to the texture, organic matter, and consequently the structure of the initially sandy soil. In fact, the 0.07 t ha⁻¹ dose applied to the soil during production would have constituted the mineral supplement to Termisol for sustainable rice production on this type of sandy soil.

From cycle 1 to cycle 2, a gradual decrease in rice plant height and yield was observed in almost all treatments. This decrease could be explained by a gradual depletion of minerals in the soil, due to uptake by the rice plants and leaching by rainwater during each growing season. Termisol therefore acts as a slow-release fertilizer, as most of the nitrogen it contains is supplied in organic form, and the mineral forms that can be used by plants are released gradually (Mokossesse et al., 2012).

The rice was harvested at 105 das with moisture levels ranging from 15.31% to 16.98% in cycle 1 and from 18.11% to 23.45% in cycle 2. In all cases, rice harvested from plots treated with Termisol had the lowest moisture content and higher yields. The application of Termisol therefore provided conditions for earlier harvesting compared to other treatments.

The paddy yield in cycle 2 is lower than that in cycle 1. Furthermore, only the yield in cycle 1 varied according to the law of less-than-proportional yield supplements, in line with the work of Soro et al. (2021). This response of rice to increasing doses of Termisol confirms its fertility potential. Termite activity in the traps over a period of three to six days therefore produced a substance with fertility potential that can be used in food crop agriculture. Furthermore, the variation in paddy yield in cycle 2 is not related to the yield in cycle 1. The differentiated effect between the yields of cycles 1 and 2 for all

treatments therefore reflects a different and more significant after-effect of Termisol.

4. CONCLUSION

The results of the effect of "Termisol" on the agronomic parameters of rice show that different doses of "Termisol" had a positive influence on the growth and productivity parameters of rice. Amendments of 4% and 6% Termisol resulted in better growth of rice plants and good yields.

All these results show that Termisol could be a sustainable solution for improving soil fertility.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that they have no known competing financial interests or non-financial interests or personal relationships that could have appeared to influence the work reported in this paper.

REFERENCES

- Akhtar, K., Ain, N., Prasad, P. V. V., Naz, M., Aslam, M. M., Djalovic, I., et al. (2024). Physiological, molecular, and environmental insights into plant nitrogen uptake, and metabolism under abiotic stresses. *The Plant Genome*, 17(2). <https://doi.org/10.1002/tpg2.20461>
- Barrett, C. B., & Sheahan, M. B. (2014). Understanding the agricultural input landscape in Sub-Saharan Africa: Recent plot, household, and community-level evidence. *LSMS|Policy Research Working Paper|No. WPS 7014*. Washington, DC: World Bank Group. <http://documents.worldbank.org/curated/en/165341468002108906>
- Bationo, A., Nandwa, S. M., Kimetu, J. M., Kinyangi, J. M., Bado, B. V., Lompo, F., et al. (2004). Sustainable intensification of crop-livestock systems through manure management in eastern and western Africa: Lessons learned and emerging

- research opportunities. *Sustainable Crop-Livestock Production in West Africa*, 173–198.
- Beugré, M. M., Ayolié, K., Yapo, S. E.-S., Silué, T. A., & Kouadio, Y. J. (2020). Soil fertilization effect of *Macrotermes bellicosus* on the vegetative growth of two cowpea (*Vigna unguiculata* (L.) Walp) accessions in Daloa, Côte d'Ivoire. *Afrique Science*, 16(6), 38–50.
- Boga, P. J. (2007). Experimental study of the impact of termite mound materials on growth, maize and rice yield, and soil fertility in South Sudanese savannahs, Boro-Borotou, Côte d'Ivoire. Doctoral thesis, University of Cocody, Côte d'Ivoire.
- Dosso, K., Koné, F., Kra, K. D., & Konaté, S. (2017). Decomposition of plant residue by termites and contribution of organic matter to the soil in the Lamto region, Côte d'Ivoire. *Journal of Animal & Plant Sciences*, 33(3), 5320–5331.
- Drouet, T. (2010). *Pedology BING-F-302*, 137.
- Du, M., Zhang, W., Gao, J., Liu, M., Zhou, Y., He, D., et al. (2022). Improvement of root characteristics due to nitrogen, phosphorus, and potassium interactions increases rice (*Oryza sativa* L.) yield and nitrogen use efficiency. *Agronomy*, 12(1), 23. <https://doi.org/10.3390/agronomy12010023>
- Gala, J. T. B., Camara, M., Assa, A., & Keli, J. Z. (2007). Problems of mineral fertilizer use in rice production areas: The case of central-western Côte d'Ivoire. *Agronomie Africaine*, 19(2), 173–185.
- Guei, M. A., Zro Bi, G. F., Bakayoko, S., & Ta Bi, D. F. (2020). Effects of vermicompost based on household waste combined with rice husks on the agronomic parameters of maize (*Zea mays* L.). *Journal of Applied Biosciences*, 154, 15862–15870.
- Kpangba, K. M. P. T., Mpika, J., Makoundou, A., Bitá, A. M., & Attibayéba. (2020). Effect of termite mound soil on the growth of three cultivars of black nightshade (*Solanum nigrum* L.: Solanaceae) cultivated in Brazzaville (Congo). *International Journal of Biological and Chemical Sciences*, 14(1), 192–203.
- Lepage, M., & Duponois, R. (2006). Termite mound powder for gardens. *Cahiers Agricultures*, 15(5), 432–443.
- Lepengue, A. N., Ebang, E. G., Ikabanga, U. D., Mouaragadja, I., Bourobou, H., Séverin Ake, S., et al. (2012). Influence of *Capsicum frutescens* L. (Solanaceae) on the germination and growth of spinach (*Spicia oleracea*, Chenopodiaceae) in Gabon. *International Journal of Biological and Chemical Sciences*, 6(1), 298–306.
- Lin, F., Jin, Y., Liu, C., Ren, Y., Jiang, C., Feng, D., et al. (2024). Termite mounds affect soil aggregate stability and aggregate-associated phosphorus forms in a tropical rubber plantation. *Plant and Soil*, 498, 93–109. <https://doi.org/10.1007/s11104-023-05880-4>
- Mokossesse, A. J., Josens, G., Mboukoulida, J., & Ledent, F. J. (2012). Effect of field application of *Cubitermes* (Isoptera: Termitidae) mound soil on growth and yield of maize in Central Africa Republic. *Agronomie Africaine*, 24(3), 241–252.
- Mokossesse, J. A., Josens, G., & Lepage, M. (2009). Growth in pots of four plant species on substrates enriched with *Cubitermes* termite mound soil. *Tropicultura*, 27(3), 168–173.
- National Center for Agronomic Research (CNRA). (2016). *Directory of improved varieties of food crops* (2016 ed., 53 p.).
- Rakotoson, T., Senthilkumar, K., Johnson, J. M., Ibrahim, A., Kihara, J., Sila, A., et al. (2023). Estimating nutrient concentrations and uptake in rice grain in sub-Saharan Africa using linear mixed-effects regression. *Field Crops Research*, 299, 108987. <https://doi.org/10.1016/j.fcr.2023.108987>
- Some, D., Hien, E., Assigbetse, K., Drevon, J. J., & Masse, D. (2015). Dynamics of carbon and nitrogen compartments in the soil of cowpea and sorghum under the zai system in the North Sudanian zone of Burkina Faso. *International Journal of Biological and Chemical Sciences*, 9(2), 954–969.
- Soro, D., Yeo, L., Konate, Z., Guei, A. M., & Zro-Bi, F. G. (2021). When termites' waste products highlight a fundamental law of fertilization under rainfed rice cultivation in West Cote d'Ivoire. *European Journal of Agriculture and Food Sciences*, 3(1), 53–60.
- Thomas, G., Michael, J., Attanda, M. I., Heiko, P., & Claudia, H. (2011). Future productivity of fallow systems in Sub-Saharan Africa: Is the effect of demographic pressure and fallow reduction more significant than climate change? *Agricultural and Forest*

- Meteorology*, 151(8), 1120–1130.
<https://doi.org/10.1016/j.agrformet.2011.03.015>
- Tumawu, A. F., Kafu, G. Y., Kangah, A., & Tumawu, M. M. M. (2025). Combating land degradation for sustainable smallholder agriculture and food sovereignty in Ghana's Volta Region. *Frontiers in Sustainable Food Systems*, 9, 1455515.
<https://doi.org/10.3389/fsufs.2025.1455515>
- Useni, S., Baboy, L., Nyembo, K., & Mpundu, M. (2012). Effects of combined biowaste and inorganic fertilizer inputs on the yield of three *Zea mays* L. varieties grown in the Lubumbashi region. *Journal of Applied Biosciences*, 54, 3935–3943.
- Wang, Q., Li, S., Li, J., & Huang, D. (2024). The utilization and roles of nitrogen in plants. *Forests*, 15(7), 1191.
<https://doi.org/10.3390/f15071191>
- Zingore, S., Wairegi, L., & Ndiaye, K. M. (2014). *Guide for the management of rice cropping systems*. African Soil Health Consortium, Nairobi.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://pr.sdiarticle5.com/review-history/144109>