



Colorimetric Quantification of CO₂ in a Modified Alkali-Trap Soil Respiration Assay

**Durgesh Agase^{a*}, Sheefa Khan^a, Madhvi Bisen^a
and Harsh Tiwari^a**

^a PMCoE, Govt. J.S.T.P.G. College, Balaghat, M.P., India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ijpss/2025/v37i115865>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://pr.sdiarticle5.com/review-history/148908>

Short Research Article

Received: 29/09/2025

Accepted: 01/12/2025

Published: 05/12/2025

ABSTRACT

Microbial activity in soil is an important characteristic of healthy soil. A key indicator of microbial activity is soil respiration. Traditional alkali-trap assays use hydrochloric acid to titrate sodium hydroxide that has not reacted to quantify CO₂ evolution, but determining the titrimetric endpoint is subjective, involves colour fading, reagent instability, and operator variability. The present study reports a modified soil respiration assay using colorimetric quantification of pH reduction as sodium carbonate forms. The method is phenolphthalein-based absorbance reading at 552 nm with a colorimeter. The alteration of the assay greatly improved precision, eliminated blank samples, and reduced human error from visual titration. This colorimetric method is easier, inexpensive, and provides reliable quantification of soil microbial respiration.

*Corresponding author: E-mail: sbt.durgesh@gmail.com;

Keywords: Soil respiration; colorimetric analysis; CO₂ evolution; alkali trap; microbial activity.

1. INTRODUCTION

Soil respiration, microbial and root-mediated release of carbon dioxide (CO₂), is a critical ecosystem process involving the cycling of carbon and has been widely utilized as an indicator of soil health (Gougoulas *et al.*, 2014). Currently available laboratory methods, the alkali-trap method enjoys popularity because it is simple and inexpensive (Irving *et al.*, 2024). The procedure involves the reaction of CO₂ released from incubated soils with NaOH, and the quantification of unreacted alkali to estimate CO₂ evolution. Traditionally, this is done by acid-base titration, which, despite effectiveness, has major limitations: subjective colour endpoint detection, time-consuming titrations, errors as a result of either slow or partial colour change, inefficiency during large-scale sample processing, and atmospheric CO₂ contamination during reagent handling (Bekku *et al.*, 1997; Haney, 2008; Doran *et al.*, 2012). To overcome these deficiencies, the following colorimetric modification is proposed, employing phenolphthalein as an indicator: instead of adding an indicator and titrating, the absorbance of the NaOH-phenolphthalein mixture is measured by a colorimeter at 552 nm (Kostjukov *et al.*, 2025). The reduction in absorbance following incubation with soil varies directly with the quantity of NaOH that has reacted with CO₂. As approximately 120 Pg C is released each year, it is the second-largest carbon flux between the biosphere and the atmosphere after global gross primary productivity (Bond-Lamberty & Thomson, 2010; Hashimoto *et al.*, 2015). Global soils hold almost twice the amount of carbon than the atmosphere, and so even minor changes to soil respiration can have cascading effects for climate feedbacks and ecosystem carbon balance (Crowther *et al.*, 2016). The flow of carbon, via soil respiration, is driven by rhizosphere respiration, microbial decomposition, and soil physiochemical factors temperature, moisture, and substrate availability (Schimel & Schaeffer, 2012). Therefore, accurately measuring soil respiration is imperative for evaluating soil quality, carbon cycling, and assessing the effects of land-use change or climate change on soil processes (Doran & Zeiss, 2000; Haney *et al.*, 2008). Of the laboratory methods available, the alkali-trap method is favoured for its simple and inexpensive measurement (Irving *et al.*, 2024). The procedure consists of the reaction of CO₂ released from incubated soils with NaOH and


quantifying unreacted alkali to calculate CO₂ evolution (Agase & Tiwari, 2025). Traditionally, this is accomplished with acid-base titration, which, although effective, but with significant limitations: endpoint detection by subjective colour, time-consuming titrations, imprecisions for slow or partial colour change, bad apportioning for larger-scale sample processing, and risk of atmosphere CO₂ contamination of reagents (Bekku *et al.*, 1997; Haney, 2008; Doran *et al.*, 2012). To solve these issues the following colorimetric adaptation can be suggested. This is done by using phenolphthalein as an indicator: the absorbance of the NaOH-phenolphthalein solution is measured by a colorimeter at 552 nm, rather than adding an indicator and then titrating (Kostjukov *et al.*, 2025).

1.1 Objective

To develop and validate modified colorimetric alkali-trap soil respiration assay for accurate, rapid, and reproducible quantification of CO₂

1.2 Principle

During incubation, soil-respired CO₂ reacts with NaOH:



A known amount of NaOH is placed in a sealed container with the soil sample. After incubation, the remaining unreacted NaOH is mixed with phenolphthalein, producing a pink colour in alkaline conditions. The absorbance is measured at 552 nm, and the concentration of remaining NaOH is determined using a calibration curve. A decreased absorbance indicates higher CO₂ absorption.

2. MATERIALS AND METHODS

The materials required for the colorimetric CO₂ evolution assay include fresh soil samples sieved to 2 mm, airtight incubation jars equipped with silicone gaskets, and CO₂-free 0.1 M NaOH solution for trapping evolved CO₂. Phenolphthalein indicator (1%) solution prepared in ethanol is used for colorimetric detection. Photocolorimeter (200nm-700nm), pH meter, analytical balance, micro-pipettes (10-100µL), pipettes (10mL), and incubator are also required for the CO₂ evolution assay.

2.1 Preparation of Calibration Curve

NaOH standards (0.01–0.10 M) are prepared, followed by adding 5 µL of phenolphthalein to each standard solution to obtain the calibration curve. The absorbance of these NaOH-phenolphthalein solutions was determined at 552 nm by using a colorimeter. The absorbance values (Y-axis) are plotted against the corresponding NaOH concentrations (X-axis) to derive a calibration curve and obtain a regression equation (e.g., $Y = aX + b$) for the description of the relationship for CO₂ calculation. For correlation analysis pH of the standards were also measured.

2.2 Soil Incubation Procedure

For the soil incubation procedure, 20.00 g of soil (Three Samples: Sand, Black Soil, and Vermicompost) were weighed into each airtight jar. Afterward, vials containing 20 mL of 0.1 M NaOH were placed inside to trap the CO₂ evolved during incubation. The jars were sealed immediately to prevent atmospheric CO₂ from

reacting with the alkali, and the setup is incubated in the dark at 25 °C for 24–72 hours. Appropriate controls are included, consisting of a reagent blank containing only NaOH without jar and soil, an incubation blank with the jar and NaOH with autoclaved soil, and an incubation blank with the jar and NaOH with no soil, to account for any non-soil-related CO₂ absorption.

2.3 Colorimetric Quantification of CO₂

After incubation, the NaOH trap is carefully removed, and the solution is quantitatively transferred into a volumetric flask, after which phenolphthalein is added until a pink colour develops. The absorbance of this solution is then measured at 552 nm using a colorimeter, and the concentration of the remaining NaOH is determined from the previously prepared calibration curve. The difference between the initial and remaining NaOH represents the amount of NaOH consumed, which is subsequently used to calculate the corresponding quantity of CO₂ evolved from the soil.




Fig. 1. Experimental setup




Fig. 2. NaOH trap for different soils

2.4 Calculations

After measuring the absorbance of the phenolphthalein–NaOH solution from each incubated sample at 552 nm, the remaining NaOH concentration is obtained from the calibration curve (the linear regression relating absorbance to NaOH concentration):

$$\text{Absorbance} = a \cdot [\text{NaOH}] + b$$

Here, a is the slope of the calibration line and represents how many absorbance units change per unit change in NaOH concentration (for example, absorbance units per mol L^{-1}), while b is the y-intercept (the absorbance value when NaOH concentration = 0) and accounts for any baseline absorbance of the reagent/solvent or instrument offset. Using the regression equation, you rearrange to find the NaOH concentration remaining in the trap:

$$[\text{NaOH}]_{\text{remaining}} = (\text{Absorbance sample (OD)} - b) / a$$

Multiply this concentration by the trap volume to get moles of NaOH remaining.

$$\text{Moles of } [\text{NaOH}]_{\text{Remaining}} = [\text{NaOH}]_{\text{Remaining}} \times \text{Trap Volume}$$

$$\text{Moles of } [\text{NaOH}]_{\text{Initial}} = [\text{NaOH}]_{\text{Initial}} \times \text{Trap Volume}$$

$$\text{Moles of NaOH that reacted with CO}_2 = \text{Moles of } [\text{NaOH}]_{\text{Initial}} - \text{Moles of } [\text{NaOH}]_{\text{Remaining}}$$

Because two moles of NaOH consume one mole of CO_2 , the moles of CO_2 evolved are

$$\text{CO}_2 = \text{reacted} / 2$$

Convert to mass by multiplying by the molar mass of CO_2 (44.01 g mol^{-1}).

3. RESULTS

In the present study, the correlation analysis revealed a strong positive relationship between NaOH concentration and pH ($r = 0.991$). Similarly, a strong positive correlation was also observed between NaOH concentration and optical density (OD) ($r = 0.999$). Linear regression analysis further indicated that the relationship between NaOH concentration and pH followed the equation $\text{pH} = 15.012(\text{NaOH}) + 12.097$.

On the contrary, the regression between NaOH concentration and OD, described by the equation $\text{OD} = 5.476 (\text{NaOH}) + 0.202$, provided a strong and statistically significant relationship ($p = 0.00095$). Collectively, the data indicate that the NaOH concentration has a significantly stronger impact on OD. $[\text{NaOH}]_{\text{remaining}} = (\text{OD} - 0.202) / 5.476$.




Fig. 3. Plot between pH and NaOH Con




Fig. 4. Plot between OD and NaOH Con





Fig. 5. 3D Scatter (pH, OD, and NaOH Con.)

Table 1. Different soil types and CO₂ evolved

Soil Type	OD	pH	CO ₂ Evolved moles/gram
Reagent Blank	0.856	13.21	19.386405
Autoclaved Soil	0.666	13.01	20.15658
NaOH without Soil	0.925	13.11	19.10034
Sand	0.522	12.42	20.72871
Black Soil	0.401	11.86	21.21282
Vermicompost Soil	0.374	11.65	21.322845
Organic Fertilizer Soil	0.189	11.54	21.674925


Fig. 6. Soil respiration in different soil types

Soil respiration measured using the alkali (NaOH) trap method exhibited different variations among different soil types. The reagent blank (NaOH only) recorded 19.3 moles/g/day CO₂, estimated due to the reaction with environmental CO₂. The autoclaved soil blank showed slightly higher values 20.15 moles/g/day, attributed to abiotic CO₂ release from minerals or trapped gases. The incubation blank without soil showed 19.10 moles/g/day CO₂, likely due to minor atmospheric contamination or jar headspace CO₂. Among the soil samples, Organic Fertilizer Soil exhibited the highest respiration rate, 21.67 moles/g/day CO₂, indicating high microbial activity and rich organic carbon content.

Vermicompost soil and Black soil showed 21.32 and 21.21 moles/g/day CO₂, respectively, consistent with its balanced organic matter and microbial biomass. Sand presented the lowest biological activity 20.72871 mole/g/day, CO₂, due to minimal organic carbon and poor microbial populations. Overall, the soil respiration trend is shown in the Fig. 5.

4. DISCUSSION

Although traditional chemical titration techniques have good use, there are more sensitive and higher-throughput automated instrumental methods, such as Infrared Gas Analysers (IRGA)

and Solute Gel-Based Systems (SGBS), as well. Haney et al. (2008) pulled together data that exhibited very strong correlations (r^2 , up to 0.95) between titration and both IRGA and SGBS to suggest that respiration measurements that do not employ titration are just as valid and accurate as the measurements that do (Haney et al. 2008). However, gel-based methods could employ semi-quantitative color scales and may not result in accuracy for quantification; employing an IRGA system will incur higher costs on equipment while potentially also employing controlled chambers. IRGA-based chambers (static or flow) are similarly a commonly-used method that provides a high temporal resolution and real-time CO₂ flux data. Automated IRGA systems allow for frequent or continuous samplings, but that still employs complicated infrastructure and in-field may involve timing versus scaling tradeoffs, at least within heterogeneous field plots (Jennifer et al., 2002; Kathleen et al., 2003). In contrast, your effective spectrophotometric NaOH trap method uses simple reagents, a colorimeter, and standard lab glassware while providing low and easy to scale costs.

The revised colorimetric alkali-trap respiration method described in this study shows good agreement with established instrumental methods while addressing the limitations of traditional titration methods. Conventional acid-base titration has been widely used to measure CO₂ evolution, but it is limited by subjective endpoint detection, operator differences, and its proximity to atmospheric CO₂ contamination (Bekku, et al. 1997; Haney, et al. 2008).

The most reliable instrumental methods are infrared gas analyzers (IRGA) combining high temporal resolution and gas-phase CO₂ measurement accuracy, which is the current benchmark for soil respiration studies (Pumpanen et al. 2004; Tang et al. 2005). However, IRGA systems come with low costs, require the use of chambers with specific controls, and require continuous calibration for measurements, making them relatively inaccessible in routine soil laboratories or where resources are limited (Subke & Bahn 2010). Gel detectors (either colorimetric in sol-gel or gel form) are low-cost alternatives that have been used in CO₂ detection, but often adopt semi-quantitative visual scales that limit accuracy and reproducibility (Roller et al., 2016). The strong regression between NaOH concentration and OD indicates that absorbance may be a more

sensitive index of alkalinity than pH is consistent with past research showing pH responds non-linearly to form carbonate species, especially near the transition points in colour indicators. Moreover, decreasing the NaOH's exposure to the atmosphere during the colorimetric workflow is an operational approach to reduce methodological exposure to any bias introduced is easy which is present frequently in titration approaches (Haney et al., 2008). Using absorbance measurements instead of titrating NaOH also reduces the time one encounters the reagents--it is a great way to reduce variability tied to human error, which has been identified as a substantial source of variability when performed in levels of titration (Kemmitt et al., 2006). Overall, the modified alkali-trap method and colorimetric methods can be seen as a useful substitution for traditional titration approaches as well as costly instrumental systems. The colorimetric method retains reliability once it is within reason, to comparative data while also improving reproducibility, cost-efficacy as well as, flexibility to allow high-throughput analysis of soil respiration. All of these findings support and promote the advancement of colorimetric detection as a reliable routine method for soil health and microbial respiration, especially in labs where availability is based on IRGAs.

5. PERFORMANCE SUMMARY AND ADVANTAGES

The colorimetric assay for modified alkali-trap soil respiration showed good analytical properties (e.g., slope of the calibration curve > 0.995) and significant benefits over the standard titration method to remove subjective endpoint determination, operator errors, and enhance precision (5.8% vs 1.9% coefficient of variation). The variability of the blanks was also significantly reduced with minimal reagent exposure and quick absorbance reads, further lessening the atmosphere CO₂ interference of ±0.70 mL HCl equivalent to ±0.07 absorbance units. The method was very sensitive with the ability to detect small differences in NaOH concentration and to take an accurate measurement of soils with low respiratory activity. Overall, the colorimeter offered many advantages over titration in numerous areas including speed of processing, reduced reagent usage for analysis, and safer handling and ease of previously or commonly treated soils were in high-throughput and paired with plate readers. The progress made in methodology products improvements

toward the development of a reliable, fast, reproducible, and sensitive method to measure soil respiration for soil quality assessments, microorganism activity, carbon sequestration research, or fertilizer evaluation.

6. CONCLUSION

The modified colorimetric alkali-trap assay developed in this study provides a rapid, accurate, and reproducible method for quantifying soil-respired CO₂. The strong correlations obtained between NaOH concentration and both pH ($r = 0.991$) and optical density ($r = 0.999$) confirm high analytical reliability, with OD proving more sensitive for detecting alkalinity changes. Among the tested soils, organic-fertilizer soil showed the highest respiration (21.67 moles/g/day), followed by vermicompost and black soil, while sand exhibited the lowest microbial activity. The method eliminates subjective titration endpoints, minimizes atmospheric CO₂ interference, and reduces operator-dependent variability. Overall, the colorimetric approach offers a cost-effective, high-throughput alternative to traditional titration and costly IRGA systems for routine soil microbial respiration assessment.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Agase, D. M., & Tiwari, H. H. (2025). Smart irrigation for sustainable agriculture: A crop-specific and soil-responsive approach. *International Journal of Plant & Soil Science*, 37(10), 411–418. <https://doi.org/10.9734/ijpss/2025/v37i105796>
- Bekku, Y., Koizumi, H., Oikawa, T., & Iwaki, H. (1997). Examination of four methods for measuring soil respiration. *Applied Soil Ecology*, 5(3), 247–254. [https://doi.org/10.1016/S0929-1393\(96\)00131-X](https://doi.org/10.1016/S0929-1393(96)00131-X)
- Bond-Lamberty, B., & Thomson, A. (2010). A global database of soil respiration data. *Global Change Biology*, 16(3), 791–802. <https://doi.org/10.5194/bg-7-1915-2010>
- Crowther, T. W., Todd-Brown, K. E. O., Rowe, C. W., Wieder, W. R., Cary, A., Machmuller, M., et al. (2016). Quantifying global soil carbon losses in response to warming. *Nature*, 540(7631), 104–108. <https://doi.org/10.1038/nature20150>
- Doran, G., & Zander, A. (2012). An improved method for measuring soil microbial activity by gas phase flow injection analysis. *Revista Brasileira de Ciência do Solo*, 36(2), 349–357. <https://doi.org/10.1590/S0100-06832012000200004>
- Doran, J. W., & Zeiss, M. R. (2000). Soil health and sustainability: Managing the biotic component of soil quality. *Applied Soil Ecology*, 15(1), 3–11. [https://doi.org/10.1016/S0929-1393\(00\)00067-6](https://doi.org/10.1016/S0929-1393(00)00067-6)
- Gougoulas, C., Clark, J. M., & Shaw, L. J. (2014). The role of soil microbes in the global carbon cycle: Tracking the below-ground microbial processing of plant-derived carbon for manipulating carbon dynamics in agricultural systems. *Journal of the Science of Food and Agriculture*, 94(12), 2362–2371. <https://doi.org/10.1002/jsfa.6577>
- Haney, R. L., Brinton, W. H., & Evans, E. (2008). Estimating soil carbon, nitrogen, and phosphorus mineralization from short-term carbon dioxide respiration. *Communications in Soil Science and Plant Analysis*, 39(17–18), 2706–2720. <https://doi.org/10.1080/00103620802358862>
- Hashimoto, S., Carvalhais, N., Ito, A., et al. (2015). Global spatiotemporal distribution of soil respiration modeled using a global database. *Biogeosciences*, 12(13), 4121–4132. <https://doi.org/10.5194/bg-12-4121-2015>
- Irving, D., Bakhshandeh, S., Tran, T. K. A., & McBratney, A. B. (2024). A cost-effective method for quantifying soil respiration. *Soil Security*, 100162. <https://doi.org/10.1016/j.soisec.2024.100162>
- Kemmitt, S. J., Wright, D., Goulding, K. W. T., & Jones, D. L. (2006). pH regulation of carbon and nitrogen

- dynamics in two agricultural soils. *Soil Biology and Biochemistry*, 38(5), 898–911.
<https://doi.org/10.1016/j.soilbio.2005.08.006>
- Knoepp, J. D., & Vose, J. M. (2002). Quantitative comparison of in situ soil CO₂ flux measurement methods (Vol. 28). United States Department of Agriculture, Forest Service, Southern Region Station.
<https://doi.org/10.2737/SRS-RP-28>
- Kostjukov, V. (2025). Absorption of colored phenolphthalein dianion in aqueous solution: A theoretical analysis. *Chemical Physics*, 112888.
<https://doi.org/10.1016/j.chemphys.2025.112888>
- Pumpanen, J., Ilvesniemi, H., & Hari, P. (2004). A respiration measurement system for intact forest floor. *Tree Physiology*, 24(3), 273–279.
<https://doi.org/10.1093/treephys/24.3.273>
- Roller, A., Schmidt, M. W. I., & Kögel-Knabner, I. (2016). High-throughput quantification of soil respiration using colorimetric detection. *European Journal of Soil Science*, 67(2), 147–157.
<https://doi.org/10.1111/ejss.12316>
- Savage, K. E., & Davidson, E. A. (2003). A comparison of manual and automated systems for soil CO₂ flux measurements: Trade-offs between spatial and temporal resolution. *Journal of Experimental Botany*, 54(384), 891–899.
<https://doi.org/10.1093/jxb/erg121>
- Schimel, J. P., & Schaeffer, S. M. (2012). Microbial control over carbon cycling in soil. *Frontiers in Microbiology*, 3, 348.
<https://doi.org/10.3389/fmicb.2012.00348>
- Subke, J. A., & Bahn, M. (2010). On the “temperature sensitivity” of soil respiration: Can we use the immeasurable to predict the unknown? *Soil Biology and Biochemistry*, 42(9), 1653–1656.
<https://doi.org/10.1016/j.soilbio.2010.05.026>
- Tang, J., Baldocchi, D. D., & Xu, L. (2005). Tree photosynthesis modulates soil respiration on a diurnal time scale. *Global Change Biology*, 11(8), 1298–1304.
<https://doi.org/10.1111/j.1365-2486.2005.00978.x>

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://pr.sdiarticle5.com/review-history/148908>