



Effects of Nano DAP on Growth and Yield of Tea (*Camellia assamica* sub-sp. *lasiocalyx* (Planch. MS))

**Krishnakhee Borah ^{a++}, Gautam Kr. Saikia ^{a#}, Mridul Deka ^{a#},
Aditi Smith Gogoi ^{a†} and Bidisha Hazarika ^{a‡}**

^a Department of Tea Husbandry and Technology, Assam Agricultural University, Jorhat, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ijpss/2025/v37i125884>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://pr.sdiarticle5.com/review-history/149397>

Original Research Article

Received: 19/10/2025

Accepted: 08/12/2025

Published: 17/12/2025

ABSTRACT

A study was conducted to investigate the growth and yield parameters of tea after application of Nano DAP on tea plants during the time frame of February, 2024 to December, 2024. The experiment was done on TV 23 clone of mature tea plants in experimental garden for plantation crops, AAU-Jorhat. The layout was in RBD with four replications and five treatments. Nano DAP was found effective towards the growth of tea plants. The growth and yield attributes of tea plants increased with application of Nano DAP. The growth and yield parameters were recorded significantly higher in treatments T₂ (Recommended doses of N, P₂O₅, K₂O + 3 sprays of Nano DAP @ 3ml/l) and T₃ (Recommended dose of N and K₂O + 75% recommended dose of P₂O₅ + 3 foliar sprays of Nano DAP @ 3 ml/l) over treatment T₁ (Recommended doses of N, P₂O₅ and K₂O). The study revealed that the application of three sprays of Nano DAP @ 3ml/l with recommended dose of fertilizers performed similar with reduction of 25 per cent of basal dose of recommended phosphate

⁺⁺M.Sc. Agriculture; [#]Professor; [†]Associate professor; [‡]PhD. Scholar;

*Corresponding author: E-mail: Krishnakhee.borah.amj22@aaau.ac.in;

and recommended dose of nitrogen and potassium along with three sprays of Nano DAP in tea plants. This approach can be a potential and economically feasible method to increase yield of tea plants and to promote the sustainable development of the tea industry.

Keywords: Nano DAP; tea; foliar application; plucking point density; fine leaf count; fresh weight of plucked shoots; dormancy index; green leaf yield.

1. INTRODUCTION

Tea (*Camellia assamica* sub-sp. *lasiocalyx* (Planch. MS)) is a commercially grown perennial crop and widely cultivated across the world. Tea is the world's second most consumed beverage after water. India's tea sector is one of the most organized industries, contributing significantly to the nation's economy (Adhikari et al., 2014).

The tea production in world reached 6422.66 million kg in 2022 (Tea statistics, Tea board of India, <https://www.teaboard.gov.in/>). India produced 1291.49 million kg during 2023-24 (Tea statistics, Tea board of India, <https://www.teaboard.gov.in/>). At the international level, China, with a production of 3090.00 million kg (in the year 2020) dominates the world tea market (Tea statistics, Tea board of India, <https://www.teaboard.gov.in/>). Total export of tea from India has estimated at 226.98 million kg in the year 2022 (Tea statistics, Tea board of India, <https://www.teaboard.gov.in/>).

The fertilizer requirement for tea crop varies at different growth stages and under different agro-climatic & soil conditions. Nitrogen, phosphorus and potassium are the most important primary macronutrients required by tea. One quintal of made tea removes 5.00 kg of nitrogen, 1.00 kg of phosphate and 2.00 kg of potash from soil (Adnan et al., 2003). Fertilizer application is an important part of the normal intensive production of tea and one of the regular field management practices with significant bearing on both yield and quality of tea. Manuring in mature tea is done on the basis of cyclic yield of the crop and available soil potash content. Soil potash status is termed as low when available potash content in soil is less than 60 ppm, medium when available potash content in soil ranges from 60 to 100 ppm & high when available potash content in soil is more than 100 ppm (Adnan et al., 2003).

Fertilization with nitrogen, phosphorus and potassium is an effective strategy for achieving high tea yield and quality (Anon, 1996). Nitrogen is an essential mineral nutrient for plant growth and reproduction. Apart from being a

fundamental building block of proteins and nucleic acids, nitrogen also participates in carbon fixation through photosynthesis as a component of chlorophyll (Anon, 2023). Nitrogenous fertilizer application can significantly improve tea yield and quality (Kumar et al., 2018). It facilitates the formation of amino acids, including theanine, arginine, and glutamic acid, which are positively correlated with tea quality (Sun et al., 2023).

The bulk of the inorganic nitrogen, phosphorus and potassium given to the soil is lost by leaching, erosion, volatilization, or is immobilized in soil organic matter. Only a small part of soil nitrogen is readily available to the plants and the majority remains unavailable (Barua, 1989).

Phosphorus is crucial for plant development and reproduction and is a key component of fertilizers for modern agriculture. Phosphorus helps in photosynthesis, energy storage, And cell division. The availability of phosphorus governs the uptake of nitrogen and potassium as well. Adequate response to nitrogen is possible only when phosphate need of the soil is well satiated (Barua & Dutta, 1972. An analysis of 30 years data of NPK experiments in Assam shows significant and positive interaction between phosphorus and nitrogen (Bernard & Habash, 2009). Positive effects of P-K interactions on yield were noticed in long-term NPK trials in tea plantations of Assam and Doors (Biswas & Biswas, 1984) as well as in Sri Lanka (Biswas & Biswas, 1984).

In acidic and highly weathered soils, 75-80 per cent of the applied phosphate is fixed onto the surfaces of iron and aluminium oxides and hydroxides (Bray & Kurtz, 1945). It has been reported that less than 15 per cent of fertilizer applied phosphorus is taken up by the crop during the year of application (Deo et al., 2022).

In the present investigation, foliar spraying of Nano DAP along with recommended basal doses of N and K₂O was found to be an efficient method for increasing growth and yield attributes in tea plants. Nanotechnologies have the potential to produce a significant boost in crop yield along with improvement of food production

systems. Nanotechnology has a greater nutrient uptake efficiency, and it will soon transform the methods used for foliar application. Nano fertilizers were an effective alternative to costly conventional fertilizers (Dutta, 1956). Moreover, foliar spraying transports nutrients quickly and efficiently to the location of food synthesis, reducing the need for fertilizer. Foliar nutrient application was found to be more beneficial than soil application in terms of reducing losses due to leaching and fixation. When moisture availability is restricted, fertilizer administration via foliar spray provides excellent absorption. While foliar spray does not replace soil treatment, it should definitely be considered as a supplemental technique (Gomez & Gomez, 1984).

2. MATERIALS AND METHOD

The experiment was conducted at Experimental Garden for Plantations Crops (EGPC), Section number 2(A), Department of Tea Husbandry and Technology, Assam Agricultural University, Jorhat from February, 2024 to December, 2024. The experiment was laid out in Randomized Block Design (RBD) with five treatments and four replications. The planting material for the experiment was TV 23 clone planted with spacing 105 cm x 70 cm. Collection of plucked tea leaves were done using nylon bags.

The treatment details of the experiment are presented below:

T₁ : Recommended doses of N, P₂O₅ and K₂O

T₂ : Recommended doses of N, P₂O₅, K₂O +3 sprays of Nano DAP @ 3ml/l

T₃ : Recommended dose of N and K₂O + 75% recommended dose of P₂O₅ + 3 foliar sprays of nano DAP @ 3 ml/l

T₄ : Recommended dose of N and K₂O + 50% recommended dose of P₂O₅ + 3 foliar sprays of nano DAP @ 3 ml/l

T₅ : Recommended dose of N and K₂O + 25% recommended dose of P₂O₅ + 3 foliar sprays of nano DAP @ 3 ml/l

2.1 Fertilizer Application

In this experiment, the recommended doses of fertilizers were N, P₂O₅, K₂O @ 100:25:75 kg/ha. The fertilizers were applied in the form of Urea, Single Super Phosphate (SSP) and Muriate of Potash (MOP) as basal application. For foliar application, Nano Di Ammonium Phosphate (Nano DAP) was used @ 3ml/l. Nano DAP

(Liquid) is a novel Nano fertilizer developed and patented by IFFCO (Indian Farmers Fertilizer Cooperative Limited) in 2023. It contains Nitrogen (8.0 % N w/v) and Phosphorus (16.0 % P₂O₅ w/v) (Jagadeesh et al., 2024).

2.2 Time Schedule for Application of Fertilizers

The application of Urea and Muriate of Potash (MOP) was applied in 2 splits viz; first split (60%) during April, 2024 and second split (40%) during August, 2024. The Single Super Phosphate (SSP) was applied as first split (100%) during April, 2024. The first, second and third foliar spray of Nano DAP @ 3ml/l was done in April, July, and October, 2024.

2.3 Methods for Estimating Plant Growth and Yield Parameters

2.3.1 Plucking point density (No./2500 cm²)

It was determined by counting numbers of shoots plucked in each plucking round. At each plucking round the plucking point density was calculated by the method described by Jayaraman, (1962). A bamboo grid of 50×50 cm² was placed on the top of the tea bushes and the number of the plucking points was counted from where the shoots were plucked. The records were expressed in no/2500 cm².

2.3.2 Fine leaf count (%)

A bulk of green leaves samples were collected from each nylon bags and weighed 100 g of green leaves. Separate the one leaf and a bud, two leaf and a bud, soft banjhis, single soft leaf and weighed all of them against the total weight of the sample and expressed in percentage (%) (Khemshetty et al., 2024).

2.3.3 Fresh weight of plucked shoot (g)

From each treatment and replication, 100 numbers of shoots were collected after each plucking round. Then weighed them in a digital weighing balance and the data was expressed in grams (g).

2.3.4 Dormancy index

The dormancy Index is the ratio between the number of shoots with buds which are less than half the length of the topmost leaf to those which are greater than half the length of the topmost leaf (Liu et al., 2021).

The dormancy index of tea was measured by using the following formula:

$$\text{Dormancy index} = \frac{\text{The no. of shoots with buds which are less than half the length of the topmost leaf}}{\text{The no. of shoots with buds which are greater than half the length of the topmost leaf}}$$

2.3.5 Green leaf yield (Kg/plot)

Plucking of green leaves was done at a time interval of 7-8 days. The green leaf yield per plot throughout the whole experiment period (February-December, 2024) was recorded against each treatment and replication using weighing balance and expressed in kg/plot.

2.4 Statistical Analysis

Analysis of variance (ANOVA) was carried out as per the design (RBD) following standard statistical procedure. The treatment mean values was compared by least significance difference based on ANOVA (Ma et al., 2021). All computations were carried out by using MS Excel 2021. The critical difference (C.D) values were calculated at 5 per cent level of significance.

3. RESULTS AND DISCUSSION

The experimental findings of the present research experiment are presented below. Field studies were conducted to study the growth and yield parameters of tea plants after application of Nano DAP. The data were statistically analyzed to come to a conclusion on influence of Nano DAP application on tea plants. The mean values were tabulated and the associated CD values at the 5 per cent probability level were computed and displayed in tables.

3.1 Effect of Nano DAP on Growth and Yield of Tea Plants

The growth and yield parameters studied in this experiment were plucking point density, fine leaf count, fresh weight of plucked shoots, dormancy index and green leaf yield. All these yield attributes were recorded with respect to all the treatments during the time period from March, 2024 to November, 2024. For evaluating the yield parameters, the cropping season was divided into three seasons viz., early season (March - May), rainy season (June- September), back-end season (October - November).

3.1.1 Plucking point density (No./2500 cm²)

The experimental findings on number of plucking point density/2500 cm² of tea plants in various treatments of the investigation are presented in Table 1.

As evident from the data presented in Table 1., Plucking point density of tea plants were significantly influenced by application of Nano DAP in all the three seasons. The treatments T₂, T₃ and T₄ showed significant difference in numbers of plucking point density/ 2500 cm² of tea plants over treatment T₁ (Recommended dose of N, P₂O₅ and K₂O) during all three seasons. The highest plucking point density (64.12 no/2500 cm², 83.77 no/2500 cm² and 83.88 no/2500 cm²) was observed in treatment T₃ (Recommended dose of N and K₂O + 75% of recommended dose of P₂O₅ + 3 foliar spray of Nano DAP @ 3 ml/l) followed by plucking point density (64.12 no/2500 cm², 83.59 no/2500 cm², 83.13 no/2500 cm²) observed in treatment T₂ (Recommended dose of N, P₂O₅, K₂O + 3 foliar spray of Nano DAP @ 3 ml/l) in all three seasons. However, treatment T₂ and treatment T₃ exhibited similar results in plucking point density during all three seasons. It was also reported that the plucking point density in tea was maximum (50 no/m²) during rain flush when they applied 0.4% Nano-urea in 3 sprays (Tea Board of India, no year). The increase in plucking point density in tea plants might be due to the synergistic effect of foliar application of Nano DAP along with basal application of recommended fertilizer. Also, the increase in number of pluckable shoots led to higher yield in tea (Saikia & Gogoi, 2023).

The plucking point density (59.50 no/2500 cm², 78.39 no/2500 cm² and 78.26 no/2500 cm²) in treatment T₄ (Recommended dose of N and K₂O + 50% of recommended dose of P₂O₅ + 3 foliar sprays of Nano DAP @ 3 ml/l) was significantly lower than the plucking point density recorded in treatment T₂ and treatment T₃ in all three seasons. The lowest plucking point density (51.50 no/2500 cm², 73.08 no/2500 cm² and 52.94 no/2500 cm²) was observed in treatment T₁ (Recommended dose of N, P₂O₅ and K₂O) followed by treatment T₅ (Recommended dose of N and K₂O + 25% P + 3 foliar sprays of Nano DAP @ 3 ml/l) in all three seasons.

3.1.2 Fine leaf count (%)

The fine leaf count (%) of plucked tea shoots during early season (March - May), rainy season (June- September) and back-end season (October - November) are shown in Table 2.

Table 1. Plucking point density (No./2500 cm²) as influenced by application of Nano DAP based fertilizers

Treatment	Plucking point density (No./2500 cm ²)		
	Early season (March - May)	Rainy season (June - September)	Back-end season (October - November)
T ₁	51.25	66.31	58.63
T ₂	64.12	83.59	83.13
T ₃	64.75	83.77	83.88
T ₄	59.50	78.39	78.26
T ₅	52.50	66.59	65.47
CD	2.34	1.66	2.22
SE(d)	1.07	0.76	1.02




Fig. 1. Plucking point density (no./2500 cm²) as influenced by application of nano DAP based fertilizers

Table 2. Fine leaf count (%) as influenced by application of Nano DAP based fertilizers

Treatment	Fine leaf count (%)		
	Early season (March - May)	Rainy season (June - September)	Back-end season (October - November)
T ₁	54.62	64.62	45.25
T ₂	56.08	68.58	58.87
T ₃	56.78	66.78	54.37
T ₄	55.25	62.75	53.5
T ₅	54.20	59.20	49.5
CD	N/A	5.42	4.57
SE(d)	1.79	2.49	2.10




Fig. 2. Fine leaf count (%) as influenced by application of nano DAP based fertilizers

From the data presented in Table 2, no significant difference was observed among the treatments in fine leaf count (%) during early season. Fine leaf count (%) was significantly higher in treatments T₂, T₃, T₄ as compared to treatment T₁ (Recommended dose of N, P₂O₅ and K₂O) at back-end season. The treatment T₂ (Recommended doses of N, P₂O₅, K₂O + 3 sprays of Nano DAP @ 3ml/l) showed highest fine leaf count (%) in rainy season (68.58%) and back-end season (58.87%). The fine leaf count (%) in treatment T₃ (Recommended dose of N and K₂O + 75% recommended dose of P₂O₅ + 3 foliar sprays of Nano DAP @ 3 ml/l) during rainy season (66.78%) and back-end season (54.37%) was not significantly different from the fine leaf count (%) observed in treatment T₂. Similar results were obtained by Tea Board of India, (no year). The fine leaf count was maximum during first flush when 0.4 per cent Nano-urea was applied in 3

sprays (Tea Board of India, no year). Increase in fine leaf count during back-end season due to application of nano DAP is expected to contribute to the production of quality made tea.

Lowest fine leaf count (%) was observed in treatment T₅ (Recommended dose of N and K₂O + 25% recommended dose of P₂O₅ + 3 foliar spray of Nano DAP @ 3 ml/l) during rainy season (59.20%) and in treatment T₁ (Recommended dose of N, P₂O₅ and K₂O) during back-end season (45.25%).

3.1.3 Fresh weight of 100 shoots (g)

The observations of the fresh weight of 100 shoots (g) during early season (March - May), rainy season (June- September) and back-end season (October - November) are presented in Table 3.

Table 3. Fresh weight of 100 shoots (g) of tea plants as influenced by application of Nano DAP based fertilizers

Treatment	Fresh weight of 100 shoots (g)		
	Early season (March - May)	Rainy season (June - September)	Back-end season (October - November)
T ₁	78.54	94.32	90.40
T ₂	85.20	98.48	94.12
T ₃	83.87	97.25	93.84
T ₄	80.90	95.36	92.01
T ₅	78.29	93.24	84.75
CD	2.58	1.33	1.67
SE(d)	1.18	0.61	0.76




Fig. 3. Fresh weight of 100 shoots (g) of tea plants as influenced by application of nano DAP based fertilizers

The fresh weight of 100 shoots of tea plants recorded in treatments T₂ and T₃ were found significant in early season, rainy season and back-end season. The treatment T₂ (Recommended dose of N, P₂O₅, K₂O + 3 foliar sprays of Nano DAP @ 3 ml/l) was found to produce significantly higher fresh weight of shoots in early season (85.20 g), rainy season (98.48 g) and back-end season (94.12 g) over treatment T₁ (Recommended dose of N, P₂O₅ and K₂O). The treatment T₃ (Recommended dose of N and K₂O + 75% recommended dose of P₂O₅ + 3 foliar sprays of Nano DAP @ 3 ml/l) exhibited similar results in fresh weight of 100 shoots during early season (83.87 g), rainy season (97.25 g) and back-end season (93.84 g) as that of treatment T₂ in all three seasons. These results are in conformity with (Salehi & Hajiboland, 2008). It was reported that an increase in overall growth of young teas (measured in terms of leaf and pruning weights) due to potash manuring in young tea (Ruan et

al., 2010). Nano phosphorus fertilization increased maize weight when compared to other treatments and controls (Poudel et al., 2023). Application of Udaipur nano rock phosphate (34% P₂O₅) considerably boosted the yield attributes like 100 grain weight in maize plants as compared to the control (Magda & Hussein, 2015).

Lowest fresh weight of 100 shoots was observed in treatment T₅ (Recommended dose of N and K₂O + 25% recommended dose of P₂O₅ + 3 foliar spray of Nano DAP @ 3 ml/l) during early season (78.29 g), rainy season (93.24 g) and in back-end season (84.75 g).

3.1.4 Dormancy index

The experimental findings on dormancy index during early season (March - May), rainy season (June- September) and back-end season (October - November) are presented below in Table 4.

Table 4. Dormancy index of tea plants as influenced by application of Nano DAP based fertilizers

Treatment	Dormancy index of tea plants		
	Early season (March - May)	Rainy season (June - September)	Back-end season (October - November)
T ₁	1.08	0.51	1.07
T ₂	0.88	0.49	0.90
T ₃	0.93	0.52	0.97
T ₄	1.01	0.54	1.05
T ₅	1.19	0.56	1.64
CD	0.07	N/A	0.24
SE(d)	0.03	0.02	0.11




Fig. 4. Dormancy index of tea plants as influenced by application of nano DAP based fertilizers

The lowest dormancy index (0.88, 0.90) was observed in treatment T₂ (Recommended dose of N, P₂O₅, K₂O + 3 foliar spray of Nano DAP @ 3 ml/l) followed by treatment T₃ (Recommended dose of N and K₂O + 75% recommended dose of P₂O₅ + 3 foliar spray of Nano DAP @ 3 ml/l) in early season (0.88) and back-end (0.90) season respectively. However, treatment T₂ and T₃ were not significantly different in both early and back-end season. These findings are in close agreement with (Sarkar et al., 2021). More than 75 per cent of the stored nitrogen within the whole plant is re-translocated to promote the sprouting of dormant buds in tea plants (Tang et al., 2021).

The dormancy index (1.08, 1.07) in treatment T₄ (Recommended dose of N and K₂O + 50% recommended dose of P₂O₅ + 3 foliar sprays of Nano DAP @ 3 ml/l) and dormancy index (1.01, 1.05) in treatment T₁ (Recommended dose of N, P₂O₅ and K₂O) were at par in both early season and back-end season. No significant difference

was observed in dormancy index during rainy season.

The treatment T₅ (Recommended dose of N and K₂O + 25% recommended dose of P₂O₅ + 3 foliar sprays of Nano DAP @ 3 ml/l) showed highest dormancy index (1.19, 1.64) in early season and back-end season respectively. The reduction in basal application of phosphate in treatment T₅ might hinder the growth of buds as compared to growth of buds in treatments T₂, T₃ and T₄ due to reduced photosynthesis rate. The growth of tea plants is directly related to photosynthesis and phosphorus deficiency inhibited photosynthesis in tea plants (Thakur et al., 2024).

3.1.5 Green leaf yield (kg/plot)

The experimental findings on green leaf yield of tea plants (kg/plot) during early season (March - May), rainy season (June- September) and back-end season (October - November) are presented in Table 5.

Table 5. Green leaf yield (kg/plot) of tea plants as influenced by application of Nano DAP based fertilizers

Treatment	Green leaf yield (kg/plot) (plot size = 44.10 m ²)		
	Early season (March - May)	Rainy season (June - September)	Back-end season (October - November)
T ₁	3.18	4.89	4.21
T ₂	4.45	6.32	6.25
T ₃	4.33	6.24	6.10
T ₄	3.83	5.75	5.71
T ₅	3.27	4.94	4.48
CD	0.24	0.14	0.26
SE(d)	0.11	0.06	0.12




Fig. 5. Green leaf yield (kg/plot) of tea plants as influenced by application of nano DAP based fertilizers

From the data presented in Table 5, it was evident that the green leaf yield recorded in treatments T₂, T₃ and T₄ showed significant difference over treatment T₁ (Recommended doses of N, P₂O₅, K₂O). The highest green leaf yield (4.45 kg/plot, 6.32 kg/plot and 6.25 kg/plot) was observed in treatment T₂ (Recommended doses of N, P₂O₅, K₂O +3 sprays of Nano DAP @ 3ml/l) followed by green leaf yield (4.33 kg/plot, 6.24 kg/plot, 6.10 kg/plot) observed in treatment T₃ (Recommended dose of N and K₂O + 75% recommended dose of P₂O₅ + 3 foliar spray of Nano DAP @ 3 ml/l) in all three seasons. The treatment T₃ exhibited similar results in green leaf yield as that of treatment T₂ during all three seasons. These findings are in close agreement with the findings of (Tea Board of India, no year) in green leaf yield of tea crop for nano urea. Similar findings were reported in Nano DAP for other crops in grain yield of wheat (Zulfiqar et al., 2019), in grain yield of rice (Okano et al., 1994), in grain yield of maize (Magda & Hussein, 2015), and in finger millet (Sharma et al., 1977).

The green leaf yield (3.83 kg/plot, 5.75 kg/plot and 5.71 kg/plot) observed in treatment T₄ (Recommended dose of N and K₂O + 50% recommended dose of P₂O₅ + 3 foliar sprays of Nano DAP @ 3 ml/l) was significantly lower than the green leaf yield observed in treatments T₂ and T₃ in all three seasons. Lowest green leaf yield (3.18 kg/plot, 4.89 kg/plot, 4.21 kg/plot) was observed in treatment T₁ (Recommended doses of N, P₂O₅ and K₂O) which was not significantly different from treatment T₅ (Recommended dose of N and K₂O + 25% recommended dose of P₂O₅ + 3 foliar sprays of Nano DAP @ 3 ml/l) in all three seasons.

4. CONCLUSION

From the present investigation, it is evident that all the yield attributes viz., plucking point density, fresh weight of 100 shoots, fine leaf count, dormancy index was recorded maximum when three foliar sprays of Nano DAP @ 3ml/l was applied along with recommended dose of N, P₂O₅, K₂O or recommended dose of N and K₂O and 75% of recommended dose of P₂O₅ in tea. Higher plucking point density, shoot weight and reduction of dormancy index might have contributed to higher yield in these treatments. Based on the superior results on these growth and yield attributes and considering reduction in cost of cultivation, foliar application of three sprays of Nano DAP @ 3ml/l along with recommended basal dose of N & K₂O and 75 per cent of recommended dose of phosphate can be suggested for application in tea plants to get higher yield.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of this manuscript

Details of the AI usage are given below:

1. QuillBot
2. ChatGPT

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Adhikari, T., Kundu, S., Meena, V., & Rao, A. S. (2014). Utilization of nano rock phosphate by maize (*Zea mays* L.) crop in a vertisol of Central India. *Journal of Agricultural Science and Technology. A.*, 4(5A).
- Adnan, A., Mavinic, D. S., & Koch, F. A. (2003). Pilot-scale study of phosphorus recovery through struvite crystallization – examining the process feasibility. *Journal of Environmental Engineering and Science*, 2(5), 315–324. <https://doi.org/10.1139/s03-040>
- Anon. (1996). The planter's Handbook. Tea Research Association. <https://www.tocklai.org/product/planters-handbook/>
- IFFCO. (2023). Nano DAP brochure. <https://www.iffco.in/products/nano-fertilisers/nano-dap-liquid>
- Barua, D. N. (1989). Science and practice in tea culture. Tea Research Association. <https://www.tocklai.net/ebooks/>
- Barua, D. N., & Dutta, K. N. (1972). Distribution of shoots on the plucking surface of a tea bush and its relation to spacing. *Two and A Bud*, 18, 8-11. <https://doi.org/10.5555/19720305138>
- Bernard, S. M., & Habash, D. Z. (2009). The importance of cytosolic glutamine synthetase in nitrogen assimilation and recycling. *New Phytologist*. <https://doi.org/10.1111/j.1469-8137.2009.02823.x>
- Biswas, A. K., & Biswas, A. K. (1984). Manuring syndrome for mature tea: a long term study. *Two and A Bud*, 31, 22-27.
- Bray, R. H., & Kurtz, L. T. (1945). Determination of total, organic, and available forms of phosphorus in soils. *Soil Science*. <https://doi.org/10.1097/00010694-194501000-00006>
- Buckman, H. O., & Brady, N. C. (1960). The nature and properties of soils. Macmillan.
- Deo, H. R., Chandrakar, T., Srivastava, L. K., Nag, N. K., Singh, D. P., & Thakur, A. (2022). Effect of Nano-DAP on yield, nutrient uptake and nutrient use efficiency by rice under Bastar plateau. *J. Pharm. Innov.*, 11(9), 1463-1465. <https://www.thepharmajournal.com/archive/s/2022/vol11issue9/PartU/11-9-10-900.pdf>
- Dutta, S. K. (1956). Manuring. Proc. 12th Annual Conf. Tocklai Experimental Station. Jorhat. pp. 5-12.
- Gomez, K. A., & Gomez, A. A. (1984). *Statistical procedures for agricultural research*. John Wiley & sons. <https://www.wiley.com/en-us/Statistical+Procedures+for+Agricultural+Research%2C+2nd+Edition-p-9780471870920>
- Jagadeesh, M. S., Vinay, H. T., Pavithra, V., & Abhishek, G. J. (2024). India's Tea Export Potential: Stirring up Global Trade Opportunities. *Journal of Experimental Agriculture International.*, 46(9) <https://doi.org/10.9734/jeai/2024/v46i92827>.
- Jayaraman, V. (1962). Fertilizer use in tea in South India. *Fertiliser News*, 7(8): 7-11. <https://www.cabidigitallibrary.org/doi/full/10.5555/19650306829>
- Khemshetty, A., Patil, D. H., Rathod, P. S., Patil, A. S., & Basavaraj, K. (2024). Studies on Nano DAP on Growth, Yield and Quality of Chickpeas under Rainfed Conditions of Northeastern Dry Zone of Karnataka. *Journal of Experimental Agriculture International*, 46(3), 139-145. <https://doi.org/10.9734/jeai/2024/v46i32332>
- Kumar, V., Kaur, J., Tanwar, B., Goyal, A., Gat, Y., Kumar, A., & Kaur, P. (2018). Tea processing. *Beverages: Processing and technology*, 190-209. ISBN: 978-93-87869-36-3
- Liu, M. Y., Tang, D., Shi, Y., Ma, L., Zhang, Q., & Ruan, J. (2021). Foliar N application on tea plant at its dormancy stage increases the N concentration of mature leaves and improves the quality and yield of spring tea. *Frontiers in Plant Science*, 12: 753086. <https://doi.org/10.3389/fpls.2021.753086>
- Ma, L., Yang, X., Shi, Y., Yi, X., Ji, L., Cheng, Y., Kang, N., & Ruan, J. (2021). Response of tea yield, quality and soil bacterial characteristics to long-term nitrogen fertilization in an eleven-year field experiment. *Applied Soil Ecology*, 166, 103976. <https://doi.org/10.1016/j.apsoil.2021.103976>
- Magda, S., & Hussein, M. M. (2015). Usage of the nano phosphorous fertilizers in enhancing the corn crop and its effect on corn borers infestations after fungi treatments. *International Journal of ChemTech Research*, 8(9), 167-173. [https://sphinxesai.com/2015/ch_vol8_no9/1/\(167-173\)V8N9CT.pdf](https://sphinxesai.com/2015/ch_vol8_no9/1/(167-173)V8N9CT.pdf)

- Mu, N., Gaddi, A. K., H, V., S, R., & MY, A. (2024). Effect of nano DAP on growth and yield of finger millet. *International Journal of Research in Agronomy*, 7(10), 521-524. <https://doi.org/10.33545/2618060X.2024.v7.i10g.1823>
- Okano, K., Komaki, S., & Matsuo, K. (1994). Remobilization of nitrogen from vegetative parts to sprouting shoots of young tea (*Camellia sinensis* L.) plants. *Japanese Journal of Crop Science*. <https://doi.org/10.1626/jcs.63.125>
- Poudel, A., Singh, S. K., Jiménez-Ballesta, R., Jatav, S. S., Patra, A., & Pandey, A. (2023). Effect of nano-phosphorus formulation on growth, yield and nutritional quality of wheat under semi-arid climate. *Agronomy*, 13(3), 768. <https://doi.org/10.3390/agronomy13030768>
- Ranganatha, V., & Natesan, S. (1985). Potassium nutrition of tea. In *Potassium in agriculture* (pp. 981-1022). <https://doi.org/10.2134/1985.potassium.c42>
- Ruan, J., Haerdter, R., & Gerendás, J. (2010). Impact of nitrogen supply on carbon/nitrogen allocation: a case study on amino acids and catechins in green tea [*Camellia sinensis* (L.) O. Kuntze] plants. *Plant Biology*, 12(5), 724-734. <https://doi.org/10.1111/j.1438-8677.2009.00288.x>
- Saikia, P., & Gogoi, A. S. (2023). Growth and Yield of Tea (*Camellia sinensis*) as Influenced by Nano Urea during Pre and Post Pruning Operation. *International Journal of Plant & Soil Science*. <https://doi.org/10.9734/IJPSS/2023/v35i203915>
- Salehi, S. Y., & Hajiboland, R. (2008). A high internal phosphorus use efficiency in tea (*Camellia sinensis* L.) plants. *Asian Journal of Plant Sciences*, 7(1), 30-36. <https://doi.org/10.3923/ajps.2008.30.36>
- Sarkar, A., Jana, K., & Mondal, R. (2021). Growth and yield of hybrid mustard (*Brassica juncea* L.) as influenced by foliar nutrition in Gangetic plains of West Bengal. *Journal of Crop and Weed*, 17(3), 35-40. <https://doi.org/10.22271/09746315.2021.v17.i3.1488>
- Sharma, P. C., Jain, N. K., & Biswas, A. K. (1977). Response to nitrogen, phosphate and potash in mature tea. In *28th Conference, Tocklai Experimental Station* (pp. 31-34).
- Sun, L., Zhang, Z., Li, Y., Ruan, J., Karak, T., & Yang, T. (2023). Editorial: Mineral nutrients on tea yield and quality formation. *Frontiers in Plant Science*, 14, 1192432. <https://doi.org/10.3389/fpls.2023.1192432>
- Tang, S., Zheng, N., Ma, Q., Zhou, J., Sun, T., Zhang, X., & Wu, L. (2021). Applying Nutrient Expert system for rational fertilization to tea (*Camellia sinensis*) reduces environmental risks and increases economic benefits. *Journal of Cleaner Production*. <https://doi.org/10.1016/j.jclepro.2021.127197>
- Tea Board of India. (n.d.). Tea statistics. <https://www.teaboard.gov.in/>
- Thakur, D., Kumar, N., Manuja, S., Hetta, G., Anshumali, Mridula, Chakraborty, M., Kumari, P., & Saqib, A. (2024). Effect of foliar spray of Nano DAP on growth of wheat (*Triticum aestivum* L.). *Himachal Journal of Agricultural Research*, 50(2), 284-290. <https://hjar.org/index.php/hjar/article/view/172604>
- Zulfiqar, F., Navarro, M., Ashraf, M., Akram, N. A., & Munné-Bosch, S. (2019). Nanofertilizer use for sustainable agriculture: Advantages and limitations. *Plant Science*. <https://doi.org/10.1016/j.plantsci.2019.110270>

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://pr.sdiarticle5.com/review-history/149397>