



Harmful Insects of Okra and Eggplant : A Study Conducted in the Lowland Areas of Daloa

Ahouty Ahouty Bernardin ^{a*},
Yao Blé Gbacla Flora Dominique ^a, Kouadio Yao Assène ^a,
Kassi Bohoussou Georges ^b, Ouattara Ouangbo Yaya ^b,
Dago Lydia Karmelle Ornella ^a and Koffi N'goran Mathurin ^b

^a Faculty of Agroforestry, Jean Lorougnon Gédé University, BP 150 Daloa, Côte d'Ivoire.

^b Faculty of Environmental Sciences, Jean Lorougnon Gédé University, BP 150 Daloa, Côte d'Ivoire.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ijpss/2025/v37i125892>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://pr.sdiarticle5.com/review-history/149450>

Original Research Article

Received: 17/10/2025

Accepted: 14/12/2025

Published: 23/12/2025

ABSTRACT

Aims: This study aims to inventorize and characterize the harmful insects associated with okra and eggplant in the city of Daloa (Central-Western Côte d'Ivoire).

Study Design: Descriptive transversal survey.

Place and Duration of Study: In various lowland areas of the city of Daloa during the dry season, from February to April 2025.

Methodology: Harmful insects were collected from okra and eggplant crops located in various lowland areas of the city of Daloa. At each site, ten (10) plants of each crop were randomly selected when okra and eggplant were cultivated together; when only one of the two species was

*Corresponding author: E-mail: bernadinahouty@yahoo.fr;

present, the same procedure was applied to the available crop. Sampling focused on the leaves and fruits of mature, productive plants. Insects were captured using transparent white plastic bags. The method consisted of enclosing a portion of the plant, including leaves and/or fruits, inside a labeled bag and then gently shaking this part of the plant to dislodge and trap the harmful insects inside the bag.

Results: In total, three species of jassids were identified, along with two additional types of pests: phytophagous bugs and aphids. The molecular approach confirmed the identification of the jassids and reliably distinguished them from the other insects collected.

Conclusion: The combination of these approaches provides a better understanding of pest distribution, their relationship with host plants, and their relevance for developing effective plant protection strategies.

Keywords: Okra; eggplant; insect pests; Jassid; aphid; bug.

1. INTRODUCTION

The agricultural sector plays a pivotal role in the economy and development of Côte d'Ivoire, both through the size of its active agricultural workforce and its substantial contribution to national wealth creation. Representing approximately one quarter of the Gross Domestic Product (GDP), agriculture employs nearly half of the working-age population (Ducroquet et al., 2017). Côte d'Ivoire ranks among the leading producers of several major agricultural commodities on both the African continent and the global market and is the undisputed world leader in cocoa bean production (Ducroquet et al., 2017). Although less emphasized, food crop production also occupies a significant position in national agriculture. Vegetable farming, in particular, constitutes a critical economic activity, contributing to food security by ensuring stable supplies of fresh produce. Despite its importance, this sector in Côte d'Ivoire remains predominantly oriented toward self-consumption and subsistence agriculture (FAO, 2018), with more than 83% of producers relying on it to meet household needs (Soro et al., 2018). Among the main food crops, okra (*Abelmoschus esculentus*) and eggplant (*Solanum melongena*) are widely cultivated across Côte d'Ivoire (Tuo et al., 2023; Konan et al. 2020). These crops serve both as essential dietary staples and as major income sources for smallholder farmers (Widadie et al., 2024). However, their production is severely threatened by a diverse range of insect pests capable of inducing substantial yield losses. Key pest groups include jassids (leafhoppers), aphids, whiteflies, thrips, caterpillars, beetles, and fruit flies (Koné et al., 2019). These insects attack various parts of the plants, such as leaves, stems, flowers, and fruits, causing deformation, wilting, and reduced productivity (Borkakati et al., 2018). The recent spread of *Amrasca biguttula* in

Côte d'Ivoire has caused drastic yield declines in crops such as okra, eggplant, and cotton, posing a significant threat to national food security and farmers' livelihoods (Kouadio et al., 2022). Pest management in these cropping systems frequently relies on the use of chemical pesticides. However, the misuse or overuse of insecticides is widespread, often resulting in increased resistance among pest populations (Sène et al., 2020). Moreover, morphological similarity among certain pest species complicates accurate field identification, which can lead to inappropriate or ineffective control strategies. Genetic variations within and among populations of the pest species could further underscore the need for integrative diagnostic approaches combining morphological and molecular tools (Seehausen et al., 2020). Accurate identification and characterization of pest species are therefore essential prerequisites for developing targeted, sustainable, and effective pest management strategies.

In this context, the present study aims to inventorize and characterize the major insect pests associated with okra and eggplant cultivated in the lowland areas of Daloa. By generating updated morphological and genetic data on local pest populations, this work seeks to support improved diagnostic accuracy and guide the development of more appropriate integrated pest management strategies for vegetable growers in Côte d'Ivoire.

2. METHODOLOGY

2.1 Study Area

This study was conducted in the city of Daloa, specifically within the lowland areas that serve as vegetable production zones, largely managed by

women. Okra and eggplant are cultivated either as monocultures or in intercropping systems. Six (6) study sites were randomly selected, based on the presence of these crop species, for insect sampling.

2.2 Sampling

harmful insects were collected from okra and eggplant crops located in various lowland areas of the city of Daloa. The study was conducted during the dry season, from February to April 2025, a period during which pests typically cause more severe damage to vegetable crops than during the rainy season (Zanzana et al., 2025 ; Prabhakar et al., 2024). At each site, ten (10) plants of each crop were randomly selected when okra and eggplant were cultivated together; when only one of the two species was present, the same procedure was applied to the available crop. Sampling focused on the leaves and fruits of mature, productive plants.

Insects were captured using transparent white plastic bags (Fig. 1). The method consisted of enclosing a portion of the plant, including leaves and/or fruits, inside a labeled bag as illustrated in Fig. 1, and then gently shaking this part of the plant to dislodge and trap the pests inside the bag. Although this procedure allowed the

collection of a considerable number of insects, it remains non-exhaustive, as some pests may escape capture. The labeled bags corresponding to each sampling site were transported to the laboratory on the same day. Collected specimens were examined under a binocular microscope for identification and subsequently preserved in labeled tubes containing alcohol.

2.3 Morphological Identification of Collected Pests

For more accurate identification, adult specimens were prioritized. Identification was carried out using the consolidated keys provided by Dmitriev and Dietrich (2009) available at :

<http://dmitriev.speciesfile.org/key.asp?key=Erythroneura&lng=En&i=1&keyN=12>.

Morphological criteria such as body size and shape, coloration, setae arrangement, wing patterns, antenna shape, and specific structures including mouthparts, legs, and the presence or absence of particular projections were examined to identify the collected specimens. Identification was performed using a microscope at 4x and 10x magnification.



Fig. 1. Trapping of insect pests using a plastic bag

2.4 Molecular Analysis

2.4.1 DNA extraction

DNA extraction from the collected specimens was performed using a 2% CTAB protocol (Aikpon et al., 2020). The procedure consisted of grinding each insect specimen in a tube containing 200 μ L of 2% CTAB, followed by the addition of 20 μ L of Proteinase K and incubation in a water bath at 65 °C for 20 minutes. After incubation, 200 μ L of chloroform was added, and the mixture was gently inverted to homogenize. The samples were then centrifuged for 15 minutes at 12,000 rpm at room temperature, after which the upper aqueous phase was transferred to a new tube. Subsequently, 200 μ L of isopropanol was added to the supernatant and mixed by gentle inversion. A second centrifugation was carried out for 15 minutes at 12,000 rpm at room temperature, and the isopropanol was carefully discarded. The pellet was drained thoroughly, then washed with 200 μ L of 70% ethanol. The samples were centrifuged for 5 minutes at 12,000 rpm at room temperature, the ethanol was removed, and the pellet was allowed to air-dry for 24 hours. Finally, 100 μ L of molecular-grade water were added to reconstitute the DNA.

2.4.2 DNA amplification by Polymerase Chain Reaction (PCR)

DNA was amplified using the universal primer pair COI-F (5'-ATAATTTTTTTTATAGTTATACC-3') and COI-R (5'-TCCTAAAAAATGTTGAGGAAA-3'), targeting the COI gene (Zhao et al., 2014). PCR conditions were set according to Hossain et al. (2023). The amplified PCR products were analyzed by gel electrophoresis following the standard protocol described by the same author.

2.5 Statistical Analysis

An Excel spreadsheet was used to construct the database. Qualitative variables were presented as frequencies or percentages.

Pearson's Chi-square test or Fisher's exact test (for $n < 5$) was used to assess the association between each insect pest specimen and the host plants. All statistical procedures were performed using computer software, particularly the R statistical package. This comprehensive analytical approach ensured the reliability and

scientific rigor of the results, providing a detailed understanding of pest infestations and the identification of the species involved.

3. RESULTS

3.1 Morphological Identification of Collected Insect Pests

Morphological identification of the insects using a microscope allowed their classification into three groups (Table 1) : leafhoppers (jassids), aphids, and true bugs. Among the insects collected from okra and eggplant during this study, leafhoppers were the most frequently encountered, followed by true bugs, and lastly aphids.

Within the leafhoppers, three species were identified on both okra and eggplant : *Amrasca biguttula*, *Jacobiasca lybica*, and *Macrosteles cristatus* (Table 1). The aphid species identified was *Macrosiphum euphorbiae*, and the true bug species was *Macrolophus pygmaeus* (Table 1).

3.2 Molecular Identification of Collected Insect Pests


PCR amplification using the universal COI primer pair enabled the distinction between leafhoppers and other insect pests, such as bugs and aphids. As shown in Fig. 2, fragments of 654 bp correspond to leafhoppers, while fragments of 900 bp correspond to bugs or aphids.

3.3 Distribution According to Host Plants

The distribution of species according to host plants indicates that only one leafhopper species, *Amrasca biguttula*, was found on okra (Table 2). In contrast, all species collected in this survey were found on eggplant (Table 2). These include *Jacobiasca lybica* (leafhopper), which was the most abundant, followed by *A. biguttula* (leafhopper), *Macrolophus pygmaeus* (bug), *Macrosiphum euphorbiae* (aphid), and finally *Macrosteles cristatus* (leafhopper). Statistical analyses revealed a highly significant association between pest species and host plants (Table 2, $p < 0.001$). These results indicate that certain leafhoppers, such as *J. lybica* and *M. cristatus*, as well as aphids (*M. euphorbiae*) and bugs (*M. pygmaeus*), prefer feeding on eggplant rather than on okra.

Table 1. Diverse insect assemblages collected from okra and eggplant

Host plants	Insects collected	Species
Okra	Jassids	<i>Amrasca biguttula</i>
Eggplant	Jassids	<i>Macrosteles cristatus</i>
		<i>Jacobiasca lybica</i>
		<i>Amrasca biguttula</i>
	Aphid	<i>Macrosiphum euphorbiae</i>
	Bug	<i>Macrolophus pygmaeus</i>

**Fig. 2. Electrophoretic profile of PCR products for insect pest identification**

Lane 1 : 100 bp DNA ladder ; Lanes 2–4 and 8–9 : leafhoppers ; Lanes 5–7, 10, and 12 : other insect pests; Lane 11: negative control

Table 2. Distribution of leafhopper species according to host plants (okra and eggplant)

Host plants	<i>Amrasca biguttula</i> (%)	<i>Jacobiasca lybica</i> (%)	<i>Macrosteles cristatus</i> (%)	<i>Macrosiphum euphorbiae</i> (%)	<i>Macrolophus pygmaeus</i> (%)	χ^2 test (p-value)
Okra	539 (89,53)	0 (0)	0 (0)	0 (0)	0 (0)	< 0,001
Eggplant	63 (10,47)	89 (100)	1 (100)	8 (100)	41 (100)	
Total	602 (100)	89 (100)	1 (100)	8 (100)	41 (100)	

4. DISCUSSION

Morphological identification of insects collected from okra (*Abelmoschus esculentus*) and eggplant (*Solanum melongena*) leaves allowed the identification of three leafhopper species (*Amrasca biguttula*, *Jacobiasca lybica*, and *Macrosteles cristatus*), one bug species (*Macrolophus pygmaeus*), and one aphid species (*Macrosiphum euphorbiae*). The higher abundance of leafhoppers on okra and eggplant compared to bugs and aphids can be explained by several ecological and biological factors. Leafhoppers show a particular affinity for these crops, which serve as preferred hosts (Sahu et al., 2024; Tamang et al., 2025). Their piercing-

sucking mouthparts are adapted for phloem feeding and the high water and nutrient content of okra and eggplant leaves promotes their establishment and multiplication (Yarou et al., 2023).

In contrast, aphids, although polyphagous, exhibit a marked preference for other plant families, such as Brassicaceae or Cucurbitaceae, which limits their density on these crops (Ahmed et al., 2022). Phytophagous bugs preferentially exploit reproductive organs (fruits and seeds), and their abundance is highly seasonal (Ouaarous et al., 2025). Agro-climatic conditions also play a determining role. Leafhoppers develop optimally under warm and dry

conditions, typical of the tropical zones where okra and eggplant are grown (Gupta et al., 2024). Aphids, on the other hand, perform better under temperate and humid climates, explaining their lower density in these agroecosystems (Parajulee, 2007; Soh et al., 2018). Population dynamics may also play a central role. Leafhoppers have a short developmental cycle and high fecundity, allowing them to rapidly and massively colonize crops (Khifif et al., 2023). Thus, the numerical dominance of leafhoppers on okra and eggplant results from a combination of their physiological adaptation to the hosts, high ecological plasticity, rapid reproductive cycles, favorable climatic conditions, and lower pressure from natural enemies. Once considered secondary pests of these crops, leafhoppers have gradually become major pests over recent decades (Kouadio et al., 2022). *Amrasca biguttula* (Shiraki, 1913) was first reported in Africa in Ghana on okra as a minor pest (Kouadio et al., 2022). Favorable climatic conditions may explain its expansion in the sub-region, particularly in Côte d'Ivoire, as there are no geographical barriers for these pests. These results are consistent with those reported by Kouadio et al. (2022); however, among the three-leafhopper species identified by these authors, two differ from those found in this study, highlighting the colonization of multiple leafhopper species in the central-western region of Côte d'Ivoire. Similar observations have been made on cotton (Koné et al., 2017).

Molecular identification in this study allowed discrimination of leafhoppers from other insects collected on okra and eggplant leaves. Molecular identification is an essential modern tool for sustainable, knowledge-based agriculture, particularly for vulnerable crops like okra and eggplant. It enables the differentiation of closely related or cryptic species, which is sometimes impossible with morphological identification alone (Wang et al., 2018). This method prevents identification errors, especially at juvenile or larval stages. Indeed, studies have shown that immature stages frequently encountered in quarantine and plant protection activities require rapid and accurate species identification (Wang et al., 2018). Molecular tools are crucial in various pest management fields, including biological control, insecticide resistance management, prevention of invasive and exotic species introductions, and management of vector insects, including the identification of biotypes and cryptic species (Subhashini et al., 2025; Wang et al., 2018).

According to Subhashini et al. (2025), molecular approaches are essential for improving crop productivity and sustainability. However, in this study, molecular identification of the different leafhopper species and other pests was not conducted.

Statistical analyses revealed a highly significant association between leafhopper species and host plants, likely because each species shows a preference for a specific plant. For instance, this study showed that only *Amrasca biguttula* was found on okra, despite some samples being collected in fields containing both okra and eggplant. In addition to okra, this species was also found on eggplant along with the two other leafhopper species, aphids, and bugs identified in this study. This is explained by the polyphagous nature of leafhoppers (Kouadio et al., 2022), which attack a wide range of crops, unlike monophagous (single-host) or oligophagous (few closely related hosts) pests. Their ability to feed on diverse plants makes them particularly problematic in agriculture. Numerous studies have shown that leafhoppers are found on a wide variety of crops, including cotton (Koné et al., 2017; Subhashini et al., 2025) and horticultural crops (Al-Hamadany et al., 2017; Kouadio et al., 2022).

The presence of all pests collected in this study on eggplant may lead to damage to the plant and crop yield. Indeed, the co-occurrence of multiple pests on the same crop could lead to co-infestations that increase the virulence of attacks, as different species share similar biological and ecological traits, particularly phloem-sucking feeding behavior (Kumaraswamy et al., 2024).

5. CONCLUSION

This study identified three distinct groups of insect pests, with jassids being the most abundant. These insects are well known for their ability to cause substantial agricultural damage through sap feeding and the potential transmission of phytopathogens. Their simultaneous occurrence on both okra and eggplant highlights their broad nutritional adaptability and underscores the urgent need for improved phytosanitary management of these crops. The combined use of morphological and molecular approaches for pest identification proves essential for promoting sustainable, knowledge-based, and environmentally sound agriculture, particularly for vulnerable crops such as okra and eggplant.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

ACKNOWLEDGEMENTS

The authors thank the farmers who granted permission for the sampling conducted in their okra and eggplant fields.

COMPETING INTERESTS

Authors have declared that they have no known competing financial interests OR non-financial interests OR personal relationships that could have appeared to influence the work reported in this paper.

REFERENCES

- Ahmed, M. A., Ban, N., Hussain, S., Batool, R., Zhang, Y. J., Liu, T. X., & Cao, H. H. (2022). Preference and performance of the green peach aphid, *Myzus persicae*, on three Brassicaceae vegetable plants and its association with amino acids and glucosinolates. *PLoS ONE*, 17(12), e0269736. <https://doi.org/10.1371/journal.pone.0269736>
- Aikpon, R., Missihoun, A., Lokossou, A., Aikpon, G., Salifou, S., Dansi, A., & Agbangla, C. (2020). Genetic heterogeneity and resistance to insecticides in malaria vector *Anopheles gambiae* s.l. in the cotton growing area in Benin. *International Journal of Biological and Chemical Sciences*, 14(8), 2724–2736. <https://doi.org/10.4314/ijbcs.v14i8.6>
- Al-Hamadany, M. N., & Al-Karboli, H. H. (2017). First record of okra leafhopper, *Amrasca biguttula biguttula* Ishida, on okra in Iraq. *International Journal of Agricultural Technology*, 13(3), 393–402. <http://www.ijat-aatsea.com>
- Borkakati, R. N., Venkatesh, M. R., & Saikia, D. K. (2019). Insect pests of brinjal and their natural enemies. *Journal of Entomology and Zoology Studies*, 7(1), 932–937. <https://www.entomoljournal.com/archives/?ArticleId=4784&issue=1&vol=7&year=2019>
- Dmitriev, D. A., & Dietrich, C. H. (2009). 3I - An Interactive Key to Genera of Empoascini. In: List of 3I Interactive Keys and Taxonomic Databases. <http://dmitriev.speciesfile.org/key.asp?key=Erythroneura&lng=En&i=1&keyN=12>
- Ducroquet, H., Tillie, P., Elouhichi, K., & Gomez Y Paloma, S. (2017). A closer look at agriculture in Côte d'Ivoire: Overview of plant and animal production sectors and review of agricultural policies. Publications Office of the European Union. <https://doi.org/10.2760/126254>
- Food and Agriculture Organization of the United Nations, International Fund for Agricultural Development, United Nations Children's Fund, World Food Programme, & World Health Organization. (2018). The state of food security and nutrition in the world: Building climate resilience for food security and nutrition. <http://www.fao.org/3/I9553EN/i9553en.pdf>
- Gupta, S. K., Yadav, A. K., Maheshwari, S., Maurya, P. K., & Kumar, P. (2024). Population dynamics of major insect pests of okra in relation to weather parameters. *Plant Archives*, 24(1), 304–308. <https://doi.org/10.51470/PLANTARCHIVES.2024.v24.no.1.043>
- Hossain, S., Rahman, M. M., Khan, H. I., Haque, M. A., Jannat, R., & Hassan, J. (2023). Molecular characterization of cotton jassid and their response against relatively newer pesticides in two cotton varieties of Bangladesh. *Malaysian Journal of Sustainable Agriculture*, 7(2), 79–87. <https://doi.org/10.26480/mjsa.02.2023.79.87>
- Khfif, K., Labaioui, Z., Koledenkova, K., Mokri, F., Sbaghi, M., Zaid, A., Brostaux, Y., & El Rhaffari, L. (2023). Population dynamics of the leafhopper *Jacobiasca lybica* (Hemiptera: Cicadellidae) within vineyards and citrus orchards of Morocco. *International Journal of Pest Management*, 71(4), 330–342. <https://doi.org/10.1080/09670874.2023.2206372>
- Konan, N. O., Akaffou, M. A., Kouadio, L., Akaffou, D. S., & Mergeai, G. (2020). Genetic diversity of exotic and local eggplants (*Solanum* spp.) cultivated in Côte d'Ivoire based on ISSR markers. *Biodiversitas*, 21(8), 3651–3658. <https://doi.org/10.13057/biodiv/d210830>
- Koné, K., Tuo, Y., Cissé, M., & Koua, H. K. (2019). Essai de Lutte Contre les

- Principaux ravageurs de la courgette par l'utilisation d'extraits de *Ocimum canum* Sims, à Korhogo dans le Nord de la Côte d'Ivoire. *European Scientific Journal*, 15(24), 72–89. <https://doi.org/10.19044/esj.2019.v15n24p72>
- Koné, P. W. E., Ochou, G. E. C., Didi, G. J. R., Dekoula, S. C., Kouakou, M., Bini, K. K. N., Mamadou, D., & Ochou, G. (2017). Spatial and temporal evolution of damage caused by the jassid *Jacobiella facialis* Jacobi, 1912 (Cicadellidae) in comparison with rainfall distribution over the past twenty years in cotton-growing areas of Côte d'Ivoire. *International Journal of Biological and Chemical Sciences*, 11(3), 1190–1201. <https://doi.org/10.4314/ijbcs.v11i3.21>
- Kouadio, H., Kouakou, M., Bini, K., Ouattara, M., Adepo-Gourène, B., & Ochou, O. (2022). Diagnosis of jassid attacks on okra and eggplant plots in the center and center-west of Côte d'Ivoire. *International Journal of Innovation and Applied Studies*, 38(2), 408–416. <http://www.ijias.issr-journals.org/>
- Kumaraswamy, S., Nasif, S. O., Siddique, A. B., Karuppannasamy, A., Chowdary, R., Mohan, M., & Chaitanya, A. K. (2024). Host Plant Resistance to Insects in Vegetable Crops. In S. Kumar & M. Furlong (Eds.), *Plant Resistance to Insects in Major Field Crops* (pp. 215-248). Springer Nature. https://doi.org/10.1007/978-981-99-7520-4_9
- Ouaarous, M., El Fakhouri, K., Taarji, N., Baouchi, A., Amri, M., Ramdani, C., Sobeh, M., Mesfioui, A., & El Bouhssini, M. (2025). Impact of field insect pests on seed and nutritional quality of some important crops: A comprehensive review. *ACS Omega*, 10(9), 8779–8792. <https://doi.org/10.1021/acsomega.4c08982>
- Parajulee, M. N. (2007). Influence of constant temperatures on life history parameters of the cotton aphid, *Aphis gossypii*, infesting cotton. *Environmental Entomology*, 36(4), 666–672. [https://doi.org/10.1603/0046-225X\(2007\)36\[666:IOCTOL\]2.0.CO;2](https://doi.org/10.1603/0046-225X(2007)36[666:IOCTOL]2.0.CO;2)
- Prabhakar, M., Rao, M. S., Timmanna, H., Prasad, T. V., & Vinod Kumar, N. (2024). Climate change and its impact on pests of dryland crops. *Indian Journal of Dryland Agricultural Research and Development*, 39(2), 121–127. <https://doi.org/10.5958/2231-6701.2024.00025.7>
- Sahu, S., Bhagat, P. K., Painkra, G. P., Painkra, K. L., Sahu, R., Jaiswal, V. K., & Miri, Y. (2024). Evaluate the efficacy of biorational pesticides against the key pests of okra, *Abelmoschus esculentus* (L.), at Northern hills of Chhattisgarh. *International Journal of Advanced Biochemistry Research*, 8(8), 542–547. <https://doi.org/10.33545/26174693.2024.v8.i8g.1790>
- Seehausen, M. L., Ris, N., Driss, L., Racca, A., Girod, P., Warot, S., Borowiec, N., Tosevski, I., & Kenis, M. (2020). Evidence for a cryptic parasitoid species reveals its suitability as a biological control agent. *Scientific Reports*, 10, 19096. <https://doi.org/10.1038/s41598-020-76180-5>
- Sène, S. O., Tendeng, E., Diatte, M., Sylla, S., Labou, B., Diallo, A. W., & Diarra, K. (2020). Insecticide resistance in field populations of the tomato fruitworm, *Helicoverpa armigera*, from Senegal. *International Journal of Biological and Chemical Sciences*, 14(1), 181-191. <https://doi.org/10.4314/ijbcs.v14i1.15>
- Shiraki, T. (1913). Report on the injurious insects of Formosa. Special Report of the Agricultural Experiment Station of Formosa, 8, 1-670.
- Soh, B. S. B., Kekeunou, S., Nanga Nanga, S., Dongmo, M., & Hanna, R. (2018). Effect of temperature on the biological parameters of the cabbage aphid *Brevicoryne brassicae*. *Ecology and Evolution* <https://doi.org/10.1002/ece3.4639>
- Soro, G., Koffi, N. M., Koné, B., Kouakou, Y. E., M'Bra, K. R., Soro, P. D., & Soro, N. (2018). Utilisation de produits phytosanitaires dans le maraîchage autour du barrage d'alimentation en eau potable de la ville de Korhogo (nord de la Côte d'Ivoire): risques pour la santé publique. *Environnement, Risques & Santé*, 17(2), 155–163. <https://doi.org/10.1684/ers.2018.1147>
- Subhashini, S., Keerthivarman, K., Rajeswari, S., Premalatha, N., Kalaimagal, T., Muthuswami, M., & Jeyakumar, P. (2025). Molecular, cytological and morphological studies on jassid resistance in cotton (*Gossypium hirsutum* L.) based on hairiness trait. *Journal of Cotton Research* <https://doi.org/10.1186/s42397-025-00217-1>
- Tamang, A., Chaudhary, P., Acharya, M., Pokharel, M., & Soti, A. (2025). Field

- efficacy of different insecticides against jassid (*Amrasca biguttula biguttula*) in brinjal at Gauradaha, Jhapa, Nepal. *Journal of Agriculture and Applied Biology*. <https://doi.org/10.11594/jaab.06.02.01>
- Tuo, S., Koné, D.S.F., Guinagui, N.B., Sanogo, S., Bamba, B., Yéo, G., Nomel, M.P., Anoman, O.J.A., & Koné, D. (2023). Assessment of the Agronomic Performance of Seven Okra (*Abelmoschus esculentus* (L.) Moench) Cultivars in Southern Côte d'Ivoire. *World Journal of Agricultural Research*, 11(4), 87-97. <https://doi.org/10.12691/wjar-11-4-1>
- Wang, H., Kennedy, G. G., Reay-Jones, F. P. F., Reisig, D. D., Toews, M. D., Roberts, P. M., Herbert, D. A., Jr., Taylor, S., Jacobson, A. L., & Greene, J. K. (2018). Molecular identification of thrips species infesting cotton in the southeastern United States. *Journal of Economic Entomology*, 111(2), 892–898. <https://doi.org/10.1093/jee/toy036>
- Widadie, F., Wulandari, E., & Lestari, R. D. (2024). Business models for engaging smallholder farmers in high-value markets: empirical evidence from vegetable value chains. *Agricultural and Resource Economics: International Scientific E-Journal*, 10(4), 179–202. <https://doi.org/10.51599/are.2024.10.04.08>
- Yarou, B. B., Francisco, A. R., De Troij, A., & Aboubakar Souana, D. (2023). *Amrasca biguttula* Ishida : Quel avenir pour la production du gombo au Bénin ?. *World Vegetable Center*. <https://agritrop.cirad.fr/607265/2/607265.pdf>
- Zanzana, K., Sinzogan, A., Tapa-Yotto, G. T., Dannon, E., Goergen, G., & Tamò, M. (2025). Seasonal and spatial distribution of fall armyworm larvae in maize fields: Implications for integrated pest management. *Insects*, 16(2), 1–17. <https://doi.org/10.3390/insects16020145>
- Zhao, W., Tan, K., Zhou, D., Wang, M., Cheng, C., Yu, Z., Miao, Y., & He, S. (2014). Phylogeographic analysis of *Apis cerana* populations on Hainan Island and southern mainland China, based on mitochondrial DNA sequences. *Apidologie*, 45(1), 21-33. <https://doi.org/10.1007/s13592-013-0223-8>

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://pr.sdiarticle5.com/review-history/149450>