



# Role of Sulphur and Molybdenum in Improving Nutritional Quality and Profitability of Cowpea (*Vigna unguiculata* L.)

Rahul R. Pisal<sup>a++</sup>, Kajal V. Vala<sup>a#</sup>, P. Sowjanya Deepthi<sup>at</sup>  
and Payal A. Patel<sup>at\*</sup>

<sup>a</sup> Department of Agronomy, N.M. College of Agriculture, Navsari Agricultural University, Navsari - 396 450, Gujarat, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/ijpss/2025/v37i125915>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://pr.sdiarticle5.com/review-history/149897>

Original Research Article

Received: 15/10/2025  
Published: 31/12/2025

## ABSTRACT

A field experiment was carried out in the summer of 2020 at the College Farm of N. M. College of Agriculture, Navsari Agricultural University, located in Navsari, Gujarat. The objective was to assess the role of sulphur and molybdenum in improving nutritional quality and profitability of cowpea (*Vigna unguiculata* L.) The soil at the experimental site was clayey in texture and slightly alkaline in reaction. The study included nine different treatment combinations. T<sub>1</sub> - RDF, T<sub>2</sub> - 25 kg S/ha, T<sub>3</sub> - 50 kg S/ha, T<sub>4</sub> -500 ppm foliar Mo, T<sub>5</sub> -0.5 g Mo/kg seed, T<sub>6</sub> - 25 kg S + 500 ppm Mo, T<sub>7</sub> - 25 kg S + 0.5 g Mo/kg seed, T<sub>8</sub> - 50 kg S + 500 ppm Mo and T<sub>9</sub> -50 kg S + 0.5 g Mo/kg seed. All treatments

<sup>++</sup> Assistant Professor; <sup>#</sup> M.sc Student; <sup>†</sup> Ph.D Scholar;  
<sup>\*</sup>Corresponding author: E-mail: [payal12062001@gmail.com](mailto:payal12062001@gmail.com);

were uniformly supplied with the recommended dose of fertilizers (RDF) at 20-40-00 kg NPK/ha. Sulphur was incorporated into the soil at the time of sowing, whereas molybdenum was applied as a foliar spray at 30 days after sowing (DAS) using 250 L of water per hectare. The experiment followed a randomized block design (RBD) with three replications. Among the treatments, T<sub>6</sub> (25 kg S + 500 ppm Mo) recorded a significant improvement in quality parameters, as well as in the nitrogen, phosphorus and potassium content and uptake by both seed and haulm. Post-harvest soil analysis revealed that the availability of nitrogen, sulphur and molybdenum was influenced by the different sulphur and molybdenum treatments. However, soil phosphorus and potassium levels after harvest were not significantly affected by any of the treatments.

**Keywords:** Cowpea; economics; fertilization; content and uptake; molybdenum; sulphur.

## 1. INTRODUCTION

Cowpea, sometimes called *lobia* (*Vigna unguiculata* L.), is a multipurpose legume that is used for grain, vegetables, fodder and to increase soil fertility. Due to its ability to withstand drought and its minimal input requirements, cowpea is primarily cultivated in arid and semi-arid regions and plays a vital role as a crop during the kharif and summer seasons in India (Reddy et al., 2023; Banerjee et al., 2021). Similar to other pulse crops, cowpea enhances soil fertility through biological nitrogen fixation and the incorporation of organic matter, facilitated by *Rhizobium leguminosarum* (Xavier et al., 2023). Despite its agronomic advantages, India continues to experience a shortfall in pulse production. According to the Ministry of Agriculture, total pulse output was estimated at 234 lakh tonnes in 2023–2024, down from 261 lakh tonnes in 2022–2023. To bridge the gap between supply and demand, India has been importing an average of 25 lakh tonnes of pulses annually, with forecasts indicating over 4.5 million tonnes of tur, urad and masoor imports in 2024. In response, the government has extended the duty-free import policy for these pulses until 2025 to help stabilize domestic availability and control prices (Anon., 2023).

Cowpea, known for its high nutritional value and extensive cultivation in tropical and subtropical regions, plays a significant role in ensuring food and nutritional security. It also contributes to improved soil fertility through atmospheric nitrogen fixation (Silva et al., 2025). Despite its potential, cowpea productivity in India remains sub optimal, primarily due to imbalanced fertilization and widespread deficiencies of key secondary and micronutrients particularly sulphur (S) and molybdenum (Mo). Sulphur is an essential secondary nutrient involved in protein synthesis, chlorophyll production and various enzymatic activities. Molybdenum, on the other

hand, functions as a crucial cofactor for enzymes like nitrogenase and nitrate reductase, thereby influencing nodulation and nitrogen fixation processes. Proper and balanced application of these nutrients can significantly boost legume yield and overall productivity (Dhaliwal et al., 2022).

In this context, the present study was undertaken to evaluate the effects of sulphur and molybdenum, applied individually and in combination through soil application, seed treatment and foliar spray on the yield, nutrient content, uptake and economics of cowpea under the agro-climatic conditions of south Gujarat.

## 2. MATERIALS AND METHODS

A field experiment was conducted during the summer season of 2020 at the College Farm, N. M. College of Agriculture, Navsari Agricultural University, Navsari, to study the influence of sulphur and molybdenum fertilization on cowpea (*Vigna unguiculata* L.). The experimental soil was clayey in texture with a slightly alkaline reaction. The study comprised nine treatment combinations: T<sub>1</sub> – RDF, T<sub>2</sub> – 25 kg S/ha, T<sub>3</sub> – 50 kg S/ha, T<sub>4</sub> – 500 ppm foliar Mo, T<sub>5</sub> – 0.5 g Mo/kg seed, T<sub>6</sub> – 25 kg S + 500 ppm Mo, T<sub>7</sub> – 25 kg S + 0.5 g Mo/kg seed, T<sub>8</sub> – 50 kg S + 500 ppm Mo, and T<sub>9</sub> – 50 kg S + 0.5 g Mo/kg seed. All treatments received a common dose of 20:40:00 NPK kg/ha. Sulphur was applied at sowing, while foliar application of molybdenum (500 ppm) was carried out at 30 DAS using 250 L/ha of water.

The experiment was conducted using a randomized block design (RBD) with three replications. Land preparation involved ploughing, harrowing and planking to achieve a fine tilth suitable for sowing. Cowpea variety GC-4 was planted on 18<sup>th</sup> February 2020 at a seed rate of 25 kg/ha, with a sowing depth of 3–4 cm

and spacing of 45 × 10 cm. To ensure uniform germination, irrigation was applied immediately after sowing. The required amounts of farmyard manure (FYM) and bio-compost were calculated based on the gross plot area. Urea, single superphosphate, gypsum and ammonium molybdate were weighed according to the treatment specifications, thoroughly mixed with the organic amendments and manually applied in pre-opened furrows. Weed control was carried out using standard herbicide practices, and plant protection measures were implemented as needed. The crop was harvested on 10 May 2020.

The nitrogen content in cowpea seeds was estimated using the micro-Kjeldahl method, as described by Jackson (1973). Chemical analyses for nitrogen, phosphorus and potassium content in both seed and haulm, as well as their uptake and the post-harvest available N, P and K status of the soil, were conducted using standard procedures. Nitrogen was determined using the modified Kjeldahl method, phosphorus by the wet digestion (diacid) method followed by the vanado-molybdo phosphoric acid yellow colorimetric technique and potassium through the flame photometric method (Jackson, 1973). Available sulphur in the soil was analyzed by extracting the sample with 1% NaCl and estimating it using the turbidimetric method (Williams and Steinbergs, 1959). Available molybdenum was extracted using acid ammonium oxalate and estimated by the colorimetric method (Chapman and Pratt, 1961). Experimental data were subjected to statistical analysis as per the procedures outlined by Panse and Sukhatme (1985). Analysis of variance (ANOVA) was used to assess the significance of treatment effects on cowpea performance. Protein content (%), protein yield (kg/ha), and nutrient uptake by seed and haulm were calculated using the following formulas:

$$\text{Protein content (\%)} = \text{Nitrogen content (\%)} \times 6.25$$

$$\text{Protein yield (kg/ha)} = \frac{\text{Protein content in grain (\%)} \times \text{Grain yield (kg/ha)}}{100}$$

$$\text{Uptake (kg/ha)} = \frac{\text{Content (\%)} \times \text{Yield(kg/ha)}}{100}$$

### 3. RESULTS AND DISCUSSION

**Quality parameters:** Seed yield (928 kg/ha) and haulm yield (2280 kg/ha) were also significantly increased with T<sub>6</sub> (25 kg S + 500 ppm Mo), but were statistically comparable to yields under T<sub>2</sub>,

T<sub>4</sub>, T<sub>7</sub>, T<sub>8</sub> and T<sub>9</sub>. Based on the data provided in Table 1, the magnitude of the protein content of cowpea seed indicated that different treatments of sulphur and molybdenum fertilization improve protein content significantly. However, treatment T<sub>6</sub> (25 kg S + 500 ppm Mo) contributed to higher protein content (24.01 %) and statistically at par with treatment T<sub>2</sub>, T<sub>4</sub>, T<sub>5</sub>, T<sub>7</sub>, T<sub>8</sub> and T<sub>9</sub>. Significantly higher protein yield of cowpea (223.24 kg/ha) was recorded under treatment of T<sub>6</sub> and was at par with the treatments of T<sub>7</sub>, T<sub>8</sub> and T<sub>9</sub>.

Sulphur plays an important role in the synthesis of amino acids, proteins, vitamins, and coenzymes. Recent studies indicate that lack of sulphur to meet plant requirements may reduce not only yields but also the quality of grain legumes (reduction of cysteine and methionine) by changing the gene expression of storage proteins in developing seeds. Molybdenum is also important for sulphur metabolism, so it can have a positive impact on the biological value of protein (Głowacka et al. (2019).

#### 3.1 N, P and K Content in Cowpea Seed and Haulm

**Nitrogen content in cowpea (%):** Data revealed that the highest nitrogen content in seed (4.46%) was recorded under treatment T<sub>6</sub>, which was statistically at par with T<sub>7</sub>, while the lowest (2.35%) was observed under T<sub>1</sub>. Similarly, nitrogen content in haulm was significantly higher with T<sub>6</sub> (0.70%), closely followed by T<sub>7</sub> and T<sub>8</sub>, whereas the minimum (0.57%) was noted under T<sub>1</sub>.

**P<sub>2</sub>O<sub>5</sub> content in cowpea (%):** Data presented that P<sub>2</sub>O<sub>5</sub> content in seed was found significantly higher under the treatment T<sub>6</sub> (2.10%) but remained at par with treatments T<sub>7</sub>, T<sub>8</sub>, T<sub>9</sub> and T<sub>2</sub>. While in case of haulm, significantly higher P<sub>2</sub>O<sub>5</sub> content in haulm was recorded with treatment T<sub>6</sub> (0.26%) but remained at par with T<sub>4</sub>, T<sub>5</sub>, T<sub>7</sub>, T<sub>8</sub> and T<sub>9</sub>. While the lowest P<sub>2</sub>O<sub>5</sub> content in seed (1.47%) and haulm (0.19%) registered under T<sub>1</sub> (RDF).

**K<sub>2</sub>O content in cowpea (%):** K<sub>2</sub>O content in seed was found significantly higher under the treatment T<sub>6</sub> (1.35%) but remained at par with treatments T<sub>7</sub>, T<sub>8</sub>, T<sub>9</sub> and T<sub>4</sub>. While there was no significant different found in case of K<sub>2</sub>O content in haulm.

This increase might be due to the fact that sulphur and molybdenum being essential plant nutrients promote nodulation, better root formation being an essential plant nutrient which in turn activated higher absorption of nitrogen, phosphorus and potash from soil. The data present in Table 1 revealed that significantly higher content of N, P<sub>2</sub>O<sub>5</sub>, K<sub>2</sub>O were observed under T<sub>6</sub> (25 kg S + 500 ppm Mo) while lower content found in T<sub>1</sub> (RDF). Similar findings were also reported by Venkatesh *et al.* (2006) and Reddy *et al.* (2023).

### 3.2 N, P and K Uptake in Cowpea Seed and Haulm

**Nitrogen uptake by cowpea (kg/ha):** Significantly the highest nitrogen uptake by seed was recorded with treatment T<sub>6</sub> (41.19 kg/ha) but remained at par with treatments T<sub>7</sub> (37.94 kg/ha). In the case of haulm, significantly higher nitrogen uptake by haulm was recorded with treatment T<sub>6</sub> (16 kg/ha), which remained at par with T<sub>7</sub> and T<sub>8</sub>. The lowest nitrogen uptake by haulm registered under T<sub>1</sub> (10.46 kg/ha).

**P<sub>2</sub>O<sub>5</sub> uptake by cowpea (kg/ha):** P<sub>2</sub>O<sub>5</sub> uptake by seed (19.47 kg/ha) and haulm (5.94 kg/ha) was significantly higher under treatment T<sub>6</sub> but remained statistically at par with treatments T<sub>7</sub>, T<sub>8</sub>, and T<sub>9</sub>.

**K<sub>2</sub>O uptake by cowpea (kg/ha):** K<sub>2</sub>O uptake by seed was significantly affected by different levels of sulphur and molybdenum fertilization. Significantly the highest K<sub>2</sub>O uptake by seed (12.52 kg/ha) and haulm (20.12 kg/ha) was recorded with T<sub>6</sub> treatment and remained at par with treatments T<sub>7</sub>, T<sub>8</sub> and T<sub>9</sub>.

It might be due to the fact that sulphur and molybdenum play vital role in growth and yield of cowpea and also increase N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O content in cowpea. Hence, it is obvious fact that application of sulphur and molybdenum improved overall uptake of N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O by seed and haulm. Similar findings were also reported by Venkatesh *et al.* (2006) and Reddy *et al.* (2023).

### 3.3 Soil Property Status

The mean data pertaining to available N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O status after harvest of cowpea are presented in Table 3.

**Available N in soil after harvest (kg/ha):** Available N in soil was significantly highest recorded with treatment T<sub>7</sub> (266.82 kg/ha). While

significantly lower soil N was recorded in treatment T<sub>1</sub> (RDF).

**Available P<sub>2</sub>O<sub>5</sub> in soil after harvest (kg/ha):** Non-significant effect of different treatments of sulphur and molybdenum fertilization treatments on available phosphorus of soil after harvest of cowpea. Data indicated that P<sub>2</sub>O<sub>5</sub> content in soil after harvest was found non-significant. The highest P<sub>2</sub>O<sub>5</sub> content in soil after harvest was recorded in treatment T<sub>3</sub> (44.61 kg/ha) while lowest registered in treatment T<sub>1</sub> (41.04 kg/ha).

**Available K<sub>2</sub>O in soil after harvest (kg/ha):** The data indicated that non-significantly the highest available K<sub>2</sub>O of soil after harvest of the crop was recorded from treatment T<sub>8</sub> (344.29 kg/ha) while lowest registered in treatment T<sub>7</sub> (335.09 kg/ha).

Similar result also reported by Reddy *et al.* (2023) and Saiyad *et al.* (2017). This might be due to molybdenum unique role in enhancing nitrogen fixation. Application of S and Mo fertilization treatments in available P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O in the soil after harvest of crop were found non significant.

**Available Sulphur in soil after harvest (ppm):** Data indicated that available sulphur in soil was significantly affected by different levels of sulphur and molybdenum fertilization. Significantly the highest available sulphur was recorded with T<sub>8</sub> (10.54 ppm) and remained at par with treatments T<sub>9</sub>, T<sub>7</sub>, T<sub>2</sub> and T<sub>3</sub>.

**Available molybdenum in soil after harvest (ppm):** Available molybdenum in soil was significantly affected by different levels of sulphur and molybdenum fertilization. Significantly the highest available molybdenum was recorded with T<sub>5</sub> (0.047 ppm) and remained at par with treatments T<sub>4</sub>, T<sub>6</sub> and T<sub>7</sub>.

The increased S and Mo content might be due to increased concentration of sulphur and molybdenum in soil solution with application of increasing levels of sulphur and molybdenum Kumar and Umesha (2023), Nehra *et al.* (2024) and Markam *et al.* (2025).

**Economics:** The data on economics pertaining to gross return, net return and benefit cost ratio (BCR) under different sulphur and molybdenum fertilization treatments The data revealed that maximum gross return of Rs. 55521.33/ha was realized with T<sub>6</sub> (25 kg S + 500 ppm Mo),

**Table 1. NPK content in seed and haulm of Cowpea as influenced by different levels of sulphur and molybdenum treatments**

Treatments	Seed yield (kg/ha)	Haulm yield (kg/ha)	Quality parameters		N content (%)		P content (%)		K content (%)	
			Protein content (%)	Protein yield (kg/ha)	Seed	Haulm	Seed	Haulm	Seed	Haulm
T <sub>1</sub> : RDF	785	1843	18.29	143.22	2.35	0.57	1.47	0.19	1.11	0.74
T <sub>2</sub> : 25 kg S/ha	861	2020	21.53	185.39	2.92	0.59	1.86	0.23	1.24	0.84
T <sub>3</sub> : 50 kg S/ha	821	1916	21.10	173.14	2.42	0.58	1.57	0.20	1.18	0.76
T <sub>4</sub> : 500 ppm foliar Mo	852	2019	21.39	182.26	3.13	0.61	1.75	0.22	1.22	0.83
T <sub>5</sub> : 0.5 g Mo/kg seeds	832	1966	21.19	175.78	3.04	0.59	1.61	0.22	1.20	0.80
T <sub>6</sub> : 25 kg S + 500 ppm Mo	928	2280	24.01	223.24	4.46	0.70	2.10	0.26	1.35	0.88
T <sub>7</sub> : 25 kg S + 0.5 g Mo/kg seeds	916	2230	23.73	217.44	4.14	0.65	2.01	0.25	1.32	0.87
T <sub>8</sub> : 50 kg S + 500 ppm Mo	902	2206	22.71	204.75	3.56	0.63	1.95	0.24	1.28	0.86
T <sub>9</sub> : 50 kg S + 0.5g Mo/kg seeds	900	2174	13.39	202.80	3.29	0.62	1.91	0.24	1.26	0.85
S.Em. ±	30.03	88.43	0.93	9.82	0.17	0.024	0.11	0.014	0.045	0.041
C.D at 5 %	90.05	265.12	2.81	29.47	0.52	0.07	0.33	0.04	0.14	NS
C.V. %	6.00	7.38	7.45	8.97	9.15	6.71	10.44	10.89	6.33	8.56

**Table 2. NPK uptake by seed and haulm of Cowpea as influenced by different levels of sulphur and molybdenum treatments**

Treatments	N uptake (kg/ha)		P uptake (kg/ha)		K uptake (kg/ha)	
	Seed	Haulm	Seed	Haulm	Seed	Haulm
T <sub>1</sub> : RDF	18.42	10.46	11.64	3.50	8.79	13.58
T <sub>2</sub> : 25 kg S/ha	25.17	11.90	16.02	4.64	10.67	16.98
T <sub>3</sub> : 50 kg S/ha	19.90	11.09	12.84	3.78	9.66	14.55
T <sub>4</sub> : 500 ppm foliar Mo	26.65	12.33	14.91	4.44	10.41	16.75
T <sub>5</sub> : 0.5 g Mo/kg seeds	25.33	11.66	13.43	4.31	10.02	15.77
T <sub>6</sub> : 25 kg S + 500 ppm Mo	41.19	16.00	19.47	5.94	12.57	20.12
T <sub>7</sub> : 25 kg S + 0.5 g Mo/kg seeds	37.94	14.40	18.43	5.57	12.05	19.39
T <sub>8</sub> : 50 kg S + 500 ppm Mo	32.09	13.91	17.61	5.36	11.54	19.00
T <sub>9</sub> : 50 kg S + 0.5g Mo/kg seeds	29.63	13.39	17.16	5.22	11.31	18.49
S.Em. ±	1.64	0.86	1.13	0.36	0.57	1.04
C.D at 5 %	4.92	2.58	3.39	1.09	1.72	3.14
C.V. %	9.98	11.64	12.47	13.21	9.22	10.56

**Table 3. Available N, P, K, S and Mo in soil after harvest of cowpea as influenced by different levels of sulphur and molybdenum treatments**

Treatments	Available N (kg/ha)	Available P <sub>2</sub> O <sub>5</sub> (kg/ha)	Available K <sub>2</sub> O (kg/ha)	Available S (ppm)	Available Mo (ppm)	Returns (₹/ha)		BCR
						Gross return	Net return	
T <sub>1</sub> : RDF	201.53	41.04	339.94	7.01	0.034	46621	28952	2.64
T <sub>2</sub> : 25 kg S/ha	216.47	41.76	342.21	8.85	0.036	51131	33234	2.86
T <sub>3</sub> : 50 kg S/ha	227.43	44.61	338.57	10.19	0.037	48731	30606	2.69
T <sub>4</sub> : 500 ppm foliar Mo	223.29	42.89	338.09	7.48	0.045	50658	32277	2.76
T <sub>5</sub> : 0.5 g Mo/kg seeds	225.73	42.74	339.04	7.19	0.047	49465	31522	2.76
T <sub>6</sub> : 25 kg S + 500 ppm Mo	233.81	42.95	338.17	8.55	0.043	55521	36912	2.98
T <sub>7</sub> : 25 kg S + 0.5 g Mo/kg seeds	266.82	43.20	335.02	9.48	0.044	54755	36584	3.01
T <sub>8</sub> : 50 kg S + 500 ppm Mo	228.31	42.48	344.29	10.54	0.040	53924	35087	2.86
T <sub>9</sub> : 50 kg S + 0.5g Mo/kg seeds	232.18	41.91	338.81	9.99	0.041	53678	35279	2.92
S.Em. ±	8.20	1.85	18.02	0.65	0.0016			
C.D at 5 %	24.58	NS	NS	1.94	0.0047			
C.V. %	6.21	7.53	9.19	12.70	6.60			
Initial status	220.12	39.50	340.25	7.86	0.035			

followed by the treatments T<sub>7</sub>, T<sub>8</sub>, T<sub>9</sub> and T<sub>2</sub> and maximum net return of Rs. 36912/ha was realized with T<sub>6</sub> (25 kg S + 500 ppm Mo) followed by the treatment T<sub>7</sub> (25 kg S + 0.5 g Mo/kg seeds) by recording net return of Rs. 36584/ha. An appraisal of data showed that maximum BCR of 3.01 was obtained with the T<sub>7</sub> (25 kg S + 0.5 g Mo/kg seeds), followed by the treatment T<sub>6</sub> (25 kg S + 500 ppm Mo) by securing BCR of 2.98.

#### 4. CONCLUSION

Until quite recently, reviews and studies confirm the positive effect of sulphur and molybdenum for a significant improve on the nutritional quality and yield of cowpea. These nutrients enhance protein content and support essential metabolic functions in the plant. The combined application of RDF with 25 kg S/ha + 500 ppm foliar Mo (T<sub>6</sub>) significantly enhanced the N, P, and K content and uptake by seed and haulm. Post-harvest soil analysis indicated that available nitrogen, sulphur, and molybdenum status were positively influenced by sulphur and molybdenum fertilization. Hence, integrated nutrient management involving sulphur and molybdenum, particularly through seed treatment, can be recommended for improving cowpea productivity and soil fertility under south Gujarat conditions.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

#### COMPETING INTERESTS

Authors have declared that they have no known competing financial interests or non-financial interests or personal relationships that could have appeared to influence the work reported in this paper.

#### REFERENCES

- Anonymous. (2023). *Annual report 2022–23*. Government of India, Ministry of Agriculture and Farmer Welfare, Directorate of Pulses Development. <https://dpd.gov.in/DPD/Pub./TR/46/2022-23.pdf>
- Banerjee, P., Das, P., & Sinha, S. (2021). Importance of molybdenum for the production of pulse crops in India. *Journal of Plant Nutrition*, 45(2), 300–310. <https://doi.org/10.1080/01904167.2021.2020831>
- Chapman, H. D., & Pratt, P. F. (1961). *Methods of analysis for soils, plants, and waters*. University of California, Division of Agricultural Sciences.
- Dhaliwal, S. S., Sharma, V., Shukla, A. K., Kaur, J., Verma, V., Kaur, M., & Hossain, A. (2022). Interactive effects of molybdenum, zinc and iron on the grain yield, quality, and nodulation of cowpea (*Vigna unguiculata* (L.) Walp.) in North-Western India. *Molecules*, 27(11), 3622. <https://doi.org/10.3390/molecules27113622>
- Głowacka, A., Gruszecki, T., Bogdan Szostak, B. and Michalek, S. (2019). The response of common bean to sulphur and molybdenum fertilization. *International Journal of Agronomy*, <https://doi.org/10.1155/2019/3830712>
- Jackson, M. L. (1973). *Soil chemical analysis*. Prentice-Hall of India Pvt. Ltd.
- Kumar, A. B., & Umesha, C. (2023). Influence of molybdenum and foliar application of boron on growth and yield of cowpea (*Vigna unguiculata*). *Journal of Experimental Agriculture International*, 45(8), 41–48. <https://doi.org/10.9734/JEAI/2023/v45i82153>
- Markam, A., Swaroop, N., & Dhruv, S. S. (2025). Response of different levels of NPK and molybdenum on growth and yield attributes of black gram (*Vigna mungo* L.) var. Shekhar-2. *International Journal of Research in Agronomy*, SP-8(1), 254–258. <https://doi.org/10.33545/2618060X.2025.v8.i1Sd.2426>
- Nehra, A., Singh, V., & Kumar, A. (2024). Effect of sulphur and molybdenum on the growth and yield of cowpea. *International Journal of Research in Agronomy*, SP-7(6), 427–429. <https://doi.org/10.33545/2618060X.2024.v7.i6Sf.896>
- Panase, V. G., & Sukhatme, P. V. (1985). *Statistical methods for agricultural workers*. Indian Council of Agricultural Research.
- Reddy, K. R. T. K., Debbarna, V., & Reddy, S. P. K. (2023). Influence of sulphur and foliar application of molybdenum on growth and yield of summer cowpea (*Vigna unguiculata* L.). *International Journal of Environment and Climate Change*, 13(6), 259–267.

- <https://doi.org/10.9734/ijecc/2023/v13i61821>
- Reddy, K., Debbarma, V. & Reddy, S. P. K. (2023). Influence of Sulphur and Foliar Application of Molybdenum on Growth and Yield of Summer Cowpea (*Vigna unguiculata* L.). *International Journal of Environment and Climate Change*, 13(6). DOI: 10.9734/ijecc/2023/v13i61821
- Saiyad, N. M., Chaudhary, N. N., Thounaojam, A. S., & Mistry, G. J. (2017). Effect of sulfur and molybdenum on quality of kharif forage cowpea (*Vigna unguiculata* (L.) Walp.). *Environment & Ecology*, 35(3A), 1883–1888.  
<https://environmentandecology.com/wp-content/uploads/2025/03/MS17-1-8.pdf>  
ISSN 0970-0420
- Silva, N. S. G. D., Oliveira, A. C. D., Júnior, B. C., Aviz, R. O. D., Pereira, K. P., Vieira, D. S. M. D. M., & Souza, E. J. O. D. (2025). Can molybdenum fertilization enhance protein content and digestibility of sorghum single cropped and intercropped with cowpea? *Grasses*, 4(3), 28.  
<https://doi.org/10.3390/grasses4030028>
- Venkatesh, M. S., Majumdar, B., & Kumar, K. (2006). Response of groundnut, *Arachis hypogaea* to sulphur, boron and FYM doses in an Ultic Hapludalf of Meghalaya. *Journal of Oilseeds Research*, 23(1), 52.
- Williams, C. H., & Steinbergs, A. (1959). Soil sulphur fractions as chemical indices of available sulphur in some Australian soils. *Australian Journal of Agricultural Research*.  
<https://doi.org/10.1071/AR9590340>
- Xavier, G. R., Jesus, E. C., Dias, A., Coelho, M. R. R., Molina, Y. C., & Rumjanek, N. G. (2023). Contribution of biofertilizers to pulse crops: From single-strain inoculants to new technologies based on microbiome strategies. *Plants*, 12(4), 954.  
<https://doi.org/10.3390/plants12040954>

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:  
The peer review history for this paper can be accessed here:  
<https://pr.sdiarticle5.com/review-history/149897>