



Microbial Volatile Compounds and Their Role in Growth, Development and Resistance in Plants

Ashutosh Gautam ^{a*}, Sachin Kumar Vaid ^b
and Rishi Richa ^c

^a Spices Board, Regional Office, Srinagar 190 008 (Jammu & Kashmir UT), India.

^b National Center for Organic and Natural Farming, Ghaziabad 201 002 (UP), India.

^c Sher-e-Kashmir University of Agriculture Sciences & Technology, Shalimar, Srinagar 190 025 (J&K), India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ijpss/2026/v38i15958>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

<https://pr.sdiarticle5.com/review-history/115878>

Review Article

Received: 15/11/2025
Published: 30/01/2026

Abstract

Microbial Volatile Compounds (MVCs) are low molecular weight and low boiling point, high vapour pressure biochemical compounds, produced by a wide array of microorganisms ranging from bacteria to fungi. MVC are forms an interface for below and above soil interaction between microorganisms and plants. Global agricultural systems are under increasing pressure to deliver sufficient, healthy food for a growing population. MVCs can be of great benefit to plants and their use in agriculture thanks to their ability to inhibit the growth and development of plant pathogens, induce the activation of plant defense, or promote plant growth and development. Owing to their natural origin, MVCs have potential as possible alternatives to synthetic pesticides, fungicides, and bactericides as well as genetic modification. Recent studies have yielded mixed results regarding

*Corresponding author: E-mail: ashutoshgtm9@gmail.com;

Cite as: Gautam, Ashutosh, Sachin Kumar Vaid, and Rishi Richa. 2026. "Microbial Volatile Compounds and Their Role in Growth, Development and Resistance in Plants". *International Journal of Plant & Soil Science* 38 (1):441-53. <https://doi.org/10.9734/ijpss/2026/v38i15958>.

the application of MVCs. Hence, there is an urgent need to explore MVCs in more detail, including dose standardization, compatibility, method of application, and cost-effectiveness to integrate them into mainstream agriculture.

Keywords: MVCs; plants; defence; growth; modulation; agriculture.

1. Background

“Organisms differ from each other in various aspects, say genetically, morphologically, anatomically, structural complexity at cellular and sub-cellular level, habitat and many more. Chemical taxonomy, exemplified by biochemical compounds produced at various levels by organisms, plays a significant role in taxonomic classification. These biochemical compounds plays major role in adaptation, defence mechanisms, evolutionary processes etc; specifically in microorganisms like fungi, bacteria, protists. Microbial Volatile Compounds (MVCs) are of low molecular weight and boiling points” (Veselova et al., 2019), simple or complex compounds like sulphides, alcohols, benzoids, terpenes, alkenes, acids, esters, ketones and similar one (Gautam et al., 2021, Schmidt et al., 2015 and Morath et al., 2012). “Though lately, it was discovered that these MVCs can be very useful for the human being in one or other way like mediators of airborne interactions (Ryu et al., 2020), foods and beverages” (Dahunsi et al.,

2020), plant vaccines (Rodriguez et al., 2018) and many more. “MVCs derived from fungi and bacteria are responsible for the aroma of particular food stuffs like wine, cheese as well as odour from degrading food materials” (Veselova et al., 2019; Li et al., 2011; Rajer et al., 2017).

“Off late, as the interest towards MVCs grown and studies were made, it also revealed that these compounds play an important role in the plant’s growth and development in one or more ways in different stages of life cycle. Most noteworthy examples include modulation of plant growth, development of systemic resistance against pathogens, insects and pests, nematodes and a range of other organisms; where MVCs plays role of repellent or attractants” (Veselova et al., 2019).

Here in this review, attempt is made to in-depth study of the role of Microbial Volatile Compounds in the development of plants as well as resistance against pathogens/pests and insects.




Fig. 1. Microbial volatile organic compounds at a glance

2. Plants and Role of MVCs

“Microbial volatile compounds are highly active owing to their chemical structure and character. They play an active role in ecological functions, which are highly competitive in nature like increased growth in plants, development of antibiosis process, reinvigoration of defense response as well as key

morphological and structural variation for the purpose of defense” (Poveda, 2021, Gautam et al. 2016). Though there have been understandings about MVCs and their roles in different domain of ecology, further investigation are needed to delve deeper and gain more insight.”




Fig. 2. MVCs and their potential roles in agriculture

Studies in last decade have shown that MVCs have wide range of involvement in crop biology, like a precursor elicitors of plant immunity, improving the plant health through antifungal and antibacterial actions, nematicidal actions etc. (Bitas et al., 2013 and Schalchli et al., 2016). Apart of this Sharifi and Ryu (2018) and Tyagi et al. (2018) also reported MVCs role in increase of yield and biomass by virtue of modulation of hormonal and physiological pathways, change in the bulk volumes of leaves like in form of leaf number, leaf size and similarly in root.

Kanchiswamy et al. (2015), Chung et al. (2016) have shown “a different aspect of MVCs, where they reported for their noteworthy potential to act as bio-stimulants and bio-protectants; and their capability to modulate metabolic activities of an organisms, as well as proteome and genome with equal efficiency”.

2.1 MVCs in Plant-Microbe Interactions and Plant Development

“There has been a wide range of MVCs identified in bacteria and studied for their role in rhizosphere and soil ecology” (Kai et al., 2009). “Numerous bacteria, particularly rhizobacteria, have a strong affinity for soil and close

association with the rhizosphere, where they play an important role in plant growth through the exploiting the rich nutrient exudates” (Bhattacharyya and Jha, 2012). “*Pseudomonas* is a very well known and extensively explored rhizobacteria which is widely used in agriculture in one or other way for crop production and protection and have been reported for its roles in the protections of crop from pathogens and maintenance of soil health” (Hol et al., 2013). “Volatile Organic Compounds (VOCs) released by rhizobacteria have been well known for their involvement in plant-pathogen interaction, plant-microbiome interaction and antimicrobial properties” (Vespermann et al., 2007; Goswami et al., 2013). While working on *Arabidosis thaliana*, Gutierrez et al. (2010) showed that “extensive communication occurs between plants and microorganisms during different stages of plant development in which signaling molecules from the two partners play an important role. They found that certain rhizospheric bacterial strains like L263, L266, L272a, L254, L265a and L265b; can modulate the plant growth and development as well as the architecture of root system like stimulation of primary root growth and lateral root development, promotion of lateral root formation etc. through the differential emission of volatile organic compounds”.




Fig. 3. Mode of action of microbial volatile organic compounds

Microbes release a wide range of volatile organic compounds with different mode and degree of action. Blom et al. (2011) screened “forty two strains of soil borne bacterial strains to study the effects of released VOCs on the growth of *Arabidosis thaliana* and sorted out thirty six compounds which had high degree of penetration to stimulate plant growth such as 1-hexanol, indole and pentadecane”. Among the screened compounds, Dimethyl disulfide (DMDS) and ammonia were found to be among the most bioactive (Kai et al., 2010); and “DMDS is known to protect the tobacco and maize plants against the infection of *Botrytis cinerea* and *Cochliobolus heterostrophus* respectively” (Huang et al., 2012). “DMDS is also reported to have the property to alter the gene expression, more specifically to reduce the level of expression e.g. methionine biosynthesis and recycling and sulfur-assimilation genes in tobacco (*Nicotiana attenuata*)” (Meldau et al., 2013). Gautam et al. (2016) have shown “a wide range of volatile oils and oleoresins in small cardamom, that play an active role in plant development and quality of capsule”.

Ryu et al. (2003) and Rudrappa (2010) identified “3-hydroxy-2-butanone also known as acetoin and 2, 3- butanediol (2, 3-BD); which are released from were released consistently from strains of *Bacillus subtilis* and *B. Amyloliquefaciens* and responsible for inducing the systemic resistance and enhancing the total leaf surface area in *A. thaliana*”. “2, 3-BD, produced by *Enterobacter aerogenes* is also

reported to induce the resistance against Northern corn leaf blight in corn” (D’Alessandro et al., 2014). “An important VOC 3-Hexanone, produced by *Burkholderia ambifaria* is known to significantly increase the biomass of *Arabidosis thaliana*; which is very similar in the action of two other VOCs namely acetophenone and DMDS” (Groenhagen et al., 2013).

2.2 MVCs and their Role in Inducing Phenotypic Plant Responses

“Owing to their wide range of activities and properties like enhancing biomass, modulation of gene action, inducing plant defence, alteration in root architecture, modulation of plant-microbe interaction and many more, Microbial volatile organic compounds have been subject of constant interest for researcher; and considerable progress has been made in recent toward these compounds. MVCs play critical roles in multitrophic interactions during different stages of plant development in form of communication between soil microorganisms and plants as well as plant-pathogen interactions. MVCs through their multifaceted inter and intra-specific interactions; are responsible for many genetic, phenotypic and morphologic alteration of the interacting organisms” (Penuelas et al., 2014 and Effmert et al., 2012).

“Recent studies revealed that a class of MVCs, acyl-homoserine lactone (AHL) are important intercellular signaling molecules used by many bacteria to monitor their population density in

quorum-sensing control of gene expression. AHLs are known to play important roles in signalling of plant-microbe interactions, such as biofilm formation, bacterial motility, expression of pathogenicity genes, plasmid transfer, production of antibiotics etc. can be recognized by plants. AHLs can activate the bacterial cells to regulate gene expression depending on population density and they can also alter gene expression in roots and shoots and modulate the defense and cell growth responses. In *Arabidopsis*, AHLs have showed a dose dependent effect on the root architecture, initiation of lateral roots and development of root hairs, and modify the growth of primary roots” (Ortiz Castro et al., 2009 and von Rad et al., 2008). Nehvi et al. (2022, 2024) have showed that volatile oils in saffron tends to change with the impact of varying climate

“Studies in recent past unravelled the puzzles of induced systemic resistance (ISR) by the bacteria, and it was showed that majority of bacteria that activate ISR are known to do so by virtue of a salicylic acid (SA) independent pathway which involves involving jasmonate (JA) and ethylene signals” (Kanchiswamy et al., 2015). Ryu et al. (2004) showed that “VOCs released from *Bacillus subtilis* strain GB03 activates systemic resistance through an ethylene dependent pathway, which is independent of the SA or JA signalling pathways; on the other hand MVCs of *Bacillus amyloliquefaciens* strain IN937a triggers the ISR

through an ethylene independent signalling pathway”.

These studies provided new information about the role of MVOCs as initiators of defence responses in plants, and revealed that when compared with water control or treatment with the *Escherichia coli* strain DH5 α ; inoculation with bacterial strains GB03 and IN937a contribute in a very significant manner to promote the growth of *Arabidopsis* and the compounds involved were identified as acetoin and 2, 3- butanediol (2, 3-BD). It is noteworthy that when applied exogenously, acetoin and 2, 3- butanediol result in the dose dependent stimulation of plant growth.

“During the activation of ISR, different defensive cascades are elicited in response to different pathogens, as evident from a study; where the application of 2, 3-BD resulted the ISR response against *Pseudomonas carotovora* sub sp. *Carotovora* but failed to do same in case of *Pseudomonas syringae* pv. *tabaci*. Interestingly, acetosin, the precursor of 2, 3-BD successfully triggered the ISR in *Arabidopsis* against the *Pseudomonas syringae*. MVCs like 3-pentanol and 2-butanone when applied on cucumber seedlings consistently, can trigger the plant systemic defense responses against *Pseudomonas syringae* pv. *lachrymans*; through the activation of gene expression of plant green leaf volatile signalling pathway”.




Fig. 4. Effects caused by MVCs in plants, fungi and bacteria

Song and Ryu (2013) showed that “these compounds led to an increase in crop yield but have not had any effect on plant growth”.

An interesting outcome from the studies by Ezquer et al. (2010) revealed that “microbial species known for MVOC emission strongly promoted starch accumulation in leaves of both mono- and dicotyledonous plants. The leaves of two days treatment of plants with these MVCs, showed a starch content comparable to potato tuber and in some cases even in higher side. To some extents, this explains the course of action of MVCs towards an enhanced yield of a crop”.

While investigating the starch content in the leaves of MVCs treated plants, Li et al. (2011) found that “MVOC induced starch accumulation process (MVOCISAP) is subjected to photoreceptor-mediated control; as evident from the starch level in the illuminated leaves of MVOCs treated *hy1/cry1*, *hy1/cry2*, and *hy1/cry1/cry2* mutants of *Arabidopsis*. Molecular studies of enzymatic and transcriptome level activities of potato leaves treated with the MVCs released by *Alternaria alternata*; showed that the phenomenon of starch accumulation due to exposure to MVCs is accompanied by multiple factors, viz. up-regulation of sucrose synthase invertase inhibitors, starch synthase (SS) class III and IV, starch branching enzyme and glucose-6-phosphate transporter”.

3. Harnessing the MVCs Crop Protection and Production and Challenges

In the rapidly changing climatic conditions and emergence of new races of insect-pests and pathogens, there is a constant threat to food production around the globe. Owing to the drastic effects on human and environment health, there has been a constant debate over the use of synthetic pesticides and fertilizers over the last two decades.

“This has paved the way for formulation and applications of bio-pesticides, bio-fertilizers, and bio-control agents derived from living microbes are becoming suitable in a larger scale as a replacement of synthetic inputs. However, the high costs, inconsistent results and field performance and reduced efficiency have been a serious concern for these replacement inputs” (Glare et al., 2012).

“In such scenario, MVCs offer an alternative for a cost effective, result oriented and sustainable crop production and protection system for food

security. Since past two decades, extensive research on MVCs and their roles in soil-plant, plant-microbes, plant-plant have provided new insights and conceptual understanding of the extremely complex and dynamic nature of these compounds and their potential role for a sustainable crop production and protection system. As discussed earlier, a wide range of MVCs have been reported and tested for their action against different pathogens, their ability to modulate the developmental process of a plant e.g. root architecture and total leaf area, and enhancing the production of biomass and economic yield e.g. enhanced starch accumulation etc. through different metabolic, physiological and transcriptional changes. It has been very well established that plants have the ability to perceive and respond to MVCs; this means MVCs have a very good prospects of application in crop production and protection. It won't be an exaggeration if we say that researchers are tapping a new era of MVCs and their potential application for a better tomorrow”.

“Application of MVCs is still in its tender stage with a limited practical and commercial availability. As of now merely 400 bacteria and fungi and 1000 MVCs released by them are identified and documented released by have been described in the literature”. “Though MVCs have shown potential for a wide range of application in agriculture system, but most of the results are from lab studies rather than the data from open field conditions. However, there have been few studies in open field condition in recent past and showed the promising results” (Cortes Barco et al., 2010a). These results back up the assumptions and idea of the application of MVCs in open field conditions for protection against herbivores, to increase pathogen resistance, and bio-control agents in general.

“There have been considerable differences in the results in lab and field conditions; also there have been ample reports which showed that MVCs may modulate growth/defense in a species-dependent manner and may be specific to conditions. One such example is 2, 3-butanediol, which in an open field condition has shown to exert its effect only as a modulator of defense but no effect as growth modulator. This makes it a pre-requisite to demonstrate and evaluate a single MVC or mixtures of MVCs on different crop species both at lab and field conditions, before generalization of the MVCs as growth or defence modulators” (Cortes Barco et al., 2010a).

Table 1. Some microbial volatile organic compounds, their source and effects

Microbe group	MVCs	Source	Effects	Reference			
Bacteria	Benzaldehyde	<i>Bacillus amyloliquefaciens</i> strain FZB42	Resistance against <i>Ralstonia solanacearum</i>	Tahir et al., (2017a)			
	Dimethylhexadecylamine	<i>Arthrobacter agilis</i>	Growth promotion in <i>Medicago sativa</i>	Velazquez Becerra et al., (2011)			
	2,5-Dimethyl pyrazine	<i>Bacillus amyloliquefaciens</i>	Antifungal activity against <i>Fusarium</i> sp. and <i>Colletotrichum gloeosporioides</i>	Guevara Avendaño et al., (2019)			
	Nonanal	<i>Bacillus subtilis</i> FA26	Resistance against <i>Xanthomonas oryzae</i>	Rajer et al., (2017)			
	Benzothiazole						
	Acetophenone						
	Decyl alcohol						
	3,5,5-Trimethylhexanol						
	Chloroacetic acid	<i>Bacillus</i> strain D13	Antifungal activity against <i>Colletotrichum gloeosporioides</i>	Xie et al., (2018)			
	Octadecane	<i>Bacillus atrophaeus</i> HAB-5	Antibacterial activity against <i>Ralstonia solanacearum</i>	Rajaofera et al., (2019)			
	Hexadecanoic acid						
	2-Undecanone	<i>Bacillus amyloliquefaciens</i> SQR-9	Antifungal activity against <i>Colletotrichum acutatum</i>	Raza et al., (2016d)			
	2-Tridecanone						
	Heptadecane						
	Undecanal						
	2-Ethyl-1-hexanol						
	2-Undecanone						
	1-undecene						
	Hydrogen cyanide				<i>Lysinibacillus</i> sp. <i>P. brassicacearum</i>	Antifungal activity against <i>Rosellinia necatrix</i> , <i>F. oxysporum</i> , <i>F. solani</i> , <i>R. solani</i> , <i>S. Sclerotium</i> , <i>Macrophomina phaseolina</i> , <i>B. cinerea</i> , <i>Fusarium equiseti</i> , and <i>Verticillium dahliae</i>	Ossowicki et al., (2017)
	Dimethyl sulfide						
S-Methyl thioacetate	<i>Pseudomonas donghuensis</i>				Oomycetocidal activity against <i>Phytophthora cactorum</i>	Macías Rubalcava et al., (2018)	
Methyl thiocyanate							
Dimethyl trisulfide							
1-undecane							
Fungi	2-Methyl-1-butanol	<i>Hypoxyton anthochroum</i>	Antifungal activity against <i>Fusarium oxysporum</i>	Macías Rubalcava et al., (2018)			
	3-Methyl-1-butanol						

Microbes emits a plethora of MVCs, and there are very chances that a particular microbial compound or a mixture of such compounds has a synergistic effect on plant resistance to diseases; and at same time it may have a positive effect on plant growth and development. This can offer an abundance of blends of MVCs and their effective application to modulate the growth and development and disease resistance of crop plants.

“Another concern with the application of MVCs in crop production is manner of application of and their exploitation in open field conditions and their side effects. Most of the MVCs are known for their rapid evaporation rates, therefore the protocols for their application are needed to standardize at greater scale. Similarly, many MVCs are known to have inhibitory effects and some point of concentration they may be toxic also. Therefore an extensive assessment of the dose-response effect on specific crops, characterization of bioactive molecules are required before their use can be safely managed” (Cortes Barco et al., 2010a).

4. Conclusion

Microbial volatile organic compounds form an interface of microbial interaction at above and below ground level. These are very complex and dynamic in nature and a better understanding about their mode of action at molecular level, ecology and evolution can pave the way for effective use towards the crop protection and production for sustainable agriculture perspective. MVCs offer a most desirable alternative of synthetic pesticides or fertilizers; as they are biodegradable and can be applied in low concentration, they are in synergy with nature and no serious hazardous effects as in the case of synthetic pesticides or fertilizers.

To understand the roles, efficacy and suitability of MVCs; more open field studies coupled with physiological and molecular studies, are required to use the microbial volatile organic compounds in a cost effective, eco-friendly and sustainable manner to realise their potential in a sustainable crop production and protection.

Disclaimer (Artificial Intelligence)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

Competing Interests

Authors have declared that no competing interests exist.

References

- Bhattacharyya, P., & Jha, D. (2012). Plant growth promoting rhizobacteria (PGPR): Emergence in agriculture. *World Journal of Microbiology and Biotechnology*, 28, 1327–1350. <https://doi.org/10.1007/s11274-011-0979-9>
- Bitas, V., Kim, H. S., Bennett, J. W., & Kang, S. (2013). Sniffing on microbes: Diverse roles of microbial volatile organic compounds in plant health. *Molecular Plant–Microbe Interactions*, 26, 835–843. <https://doi.org/10.1094/MPMI-10-12-0249-CR>
- Blom, D., Fabbri, C., Connor, E., Schiestl, F., Klausner, D., & Boller, T. (2011). Production of plant growth modulating volatiles is widespread among rhizosphere bacteria and strongly depends on culture conditions. *Environmental Microbiology*, 13, 3047–3058. <https://doi.org/10.1111/j.1462-2920.2011.02582.x>
- Che, J., Liu, B., Liu, G., Chen, Q., & Lan, J. (2017). Volatile organic compounds produced by *Lysinibacillus* sp. FJAT-4748 possess antifungal activity against *Colletotrichum acutatum*. *Biocontrol Science and Technology*, 27, 1349–1362. <https://doi.org/10.1080/09583157.2017.1397600>
- Chung, J. H., Song, G. C., & Ryu, C. M. (2016). Sweet scents from good bacteria: Case studies on bacterial volatile compounds for plant growth and immunity. *Plant Molecular Biology*, 90, 677–687. <https://doi.org/10.1007/s11103-015-0344-8>
- Cortés Barco, A., Goodwin, P., & Hsiang, T. (2010a). Comparison of induced resistance activated by benzothiadiazole, (2R,3R)-butanediol and an isoparaffin mixture against anthracnose of *Nicotiana benthamiana*. *Plant Pathology*, 59, 643–653. <https://doi.org/10.1111/j.1365-3059.2010.02283.x>
- D’Alessandro, M., Erb, M., Ton, J., Brandenburg, A., Karlen, D., & Zopfi, J. (2014). Volatiles produced by soil-borne endophytic bacteria increase plant pathogen resistance and affect tritrophic interactions. *Plant, Cell &*

- Environment*, 37, 813–826.
<https://doi.org/10.1111/pce.12220>
- Dahunsi, A. T. A., Dahunsi, S. O., & Olayanju, A. (2020). Synergistic microbial interaction between lactic acid bacteria and yeasts during production of Nigerian indigenous fermented foods and beverages. *Food Control*, 110, 106963.
- Di Giorgio, A., De Stradis, A., Cantore, P., & Iacobellis, N. S. (2015). Biocide effects of volatile organic compounds produced by potential biocontrol rhizobacteria on *Sclerotinia sclerotiorum*. *Frontiers in Microbiology*, 6, 1056.
<https://doi.org/10.3389/fmicb.2015.01056>
- Effmert, U., Kalderas, J., Warnke, R., & Piechulla, B. (2012). Volatile mediated interactions between bacteria and fungi in the soil. *Journal of Chemical Ecology*, 38, 665–703. <https://doi.org/10.1007/s10886-012-0135-5>
- Ezquer, I., Li, J., Ovecka, M., Baroja-Fernández, E., Muñoz, F. J., & Montero, M. (2010). Microbial volatile emissions promote accumulation of exceptionally high levels of starch in leaves in mono- and dicotyledonous plants. *Plant Cell Physiology*, 51, 1674–1693.
<https://doi.org/10.1093/pcp/pcq126>
- Gautam, A., Bharat Kumar, C., Pradip Kumar, K., & Rao, Y. S. (2016). Profiling of oils and oleo-resins in small cardamom (*Elettaria cardamomum*) as a prerequisite parameter of quality. *Progressive Research – An International Journal*, 11(Special II), 1415–1418.
- Gautam, A., Leela, N. K., Chhetri, R., Bora, S. S., Deka, T. N., Pradip Kumar, K., Krishnamoorthy, K. S., & Sharon, A. (2021). Study on composition of essential oils of large cardamom (*Amomum subulatum* Roxb.) and its wild relatives found in Himalayan region, Sikkim. *Proceedings of the 24th Plantation Crop Symposium*, 82–83.
- Glare, T., Caradus, J., Gelernter, W., Jackson, T., Keyhani, N., & Kohl, J. (2012). Have bio-pesticides come of age? *Trends in Biotechnology*, 30, 250–258.
<https://doi.org/10.1016/j.tibtech.2012.01.003>
- Goswami, D., Vaghela, H., Parmar, S., Dhandhukia, P., & Thakker, J. N. (2013). Plant growth promoting potentials of *Pseudomonas* spp. strain OG isolated from marine water. *Journal of Plant Interactions*, 8, 281–290.
<https://doi.org/10.1080/17429145.2013.768360>
- Groenhagen, U., Baumgartner, R., Bailly, A., Gardiner, A., Eberl, L., & Schulz, S. (2013). Production of bioactive volatiles by different *Burkholderia ambifaria* strains. *Journal of Chemical Ecology*, 39, 892–906.
<https://doi.org/10.1007/s10886-013-0315-y>
- Guevara-Avenidaño, E., Bejarano-Bolívar, A. A., Kiel-Martínez, A. L., Ramírez-Vázquez, M., Méndez-Bravo, A., & von Wobeser, E. A. (2019). Avocado rhizobacteria emit volatile organic compounds with antifungal activity against *Fusarium solani*, *Fusarium* sp. associated with Kuroshio shot hole borer, and *Colletotrichum gloeosporioides*. *Microbiological Research*, 219, 74–83.
<https://doi.org/10.1016/j.micres.2018.11.009>
- Gutiérrez-Luna, F. M., López-Bucio, J., Altamirano-Hernández, J., Valencia-Cantero, E., de la Cruz, H. R., & Macías-Rodríguez, L. (2010). Plant growth promoting rhizobacteria modulate root system architecture in *Arabidopsis thaliana* through volatile organic compound emission. *Symbiosis*, 51, 75–83.
<https://doi.org/10.1007/s13199-010-0066-2>
- Hassan, S. R. U., Strobel, G. A., Geary, B., & Sears, J. (2013). An endophytic *Nodulisporium* sp. from Central America producing volatile organic compounds with both biological and fuel potential. *Journal of Microbiology and Biotechnology*, 23, 29–35. <https://doi.org/10.4014/jmb.1208.04062>
- Hol, W. H. G., Bezemer, T. M., & Biere, A. (2013). Getting the ecology into interactions between plants and the plant growth-promoting bacterium *Pseudomonas fluorescens*. *Frontiers in Plant Science*, 4, 81.
<https://doi.org/10.3389/fpls.2013.00081>
- Huang, C. J., Tsay, J. F., Chang, S. Y., Yang, H. P., Wu, W. S., & Chen, C. Y. (2012). Dimethyl disulfide is an induced systemic resistance elicitor produced by *Bacillus cereus* C1L. *Pest Management Science*, 68, 1306–1310.
<https://doi.org/10.1002/ps.3301>
- Kai, M., Crespo, E., Cristescu, S. M., Harren, F. J., Francke, W., & Piechulla, B. (2010). *Serratia odorifera*: Analysis of volatile emission and biological impact of volatile compounds on *Arabidopsis thaliana*. *Applied Microbiology and Biotechnology*, 88, 965–976.
<https://doi.org/10.1007/s00253-010-2810-1>

- Kai, M., Haustein, M., Molina, F., Petri, A., Scholz, B., & Piechulla, B. (2009). Bacterial volatiles and their action potential. *Applied Microbiology and Biotechnology*, 81, 1001–1012. <https://doi.org/10.1007/s00253-008-1760-3>
- Kanchiswamy, C. N., Malnoy, M., & Maffei, M. E. (2015). Bioprospecting bacterial and fungal volatiles for sustainable agriculture. *Trends in Plant Science*, 20, 206–211. <https://doi.org/10.1016/j.tplants.2015.01.004>
- Kanchiswamy, C. N., Malnoy, M., & Maffei, M. E. (2015). Chemical diversity of microbial volatiles and their potential for plant growth and productivity. *Frontiers in Plant Science*, 6, Article 151. <https://doi.org/10.3389/fpls.2015.00151>
- Li, J., Ezquer, I., Bahaji, A., Montero, M., Ovecka, M., & Baroja-Fernández, E. (2011). Microbial volatile-induced accumulation of exceptionally high levels of starch in *Arabidopsis* leaves is a process involving NTRC and starch synthase classes III and IV. *Molecular Plant–Microbe Interactions*, 24, 1165–1178. <https://doi.org/10.1094/MPMI-05-11-0112>
- Li, N., Alfiky, A., Wang, W., Islam, M., Nourollahi, K., Liu, X., & Kang, S. (2018a). Volatile compound-mediated recognition and inhibition between *Trichoderma* biocontrol agents and *Fusarium oxysporum*. *Frontiers in Microbiology*, 9, 2614. <https://doi.org/10.3389/fmicb.2018.02614>
- Macías-Rubalcava, M. L., Sánchez-Fernández, R. E., Roque-Flores, G., Lappe-Oliveras, P., & Medina-Romero, Y. M. (2018). Volatile organic compounds from *Hypoxyylon anthochroum* endophytic strains as postharvest mycofumigation alternative for cherry tomatoes. *Food Microbiology*, 76, 363–373. <https://doi.org/10.1016/j.fm.2018.06.014>
- Meldau, D. G., Meldau, S., Hoang, L. H., Underberg, S., Wuensche, H., & Baldwin, I. T. (2013). Dimethyl-disulfide produced by the naturally associated bacterium *Bacillus* sp. B55 promotes *Nicotiana attenuata* growth by enhancing sulfur nutrition. *Plant Cell*, 25, 2731–2747. <https://doi.org/10.1105/tpc.113.114744>
- Morath, S. U., Hung, R., & Bennett, J. W. (2012). Fungal volatile organic compounds: A review with emphasis on their biotechnological potential. *Fungal Biology Reviews*, 26, 73–83. <https://doi.org/10.1016/j.fbr.2012.07.001>
- Naznin, H. A., Kimura, M., Miyazawa, M., & Hyakumachi, M. (2013). Analysis of volatile organic compounds emitted by plant growth-promoting fungus *Phoma* sp. GS8-3 for growth promotion effects on tobacco. *Microbes and Environments*, 28, 42–49. <https://doi.org/10.1264/jsme2.ME12085>
- Nehvi, F. A., Dar, N. A., Elai, B. A., Khan, M. H., Iqbal, A. M., Mir, G. H., Gautam, A., & Qayoom, S. (2022). Effect of trigger weather on saffron phenology under temperate conditions of Jammu and Kashmir, India. *International Journal of Agriculture Sciences*, 14(2), 11085–11089.
- Nehvi, F. A., Gautam, A., Qayoom, S., Kanth, R. H., Elai, B. A., Dar, N. A., Nasir, S., Nagoo, S. A., Khan, M. H., Mir, G. H., Iqbal, A. M., & Sofi, M. D. (2024). Comparative yield gaps, weather impact and constraint analysis in saffron heritage site of Pampore, Kashmir, India. *Vegetos – An International Journal of Plant Research & Biotechnology*, 38(1), 236–249.
- Nieto-Jacobo, M. F., Steyaert, J. M., Salazar-Badillo, F. B., Nguyen, D. V., Rostás, M., & Braithwaite, M. (2017). Environmental growth conditions of *Trichoderma* spp. affect indole acetic acid derivatives, volatile organic compounds, and plant growth promotion. *Frontiers in Plant Science*, 8, 102. <https://doi.org/10.3389/fpls.2017.00102>
- Ortiz-Castro, R., Contreras-Cornejo, H. A., Macías-Rodríguez, L., & López-Bucio, J. (2009). The role of microbial signals in plant growth and development. *Plant Signaling & Behavior*, 4, 701–712. <https://doi.org/10.4161/psb.4.8.9047>
- Ossowicki, A., Jafra, S., & Garbeva, P. (2017). The antimicrobial volatile power of the rhizospheric isolate *Pseudomonas donghuensis* P482. *PLOS ONE*, 12, e0174362. <https://doi.org/10.1371/journal.pone.0174362>
- Peñuelas, J., Asensio, D., Tholl, D., Wenke, K., Rosenkranz, M., & Piechulla, B. (2014). Biogenic volatile emissions from the soil. *Plant, Cell & Environment*, 37, 1866–1891. <https://doi.org/10.1111/pce.12340>
- Poveda, J. (2021). Beneficial effects of microbial volatile organic compounds (MVOCs) in plants. *Applied Soil Ecology*, 168, 104118.
- Rajaofera, M. J. N., Wang, Y., Dahar, G. Y., Jin, P., Fan, L., Xu, L., et al. (2019). Volatile

- organic compounds of *Bacillus atrophaeus* HAB-5 inhibit the growth of *Colletotrichum gloeosporioides*. *Pesticide Biochemistry and Physiology*, 156, 170–176. <https://doi.org/10.1016/j.pestbp.2019.02.019>
- Rajer, F. U., Wu, H., Xie, Y., Xie, S., Raza, W., Tahir, H. A. S., & Gao, X. (2017). Volatile organic compounds produced by a soil isolate, *Bacillus subtilis* FA26, induce adverse ultrastructural changes to the cells of *Clavibacter michiganensis* subsp. *sepedonicus*. *Microbiology*, 163, 523–530.
- Raza, W., Ling, N., Yang, L., Huang, Q., & Shen, Q. (2016). Response of tomato wilt pathogen *Ralstonia solanacearum* to the volatile organic compounds produced by a biocontrol strain *Bacillus amyloliquefaciens* SQR-9. *Scientific Reports*, 6, 24856. <https://doi.org/10.1038/srep24856>
- Rezende, D. C., Maurício, B. F., Simone, C. B., Silvia, B., & Sergio, F. P. (2015). Antimicrobial activity of volatile organic compounds and their effect on lipid peroxidation and electrolyte loss in *Colletotrichum gloeosporioides* and *Colletotrichum acutatum* mycelia. *African Journal of Microbiology Research*, 9, 1527–1535.
- Rodríguez, E. Q., Durán, D., Heil, M., & Coronel, X. C. (2018). Damage-associated molecular patterns (DAMPs) as future plant vaccines that protect crops from pests. *Scientia Horticulturae*, 237, 207–220. <https://doi.org/10.1016/j.scienta.2018.03.026>
- Rudrappa, T., Biedrzycki, M. L., Kunjeti, S. G., Donofrio, N. M., Czymbek, K. J., & Paré, P. W. (2010). The rhizobacterial elicitor acetoin induces systemic resistance in *Arabidopsis thaliana*. *Communicative & Integrative Biology*, 3, 130–138. <https://doi.org/10.4161/cib.3.2.10584>
- Ryu, C. M., Farag, M. A., Hu, C. H., Reddy, M. S., Kloepper, J. W., & Paré, P. W. (2004). Bacterial volatiles induce systemic resistance in *Arabidopsis*. *Plant Physiology*, 134, 1017–1026. <https://doi.org/10.1104/pp.103.026583>
- Ryu, C. M., Farag, M. A., Hu, C. H., Reddy, M. S., Wei, H. X., & Paré, P. W. (2003). Bacterial volatiles promote growth in *Arabidopsis*. *Proceedings of the National Academy of Sciences of the United States of America*, 100, 4927–4932. <https://doi.org/10.1073/pnas.0730845100>
- Ryu, C. M., Weisskopf, L., & Piechulla, B. (2020). *Bacterial volatile compounds as mediators of airborne interactions*. Springer Nature Singapore. <https://doi.org/10.1007/978-981-15-7293-7>
- Sánchez-Ortiz, B. L., Sánchez-Fernández, R. E., Duarte, G., Lappe-Oliveras, P., & Macías-Rubalcava, M. L. (2016). Antifungal, antioomycete and phytotoxic effects of volatile organic compounds from the endophytic fungus *Xylaria* sp. strain PB3f3 isolated from *Haematoxylon brasiletto*. *Journal of Applied Microbiology*, 120, 1313–1325. <https://doi.org/10.1111/jam.13101>
- Schalchli, H., Tortella, G. R., Rubilar, O., Parra, L., Hormazabal, E., & Quiroz, A. (2016). Fungal volatiles: An environmentally friendly tool to control pathogenic microorganisms in plants. *Critical Reviews in Biotechnology*, 36, 144–152. <https://doi.org/10.3109/07388551.2014.946466>
- Schmidt, R., Cordovez, V., de Boer, W., Raaijmakers, J., & Garbeva, P. (2015). Volatile affairs in microbial interactions. *The ISME Journal*, 9, 2329–2335. <https://doi.org/10.1038/ismej.2015.42>
- Sharifi, R., & Ryu, C. M. (2018). Revisiting bacterial volatile-mediated plant growth promotion: Lessons from the past and objectives for the future. *Annals of Botany*, 122, 349–358. <https://doi.org/10.1093/aob/mcy108>
- Singh, S. K., Strobel, G. A., Knighton, B., Geary, B., Sears, J., & Ezra, D. (2011). An endophytic *Phomopsis* sp. possessing bioactivity and fuel potential with its volatile organic compounds. *Microbial Ecology*, 61, 729–739. <https://doi.org/10.1007/s00248-011-9818-7>
- Song, G. C., & Ryu, C. M. (2013). Two volatile organic compounds trigger plant self-defense against a bacterial pathogen and a sucking insect in cucumber under open field conditions. *International Journal of Molecular Sciences*, 14, 9803–9819. <https://doi.org/10.3390/ijms14059803>
- Stinson, M., Ezra, D., Hess, W. M., Sears, J., & Strobel, G. (2003). An endophytic *Gliocladium* sp. of *Eucryphia cordifolia* producing selective volatile antimicrobial compounds. *Plant Science*, 165, 913–922. [https://doi.org/10.1016/S0168-9452\(03\)00299-1](https://doi.org/10.1016/S0168-9452(03)00299-1)
- Tahir, H. A. S., Gu, Q., Wu, H., Niu, Y., Huo, R., & Gao, X. (2017). *Bacillus* volatiles

- adversely affect the physiology and ultrastructure of *Ralstonia solanacearum* and induce systemic resistance in tobacco against bacterial wilt. *Scientific Reports*, 7, 40481.
- Tyagi, S., Mulla, S. I., Lee, K. J., Chae, J. C., & Shukla, P. (2018). VOCs-mediated hormonal signaling and crosstalk with plant growth-promoting microbes. *Critical Reviews in Biotechnology*, 38, 1277–1296. <https://doi.org/10.1080/07388551.2018.1472551>
- Velázquez-Becerra, C., Macías-Rodríguez, L. I., López-Bucio, J., Altamirano-Hernández, J., Flores-Cortez, I., & Valencia-Cantero, E. (2011). A volatile organic compound analysis from *Arthrobacter agilis* identifies dimethylhexadecylamine, an amino-containing lipid modulating bacterial growth and *Medicago sativa* morphogenesis in vitro. *Plant and Soil*, 339, 329–340. <https://doi.org/10.1007/s11104-010-0583-z>
- Veselova, M. A., Plyuta, V. A., & Khmel, I. A. (2019). Volatile compounds of bacterial origin: Structure, biosynthesis, and biological activity. *Microbiology*, 88, 261–274. <https://doi.org/10.1134/S0026261719030160>
- Vespermann, A., Kai, M., & Piechulla, B. (2007). Rhizobacterial volatiles affect the growth of fungi and *Arabidopsis thaliana*. *Applied and Environmental Microbiology*, 73, 5639–5641. <https://doi.org/10.1128/AEM.01078-07>
- von Rad, U., Klein, I., Dobrev, P. I., Kottová, J., Zazimalová, E., & Fekete, A. (2008). Response of *Arabidopsis thaliana* to N-hexanoyl-DL-homoserine lactone, a bacterial quorum sensing molecule produced in the rhizosphere. *Planta*, 229, 73–85. <https://doi.org/10.1007/s00425-008-0811-4>
- Xie, S., Zang, H., Wu, H. J., Rajer, F. U., & Gao, X. (2018). Antibacterial effects of volatiles produced by *Bacillus* strain D13 against *Xanthomonas oryzae* pv. *oryzae*. *Molecular Plant Pathology*, 19, 49–58.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2026): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://pr.sdiarticle5.com/review-history/115878>