



# Evaluation of Tomato (*Solanum lycopersicum* L.) Genotypes for Growth and Yield Attributing Traits

Puli Vijaya Babu <sup>a\*</sup>, D. Laxminarayana <sup>b</sup>, P. Prasanth <sup>b</sup>,  
S. Mallesh <sup>a</sup>, B. Sai Krishna Nikhil <sup>c</sup> and P. Pranusha <sup>d</sup>

<sup>a</sup> Department of Vegetable Science, Sri Konda Laxman Telangana Horticultural University, Mulugu, Telangana, India.

<sup>b</sup> Department of Horticulture, Sri Konda Laxman Telangana Horticultural University, Mulugu, Telangana, India.

<sup>c</sup> Department of Genetics and Plant breeding, Sri Konda Laxman Telangana Horticultural University, Mulugu, Telangana, India.

<sup>d</sup> Department of Crop Physiology, National Bureau of Plant Genetic Resources, Regional Station, Hyderabad, India.

## *Authors' contributions*

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## *Article Information*

DOI: <https://doi.org/10.9734/ijpss/2026/v38i46045>

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://pr.sdiarticle5.com/review-history/156520>

**Original Research Article**

**Received: 17/02/2026**

**Accepted: 10/04/2026**

**Published: 13/04/2026**

## **Abstract**

Tomato (*Solanum lycopersicum* L.) is an economically important vegetable crop, and improving its yield potential through the evaluation of diverse genotypes remains a key objective in breeding programmes. The identification of superior genotypes based on growth and yield attributes is essential for enhancing productivity under varying agro-climatic conditions. The study was conducted to evaluate the tomato

\*Corresponding author: E-mail: [pulivijaybabu@gmail.com](mailto:pulivijaybabu@gmail.com);

genotypes for growth and yield attributing traits during *summer* 2024. The present investigation was conducted at PG Students Research Farm, College of Horticulture, Sri Konda Laxman Telangana Horticultural University, Rajendranagar, Hyderabad, India. The experimental material consisted of twenty-four tomato genotypes and followed randomized block design with two replications. Growth and yield attributing parameters were recorded such as plant spread, stem girth, leaf area, days to first flowering, number of flowers per cluster, days to last fruit harvest, number of fruits per cluster and number of fruit clusters per plant. Among the genotypes, maximum leaf area was recorded in the genotype EC 620516 (113.18 cm<sup>2</sup>). Minimum days for first flowering were recorded in EC 620385 (25.50). Among the genotypes, maximum number of fruits per cluster were observed in EC 631373 (4.80). Maximum number of fruit clusters per plant were recorded in EC 620516 (8.60). The superior performed tomato genotypes for different growth and yield attributing traits can be used in further crop improvement programme.

**Keywords:** Evaluation; tomato genotypes; growth; yield; traits.

## 1. Introduction

Tomato (*Solanum lycopersicum* L.), a member of the Solanaceae family with a chromosome number of  $2n = 24$  (Peralta *et al.*, 2005), is believed to have originated in the Peru–Ecuador–Bolivia region (Rick, 1969). It is one of the most important solanaceous vegetable crops cultivated globally and plays a pivotal role in both subsistence and commercial agriculture. Owing to its versatility, tomato serves as a fundamental ingredient in a wide range of culinary preparations, contributing significantly to human nutrition through its rich content of vitamins, minerals, and antioxidants.

India is the second-largest producer of vegetables after China, with a total production of 219,674.29 thousand metric tonnes from an area of 11,750.76 thousand hectares. Within this, tomato occupies a prominent position, being cultivated over 821.98 thousand hectares with a production of 20,752.36 thousand metric tonnes. Major tomato-producing states in India include Andhra Pradesh, Karnataka, Madhya Pradesh, Telangana, Gujarat, Tamil Nadu, Odisha, and West Bengal. In Telangana alone, tomato is grown over an area of 12.06 thousand hectares, yielding 352.51 thousand metric tonnes (Ministry of Agriculture and Farmers Welfare, Government of India, 2024).

Despite its widespread cultivation, there remains considerable scope for improving yield potential and quality attributes through systematic evaluation and selection of superior genotypes. Therefore, the present study was undertaken to assess diverse tomato cultivars and breeding lines from varied agro-ecological and genetic backgrounds for key yield-attributing traits. The ultimate objective is to identify or develop high-performing genotypes that combine enhanced productivity with desirable quality characteristics, thereby contributing to sustainable tomato production.

## 2. Material and Methods

### 2.1 Experiment Details

The present investigation was conducted at PG Students Research Farm, College of Horticulture, Sri Konda Laxman Telangana Horticultural University, Rajendranagar, Hyderabad during *Summer*, 2024. Randomized block design was followed with two replications.

### 2.2 Experimental Material

The experimental material comprising twenty-four tomato genotypes (Table 1) among them twenty-two exotic collections from the National Bureau of Plant Genetic Resources, Regional Station, Hyderabad, Arka Vikas from Indian Institute of Horticultural Research, Bengaluru and PKM-1 from Tamil Nadu Agricultural University, Coimbatore were collected for conducting the experiment.

### 2.3 Seedlings Raising, Transplanting and Cultural Operations

Seeds of all tomato genotypes were sown in 98-celled plastic pro-trays filled with cocopeat, germination was observed at 5–7 days after sowing. Seedlings were maintained under shade net conditions with regular watering as required and proper care was provided until transplanting.

**Table 1. Details of tomato genotypes**

| Sl. No. | Genotypes  | Source                                               |
|---------|------------|------------------------------------------------------|
| 1       | EC 615039  | NBPGR, Regional station, Hyderabad-30                |
| 2       | EC 620366  | NBPGR, Regional station, Hyderabad-30                |
| 3       | EC 620385  | NBPGR, Regional station, Hyderabad-30                |
| 4       | EC 620510  | NBPGR, Regional station, Hyderabad-30                |
| 5       | EC 620516  | NBPGR, Regional station, Hyderabad-30                |
| 6       | EC 620543  | NBPGR, Regional station, Hyderabad-30                |
| 7       | EC 631399  | NBPGR, Regional station, Hyderabad-30                |
| 8       | EC 631957  | NBPGR, Regional station, Hyderabad-30                |
| 9       | EC 638522  | NBPGR, Regional station, Hyderabad-30                |
| 10      | EC 620414  | NBPGR, Regional station, Hyderabad-30                |
| 11      | EC 687423  | NBPGR, Regional station, Hyderabad-30                |
| 12      | EC 631477  | NBPGR, Regional station, Hyderabad-30                |
| 13      | EC 631455  | NBPGR, Regional station, Hyderabad-30                |
| 14      | EC 620388  | NBPGR, Regional station, Hyderabad-30                |
| 15      | EC 615040  | NBPGR, Regional station, Hyderabad-30                |
| 16      | EC 638518  | NBPGR, Regional station, Hyderabad-30                |
| 17      | EC 631436  | NBPGR, Regional station, Hyderabad-30                |
| 18      | EC 690982  | NBPGR, Regional station, Hyderabad-30                |
| 19      | EC 690989  | NBPGR, Regional station, Hyderabad-30                |
| 20      | EC 631373  | NBPGR, Regional station, Hyderabad-30                |
| 21      | EC 617055  | NBPGR, Regional station, Hyderabad-30                |
| 22      | EC 620360  | NBPGR, Regional station, Hyderabad-30                |
| 23      | PKM -1     | Tamil Nadu Agricultural University, Coimbatore       |
| 24      | Arka Vikas | Indian Institute of Horticulture Research, Bengaluru |

*EC: Exotic collection, NBPGR: National Bureau of Plant Genetic Resources*

The main field was ploughed three times and brought to a fine tilth using a tractor-drawn cultivator. After leveling, raised beds were prepared according to the experimental design, drip lines were installed and farmyard manure along with NPK fertilizers (120:60:60 kg ha<sup>-1</sup>) were applied as recommended by Sri Konda Laxman Telangana Horticultural University's Udyana Panchangam.

Mulching was applied and 30-day-old tomato seedlings were transplanted into the plots with 60 cm row-to-row and 45 cm plant-to-plant spacing. Wooden stakes provided support as needed, irrigation followed immediately post-transplanting for uniform establishment and subsequent watering occurred based on requirements, with hand weeding every 25–30 days.

Prophylactic plant protection involved sprays of insecticides (Imidacloprid 17.8% SL and Thiamethoxam 25% WG) and fungicides (Carbendazim 12% + Mancozeb 63% WP; Metalaxyl 4% + Mancozeb 64% WP).

## 2.4 Observations Recorded

### 2.4.1 Growth Traits

Plant spread was measured from tagged plants using a measuring scale in the east-west and north-south directions and the average value was expressed in centimeters (cm). The plant's stem girth was measured using a digital vernier caliper and the millimeter value was converted and expressed in centimeters. The leaf area was recorded using a leaf area meter (LI-3100 Area Meter) as well as expressed in square centimeters.

### 2.4.2 Yield Attributing Traits

The number of days taken from the date of transplanting to the day when the first flower appeared in each treatment was counted and noted at the flowering stage. The number of flowers in each selected cluster of tagged plants was counted and their mean was calculated. Days taken from the date of transplanting to the date of last fruit picking at the marketable stage were counted and recorded. Before the harvesting, the fruits in each

selected cluster of the tagged plants were counted and recorded. The total number of fruit bearing clusters were counted up to the end of the crop cycle and recorded from five tagged plants of each replication.

## 2.5 Statistical Analysis

### 2.5.1 Analysis of Variance (ANOVA)

Combined analysis of variance (ANOVA) was carried out for each trait, with environments and genotypes included as factors. Mean sums of squares were determined for genotypes, environments and their interaction.

### 2.5.2 Mean

The mean value of each character was worked out by dividing the totals by the corresponding number of observations.

$$\text{Mean (X)} = \frac{\sum X_{ij}}{N}$$

Where,

$X_{ij}$  = Any observation in the  $i^{\text{th}}$  genotype and  $j^{\text{th}}$  replication  
 $N$  = Total number of observations.

### 2.5.3 Standard Error (S.E.)

The standard errors (S.E. (m)±) for genotypes were calculated with the help of the mean square due to error from the analysis of variance table by the following formula:

$$\text{SE (m)} \pm = \frac{\sqrt{MSE}}{r}$$

Where,

MSE = Mean of squares due to error  
 $r$  = Number of replications

### 2.5.4 Critical Difference

Critical difference was calculated to find out the superiority of one variety over the other by following the formula.

$$CD = S.E (d) \times t \text{ value at } 5 \% \text{ and } 1 \% \text{ error degrees of freedom.}$$

Where,

$$S.E (d) = \frac{\sqrt{2 \times MSE}}{r}$$

$t$  = Table value of 't' distribution at error degrees of freedom on  $P < 0.05$  and  $0.01$ .

### 2.5.5 Coefficient of Variation (C.V.)

$$CV \% = \frac{\sqrt{MSE}}{X} \times 100$$

Where,

$X$  = Grand mean

### 3. Results and Discussion

#### 3.1 Growth Traits

##### 3.1.1 Plant Spread (cm)

The analysis for plant spread in the genotypes ranged (Table 2) from 34.60 (cm) to 54.40 (cm) with overall mean (Fig. 1) of 45.67 (cm). Among the genotypes, minimum plant spread was recorded in EC 690982 (34.60 cm) followed by EC 690989 (36.35 cm), while maximum plant spread was recorded in EC 631399 (54.40 cm). The results for the trait plant spread are in confirmation with the findings of Raut *et al.* (2021).

**Table 2. Mean performance of twenty-four tomato genotypes for growth traits**

| Sl. No. | Genotypes      | Plant spread (cm) | Stem girth (cm) | Leaf area (cm <sup>2</sup> ) |
|---------|----------------|-------------------|-----------------|------------------------------|
| 1       | EC 615039      | 37.90             | 0.99            | 61.57                        |
| 2       | EC 620366      | 46.90             | 1.51            | 51.03                        |
| 3       | EC 620385      | 40.60             | 1.64            | 70.88                        |
| 4       | EC 620510      | 51.00             | 1.25            | 88.64                        |
| 5       | EC 620516      | 53.80             | 1.68            | 113.18                       |
| 6       | EC 620543      | 49.70             | 1.53            | 55.03                        |
| 7       | EC 631399      | 54.40             | 2.29            | 67.37                        |
| 8       | EC 631957      | 41.80             | 1.55            | 50.39                        |
| 9       | EC 638522      | 50.30             | 1.54            | 95.13                        |
| 10      | EC 620414      | 52.00             | 2.27            | 94.35                        |
| 11      | EC 687423      | 49.70             | 1.56            | 83.18                        |
| 12      | EC 631477      | 41.80             | 1.63            | 93.01                        |
| 13      | EC 631455      | 49.75             | 2.25            | 63.82                        |
| 14      | EC 620388      | 43.00             | 1.25            | 65.60                        |
| 15      | EC 615040      | 49.10             | 1.31            | 95.49                        |
| 16      | EC 638518      | 47.30             | 1.55            | 63.20                        |
| 17      | EC 631436      | 37.70             | 1.64            | 53.96                        |
| 18      | EC 690982      | 34.60             | 1.15            | 68.58                        |
| 19      | EC 690989      | 36.35             | 1.11            | 58.48                        |
| 20      | EC 631373      | 39.10             | 1.08            | 61.41                        |
| 21      | EC 617055      | 48.40             | 1.57            | 76.57                        |
| 22      | EC 620360      | 49.90             | 1.19            | 91.44                        |
| 23      | PKM -1         | 48.40             | 1.55            | 73.15                        |
| 24      | Arka Vikas     | 42.60             | 1.52            | 78.72                        |
|         | <b>Mean</b>    | 45.67             | 1.53            | 73.92                        |
|         | <b>SE.(m)</b>  | 1.82              | 0.10            | 3.40                         |
|         | <b>SE.(d)</b>  | 2.57              | 0.14            | 4.80                         |
|         | <b>CD (5%)</b> | 5.32              | 0.30            | 9.95                         |
|         | <b>CV (%)</b>  | 5.63              | 9.48            | 6.51                         |

CD: Critical difference, CV: Coefficient of variance, SE.(m): Standard error mean, SE.(d): Standard error deviation

##### 3.1.2 Stem Girth (cm)

The mean values for stem girth in the genotypes were varied (Table 2) from 0.99 (cm) to 2.29 (cm) with overall mean of 1.53 (cm). The genotype EC 631399 (2.29 cm) recorded maximum stem girth followed by EC 620414 (2.27 cm), whereas minimum stem girth was recorded in EC 615039 (0.99 cm). The results are in line with Swain (2022).

##### 3.1.3 Leaf Area

Among the genotypes, mean values for leaf area were varied (Table 2) from 50.39 (cm<sup>2</sup>) to 113.18 (cm<sup>2</sup>) with overall mean of 73.92 (cm<sup>2</sup>). Maximum leaf area was recorded (Fig. 2) in the genotype EC 620516 (113.18 cm<sup>2</sup>)

followed by EC 615040 (95.49 cm<sup>2</sup>), while minimum leaf area was found in EC 631957 (50.39 cm<sup>2</sup>). The observations are in accordance with the findings of Oladimeji (2025).




Fig. 1. Mean performance of twenty-four tomato genotypes for plant spread (cm)




Fig. 2. Mean performance of twenty-four tomato genotypes for leaf area (cm<sup>2</sup>)

### 3.2 Yield Attributing Traits


#### 3.2.1 Days taken to First Flowering

Number of days taken to first flowering among the genotypes were varied from 25.50 to 30.50 days with overall mean of 28.21 days (Table 3). The genotypes EC 631957, EC 638522 and EC 690989 recorded maximum (30.50) number of days for first flowering followed by EC 620510 and EC 620360 (29.50), whereas, minimum days for first flowering were recorded in EC 620385 (25.50). The results are comparable with findings of Maurya *et al.* (2022).

#### 3.2.2 Number of Flowers per Cluster

Among the genotypes number of flowers per cluster were ranged (Fig. 3) from 3.30 cm to 6.90 with overall mean of 5.02 (Table 3). Among the genotypes, EC 631373 (6.90) recorded highest number of flowers per cluster followed by EC690982 (6.50), while lowest (3.30) was observed in genotype, Arka Vikas. The observations are

in accordance with the findings of Kumar *et al.* (2016), Maurya *et al.* (2022), Rasheed *et al.* (2023) and Oladimeji (2025).



**Fig. 3. Mean performance of twenty-four tomato genotypes for number of flowers per cluster**

**Table 3. Mean performance of twenty-four tomato genotypes for yield attributing traits**

| SI. No | Genotypes      | Days taken to first flowering | Number of flowers per cluster | Days taken for last fruit harvest | Number of fruits per cluster | Number of fruit clusters per plant |
|--------|----------------|-------------------------------|-------------------------------|-----------------------------------|------------------------------|------------------------------------|
| 1      | EC 615039      | 28.50                         | 4.90                          | 97.50                             | 2.40                         | 5.50                               |
| 2      | EC 620366      | 27.50                         | 4.70                          | 92.50                             | 2.60                         | 6.40                               |
| 3      | EC 620385      | 25.50                         | 5.30                          | 84.50                             | 2.60                         | 6.50                               |
| 4      | EC 620510      | 29.50                         | 5.50                          | 109.50                            | 2.50                         | 7.70                               |
| 5      | EC 620516      | 28.50                         | 5.70                          | 107.50                            | 4.20                         | 8.60                               |
| 6      | EC 620543      | 28.50                         | 5.50                          | 103.50                            | 3.10                         | 8.30                               |
| 7      | EC 631399      | 28.50                         | 5.30                          | 98.00                             | 3.30                         | 7.70                               |
| 8      | EC 631957      | 30.50                         | 5.50                          | 101.00                            | 3.50                         | 4.70                               |
| 9      | EC 638522      | 30.50                         | 4.50                          | 103.50                            | 2.80                         | 6.30                               |
| 10     | EC 620414      | 28.50                         | 5.50                          | 104.00                            | 3.50                         | 8.30                               |
| 11     | EC 687423      | 28.50                         | 5.50                          | 98.00                             | 2.70                         | 5.20                               |
| 12     | EC 631477      | 28.50                         | 4.50                          | 100.50                            | 2.90                         | 5.90                               |
| 13     | EC 631455      | 26.50                         | 5.00                          | 89.50                             | 2.50                         | 5.90                               |
| 14     | EC 620388      | 27.50                         | 5.50                          | 89.00                             | 3.30                         | 8.50                               |
| 15     | EC 615040      | 27.50                         | 4.50                          | 90.00                             | 3.00                         | 6.50                               |
| 16     | EC 638518      | 26.50                         | 5.50                          | 95.00                             | 3.60                         | 4.30                               |
| 17     | EC 631436      | 29.50                         | 3.60                          | 99.00                             | 2.30                         | 7.70                               |
| 18     | EC 690982      | 27.50                         | 6.50                          | 96.50                             | 3.50                         | 6.00                               |
| 19     | EC 690989      | 30.50                         | 5.30                          | 96.00                             | 3.50                         | 5.70                               |
| 20     | EC 631373      | 28.50                         | 6.90                          | 95.50                             | 4.80                         | 5.90                               |
| 21     | EC 617055      | 27.50                         | 4.70                          | 96.50                             | 3.40                         | 6.10                               |
| 22     | EC 620360      | 29.50                         | 3.90                          | 98.50                             | 2.60                         | 5.30                               |
| 23     | PKM -1         | 26.50                         | 3.40                          | 89.50                             | 2.50                         | 8.30                               |
| 24     | Arka Vikas     | 26.50                         | 3.30                          | 94.50                             | 2.30                         | 8.50                               |
|        | <b>Mean</b>    | 28.21                         | 5.02                          | 97.06                             | 3.06                         | 6.66                               |
|        | <b>SE.(m)</b>  | 0.49                          | 0.12                          | 0.81                              | 0.11                         | 0.29                               |
|        | <b>SE.(d)</b>  | 0.69                          | 0.16                          | 1.13                              | 0.16                         | 0.41                               |
|        | <b>CD (5%)</b> | 1.45                          | 0.35                          | 2.36                              | 0.33                         | 0.86                               |
|        | <b>CV (%)</b>  | 2.48                          | 3.37                          | 1.17                              | 5.27                         | 6.26                               |

CD: Critical difference, CV: Coefficient of variance, SE.(m): Standard error mean, SE.(d): Standard error deviation

### 3.2.3 Days Taken for Last Fruit Harvest

Number of days taken for last fruit harvest among the genotypes varied from 84.50 to 109.50 with overall mean of 97.06 (Table 3). The genotype EC 620510 (109.50 cm) recorded maximum days for last fruit harvest followed by EC 620516 (107.50 cm), whereas minimum days for last fruit harvest were recorded in EC 620385 (84.50). The results are in line with the findings of Reddy *et al.* (2014), Meitei *et al.* (2014), Kumar *et al.* (2016) and Shankar (2016).

### 3.2.4 Number of Fruits per Cluster

The analysis for number of fruits per cluster in the genotypes ranged from 2.30 to 4.80 with overall mean of 3.06 (Table 3). Among the genotypes, minimum number of fruits per cluster were recorded in EC 631436 (2.30) followed by EC 615039 (2.40), while maximum number of fruits per cluster were observed in EC 631373 (4.80). The results are in accordance with the findings of Ullah *et al.* (2015), Kumar *et al.* (2016), Rasheed *et al.* (2023) and Oladimeji (2025).

### 3.2.5 Number of Fruit Clusters per Plant

Among the genotypes number of fruit clusters per plant were ranged (Fig. 4) from 4.30 to 8.60 with overall mean of 6.66 (Table 3). Among the genotypes, EC 620516 (8.60) recorded highest number of fruit clusters per plant followed by EC 620388 and Arka Vikas (8.50), while lowest number of fruit clusters (4.30) were observed in EC 638518. The results are in accordance to the findings of Mehta and Asati (2008), Meena and Bahadur (2015), Jatav *et al.* (2016) and Rasheed *et al.* (2023).




Fig. 4. Mean performance of twenty-four tomato genotypes for number of fruit clusters per plant

## 4. Conclusion

The study revealed significant variations in growth and yield attributing traits of tomato genotypes, including plant spread, stem girth, leaf area, days to first flowering, number of flowers per cluster, days to last fruit harvest, number of fruits per cluster and number of fruit clusters per plant. Genotypes such as EC 620516, EC 620388, Arka Vikas, EC 631399, EC 620385, EC 631373, EC690982 and EC 620510 stood out for their superior performance in terms of growth and yield attributing traits, respectively. Therefore, these genotypes can be effectively employed in future breeding initiatives to enhance fruit yield attributes.

## Disclaimer (Artificial Intelligence)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

## Competing Interests

Authors have declared that no competing interests exist.

## References

- Jatav, P. K., Dhangal, V. P. S., & Sachin, S. C. (2016). Studies on genetic variability, heritability and genetic advance for growth, yield and quality parameters in elite genotypes of tomato (*Lycopersicum esculentum* Mill.). *Advances in Life Sciences*, 5(20), 9201–9204.
- Kumar, A. P., Reddy, R. K., Reddy, R. V. S. K., Pandravada, S. R., & Saidaiah, P. (2016). Heritability studies in dual purpose tomato genotypes for growth, yield and quality attributes. *Plant Archives*, 16(2), 885–889.  
[https://www.researchgate.net/publication/319000000\\_Heritability\\_studies\\_in\\_dual\\_purpose\\_tomato\\_genotypes\\_for\\_growth\\_yield\\_and\\_quality\\_attributes](https://www.researchgate.net/publication/319000000_Heritability_studies_in_dual_purpose_tomato_genotypes_for_growth_yield_and_quality_attributes)
- Maurya, D., Akhtar, S., Chattopadhyay, T., Kumar, R., Sahay, S., Sangam, S., et al. (2022). Genetic variability and character association in tomato (*Solanum lycopersicum* L.). *Bangladesh Journal of Botany*, 51(4), 747–757. <https://doi.org/10.3329/bjb.v51i4.63494>
- Meena, O. P., & Bahadur, V. (2015). Genetic associations analysis for fruit yield and its contributing traits of indeterminate tomato (*Solanum lycopersicum* L.) germplasm under open field condition. *Journal of Agricultural Science*. <https://doi.org/10.5539/jas.v7n3p148>
- Mehta, N., & Asati, B. S. (2008). Genetic relationship of growth and development traits with fruit yield in tomato (*Solanum lycopersicum* L.). *Karnataka Journal of Agricultural Science*, 21(1), 92–96.
- Meitei, K. M., Bora, G. C., Singh, S. J., & Sinha, A. K. (2014). Morphology based genetic variability analysis and identification of important characters for tomato (*Solanum lycopersicum* L.) crop improvement. *American-Eurasian Journal of Agricultural & Environmental Science*, 14(10), 1105–1111. <https://doi.org/10.5829/idosi.aejas.2014.14.10.12434>
- Ministry of Agriculture and Farmers Welfare, Government of India. (2024). *Second advance estimates of production of foodgrains, oilseeds and other commercial crops for the year 2023–24*. Department of Agriculture & Farmers Welfare. <https://upag.gov.in/imagedefault/estimates>
- Oladimeji, A. A. (2025). Estimation of variance components, heritability and genetic advance of yield and yield related traits in tomatoes (*Solanum lycopersicum* L.). *Journal of Agricultural and Environmental Science Research*, 8(1), 44–56. <https://doi.org/10.70382/hujaesr.v8i1.028>
- Peralta, I. E., Knapp, S., & Spooner, D. M. (2005). New species of wild tomatoes (*Solanum* section *Lycopersicon*: Solanaceae) from Northern Peru. *Systematic Botany*, 30(2), 424–434. <https://doi.org/10.1600/0363644054223657>
- Rasheed, A., Ilyas, M., Khan, T. N., Mahmood, A., Riaz, U., Chattha, M. B., et al. (2023). Study of genetic variability, heritability, and genetic advance for yield-related traits in tomato (*Solanum lycopersicon* MILL.). *Frontiers in Genetics*. <https://doi.org/10.3389/fgene.2022.1030309>
- Raut, N., Patil, H. B., Basavaraj, N., Jagadeesha, R. C., Fakrudin, B., Cholin, S., et al. (2021). Assessment of genetic variability in tomato (*Solanum lycopersicum* L.) for yield and yield attributing traits. *The Pharma Innovation Journal*, 10(4), 399–403.
- Reddy, B. R., Begum, H., Reddy, M. A., Sunil, A., & Reddy, M. P. (2014). Genetic variability and effect relationship for yield & quantitative traits in exotic lines of tomato. *Green Farming*, 5(1), 104–107. <https://www.greenfarming.in/>
- Rick, C. M. (1969). Origin of cultivated tomato, current status of the problem. *Abstract XI International Botanical Congress*, 180.
- Shankar, A. (2016). *Studies on heterosis, combining ability and stability for yield and its components in tomato (Solanum lycopersicum L.)* (Ph.D. (Horti.) thesis). Dr. Y. S. R Horticultural University, Andhra Pradesh.
- Swain, S. (2022). *Genetic variability study in advance breeding lines of tomato (Lycopersicum esculentum L.)* (M.Sc. (Agri.) thesis). Odisha University of Agriculture and Technology, Bhubaneswar, India.

Ullah, M. Z., Hassan, L., Shahid, S. B., & Patwary, A. K. (2015). Variability and inter relationship studies in tomato (*Solanum lycopersicum* L.). *Journal of the Bangladesh Agricultural University*, 13(1), 65–69. <https://doi.org/10.3329/jbau.v13i1.28716>

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2026): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://pr.sdiarticle5.com/review-history/156520>