



Influence of Weed and Nitrogen Management on Growth, Productivity and Economics of Barley (*Hordeum vulgare* L.)

Lokesh Kumar ^a, Karmnath Kumar ^a, Sohit ^a, Sucheta Dahiya ^{a*}
and Tinku Raj Singh ^a

^a Department of Natural Resource Management (Agronomy), Faculty of Agricultural Sciences (FASC), SGT University, Gurugram, Haryana, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ijpss/2026/v38i66116>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://pr.sdiarticle5.com/review-history/157607>

Original Research Article

Received: 27/02/2026

Accepted: 26/04/2026

Published: 03/06/2026

Abstract

Background: Barley (*Hordeum vulgare* L.) productivity is often constrained by severe weed competition and improper nitrogen management, which adversely affect crop growth, yield, and nutrient use efficiency. Integrated weed and nitrogen management is therefore essential to enhance productivity and profitability under semi-arid conditions.

Methods: A field experiment was conducted during the *Rabi* season of 2021–22 at the Agronomy Research Farm, SGT University, Gurugram, Haryana. The experiment was laid out in a factorial randomized block design with three replications, comprising four weed management treatments (W₁: weedy check, W₂: weed-free, W₃: metsulfuron, W₄: 2,4-D) Observations on growth, yield attributes, yield, nitrogen uptake, and economics were recorded and analyzed statistically.

Results: Weed management and nitrogen levels significantly influenced growth, yield attributes, and productivity of barley. Treatment W₂ and W₃ improved plant growth, dry matter accumulation, and yield

*Corresponding author: E-mail: candidsucheta@gmail.com;

attributes over W_1 . Among nitrogen levels, N_3 recorded superior performance, followed by N_2 . Grain yield, biological yield, and nitrogen uptake increased with effective weed control and higher nitrogen levels. Among interaction $W_2 \times N_2$ showed maximum productivity and nutrient uptake. W_3 resulted in higher profitability, while N_3 recorded maximum returns, though N_2 was more cost-effective.

Keywords: Barley; weed management; nitrogen levels; yield; nutrient uptake.

1. Introduction

Barley (*Hordeum vulgare* L.) is one of the oldest and most important cereal crops cultivated worldwide, ranking fourth after wheat, rice, and maize. It serves as a major source of food, feed, and raw material for malting and brewing industries. Owing to its wide adaptability to diverse agro-climatic conditions, barley plays a crucial role in ensuring food security, particularly in semi-arid and temperate regions (Kaur *et al.*, 2024; Cammarano *et al.*, 2024).

Barley productivity is often constrained by weed infestation and improper nitrogen management. Weeds compete for nutrients, moisture, light, and space during early growth, reducing crop vigor and yield by 20–50% depending on conditions (Mendera *et al.*, 2024; Dorado *et al.*, 2025; Naeem *et al.*, 2022; Safdar *et al.*, 2022). Timely weed control is essential.

Effective weed management is essential for improving barley productivity. Herbicide application is widely adopted due to its efficiency and cost-effectiveness. Proper weed control enhances growth, yield attributes, and economic returns in barley cultivation Kumar & Jha, (2017).

Nitrogen is essential for plant growth, aiding chlorophyll formation, photosynthesis, and protein synthesis. Adequate supply enhances growth and yield, while inefficient management reduces efficiency and causes environmental issues, requiring optimization (Sisay *et al.*, 2025; Scapino *et al.*, 2025). Efficient nitrogen management not only increases crop yield but also improves economic returns and minimizes environmental risks by reducing input losses (Loukakis *et al.*, 2025).

The interaction between weed management and nitrogen fertilization plays a crucial role in crop performance. While nitrogen enhances both crop and weed growth, effective weed control improves nutrient use efficiency. Thus, integrating suitable weed management with optimum nitrogen levels is essential for maximizing productivity and economic returns in barley cultivation. (Bogale *et al.*, 2024; Al-Jayashi *et al.*, 2024; Singh *et al.*, 2025; Hirich *et al.*, 2026; Shpanev and Smuk, 2025). Therefore, the present study was undertaken to evaluate the influence of weed and nitrogen management on growth, productivity, nutrient uptake, and economics of barley.

2. Materials and Methods

The present field investigation was conducted during the *Rabi* season of 2021–22 at the Agronomy Research Farm, Faculty of Agricultural Sciences, SGT University, Gurugram, (Haryana), India. The experimental site is situated in the semi-arid region of the north-western Indo-Gangetic plains, characterized by typical climatic conditions of the area.

The soil of the experimental field was sandy loam in texture, slightly alkaline in reaction, and had low to medium fertility status based on initial soil analysis. Soil samples were collected from 0–15 cm depth before sowing and analyzed using standard procedures, including alkaline $KMnO_4$ method for nitrogen (Subbaiah and Asija, 1956), Olsen's method for phosphorus (Olsen *et al.*, 1954), and flame photometric method for potassium determination (Jackson, 1973).

The experiment was laid out in a Factorial Randomized Block Design (FRBD) with three replications. The treatments consisted of two factors: weed management and nitrogen levels. Weed management included four treatments, *viz.*, weedy check (W_1), weed-free (W_2), metsulfuron at 4 g ha⁻¹ applied at 30 DAS, (W_3), and 2,4-D at 0.5 kg ha⁻¹ applied at 30 DAS (W_4). Nitrogen was applied at four levels: 0 kg ha⁻¹ (N_0), 30 kg ha⁻¹ (N_1), 60 kg ha⁻¹ (N_2), and 90 kg ha⁻¹ (N_3). Thus, a total of 16 treatment combinations were evaluated.

Each plot measured 3.0 m × 2.7 m, and the total number of experimental plots was 48. Barley variety BH 393 was sown on 22 November 2021-22 using a seed rate of 100 kg ha⁻¹, maintaining a row spacing of 22.5 cm. The field was prepared by ploughing followed by harrowing and planking to obtain a fine tilth. A pre-sowing irrigation was applied to ensure uniform germination.

Fertilizers were applied as per treatments and phosphorus at 40 kg ha⁻¹ as basal. Irrigation was given at CRI and subsequent stages. Herbicides were applied at 30 DAS as per treatment in the treated plots, while weed-free plots were manually throughout the crop season. Growth parameters such as plant height and dry matter accumulation at different growth stages were observed. Yield attributes including effective tillers, spike length, grains per spike, and test weight were also recorded. Yield parameters such as grain yield, straw yield, biological yield, and harvest index (Grain yield/Biological yield *100) were computed following standard agronomic procedures. In addition, nitrogen content and uptake in grain and straw, as well as economic parameters including cost of cultivation, gross return, net return, and benefit–cost ratio, were calculated using established methods (Jackson, 1973).

The experimental data were statistically analyzed using analysis of variance (ANOVA) appropriate for Factorial Randomized Block Design, and treatment means were compared at the 5% level of significance using the critical difference (CD) test, as described by (Gomez and Gomez, 1984).

3. Results and Discussion

The present investigation clearly demonstrated that both weed management practices and nitrogen levels significantly influenced growth, yield attributes, productivity, nutrient uptake, and economic returns of barley.

3.1 Crop Establishment and Plant Growth

The initial plant population was not significantly influenced by weed management or nitrogen levels (Table 1), indicating uniform crop establishment across treatments. Plant height increased progressively with crop growth (Table 2). Among weed management treatments, W₂ recorded the highest plant height at harvest, followed by W₃, which remained statistically at par, while W₁ showed comparatively lower growth.” This improvement may be attributed to reduced competition for light, nutrients, and moisture, as also reported by (Al-Gburi *et al.*, 2024; Kumar & Singh 2018; Naeem *et al.*, 2022; Cammarano *et al.*, 2024; Sisay *et al.*, 2025).

Nitrogen levels significantly influenced plant height, with N₃ producing the tallest plants, followed by N₂, N₁, and N₀ (Table 2). Dry matter accumulation followed a similar pattern (Table 3), with maximum biomass under W₂ and N₃. Interaction effects (Tables 4 and 5) further confirmed enhanced growth under combined treatments.

3.2 Yield Attributes

Yield attributes were significantly influenced by both weed and nitrogen management practices (Table 6). Among weed treatments, W₂ recorded superior performance in terms of yield attributes, which was statistically at par with W₃, while W₁ showed comparatively lower values. This indicates that effective chemical weed control can serve as a viable alternative to manual weeding.

Among nitrogen levels, N₃ recorded the highest values of yield attributes, followed by N₂, N₁, and N₀ (Table 6). This improvement may be attributed to enhanced nutrient availability and increased photosynthetic efficiency (Loukakis *et al.*, 2025; Saikia *et al.*, 2018; Scapino *et al.*, 2025).

The interaction effect on spike length (Table 7) further revealed that combinations of effective weed management (W₂ and W₃) with higher nitrogen levels (N₂ and N₃) resulted in improved spike development, emphasizing the importance of integrated weed and nutrient management practices.

3.3 Yield Performance

Grain yield, straw yield, and biological yield were significantly influenced by weed and nitrogen management practices (Table 8). Among weed treatments, W₂ recorded the highest grain yield, which was statistically at par

with W₃, while W₁ resulted in the lowest yield due to severe weed competition. This clearly indicates the detrimental effect of uncontrolled weeds on crop productivity.

Nitrogen application significantly enhanced yield, with N₃ producing the highest yield, followed by N₂, N₁, and N₀ (Table 8). However, the response beyond N₂ was comparatively less pronounced, suggesting diminishing returns at higher nitrogen levels. Similar trends have been reported by (Sisay *et al.*, 2025; Pooniya *et al.*, 2021; Kumar *et al.*, 2022; Kantwa *et al.*, 2025; Bogale *et al.*, 2024).

Interaction effects (Tables 9 and 10) revealed that combinations of effective weed management (W₂ and W₃) with higher nitrogen levels (N₂ and N₃) significantly improved straw and biological yield, indicating better utilization of available resources.

3.4 Harvest Index

Harvest index was significantly influenced by weed and nitrogen management practices (Table 8). Among weed control treatments, W₂ recorded the highest harvest index, followed by W₃ and W₄, while W₁ showed lower values, indicating better assimilate partitioning under effective weed control. Similar findings were reported by (Arshad *et al.*, 2025; Puniya *et al.*, 2023). Interaction effects (Table 11) showed that W₂ combined with higher nitrogen levels (N₂ and N₃) improved harvest index. This may be due to synergistic effects of nitrogen and weed control on photosynthesis and dry matter distribution, as also reported by (Dolijanović *et al.*, 2025; Shekhawat *et al.*, 2021).

3.5 Nitrogen Content and Uptake

Nitrogen content and uptake were significantly influenced by weed and nitrogen management practices (Table 12). Among weed treatments, W₂ recorded higher nitrogen uptake, followed by W₃, while W₁ showed lower uptake due to nutrient competition. This may be attributed to reduced crop–weed competition and improved nutrient availability, as reported by (Kumar *et al.*, 2022; Naeem *et al.*, 2022).

Among nitrogen levels, N₃ recorded the highest uptake, followed by N₂, N₁, and N₀, indicating better nutrient utilization under higher nitrogen application. Similar trends were observed by (Loukakis *et al.*, 2025; Kantwa *et al.*, 2025; Scapino *et al.*, 2025; Saikia *et al.*, 2018).

3.6 Economic Analysis

Economic analysis (Table 13) revealed that among weed management treatments, W₃ recorded the highest net return and benefit–cost ratio, outperforming W₂, W₄, and W₁, mainly due to reduced cost of weed control and efficient resource utilization.

Among nitrogen levels, N₃ recorded the highest net return and B:C ratio, followed by N₂, N₁, and N₀. However, considering input costs and marginal returns, N₂ may be a more economically viable option under limited resource conditions.

These findings are in agreement with (Singh *et al.*, 2025; Sunil *et al.*, 2025; Bogale *et al.*, 2024), who also reported higher profitability under integrated weed and nutrient management practices.

Table 1. Effect of different treatments on initial population of barley

Treatments	Initial population (no. m ⁻¹) at 30 DAS
Weed management	
W ₁ : Weedy check (no weed control)	40.16
W ₂ : Weed-free	45.83
W ₃ : Application of Metsulfuron-methyl at 4 g ha ⁻¹ at 30 DAS	41.67
W ₄ : Application of 2,4-D at 0.5 kg ha ⁻¹ at 30 DAS	40.00
SE(m) ±	2.46
C.D. at 5 %	NS

Treatments	Initial population (no. m ⁻¹) at 30 DAS
Nitrogen management	
No: Control (0 kg N ha ⁻¹)	39.83
N ₁ : 30 kg N ha ⁻¹	41.25
N ₂ : 60 kg N ha ⁻¹	42.00
N ₃ : 90 kg N ha ⁻¹	44.58
SEm±	2.46
CD (P = 5%)	NS

Table 2. Effect of different treatments on plant height of barley at different growth stages

Treatment	Plant height (cm)			
	30 DAS	60 DAS	90 DAS	At harvest
Weed management				
W ₁ : Weedy check (no weed control)	5.87	45.90	64.99	69.26
W ₂ : Weed-free	8.33	51.68	69.10	73.57
W ₃ : Application of Metsulfuron-methyl at 4 g ha ⁻¹ at 30 DAS	7.20	50.35	68.57	72.77
W ₄ : Application of 2,4-D at 0.5 kg ha ⁻¹ at 30 DAS	6.83	47.97	66.81	72.31
SE(m)±	0.80	2.12	1.00	1.07
C.D. at 5 %	NS	NS	2.91	3.11
Nitrogen management				
No: Control (0 kg N ha ⁻¹)	6.95	40.75	58.01	62.31
N ₁ : 30 kg N ha ⁻¹	6.79	46.26	65.80	69.11
N ₂ : 60 kg N ha ⁻¹	7.04	51.99	72.46	78.63
N ₃ : 90 kg N ha ⁻¹	7.45	56.90	73.20	77.87
SEm±	0.80	2.12	1.00	1.07
CD (P = 5%)	NS	6.16	2.91	3.11

Table 3. Effect of weed and nitrogen management on plant dry weight of barley at different growth stages

Treatment	Plant dry weight (g plant ⁻¹)			
	30 DAS	60 DAS	90 DAS	At Harvest
Weed management				
W ₁ : Weedy check (no weed control)	5.87	36.35	120.98	165.56
W ₂ : Weed-free	8.33	42.68	151.76	209.68
W ₃ : Application of Metsulfuron-methyl at 4 g ha ⁻¹ at 30 DAS	7.20	38.79	136.12	193.68
W ₄ : Application of 2,4-D at 0.5 kg ha ⁻¹ at 30 DAS	6.83	38.55	131.55	179.20
SE(m)±	0.80	3.72	2.40	2.69
C.D. at 5 %	NS	NS	6.95	7.81
Nitrogen management				
No: Control (0 kg N ha ⁻¹)	6.79	33.12	113.79	144.64
N ₁ : 30 kg N ha ⁻¹	6.95	36.81	122.30	174.07
N ₂ : 60 kg N ha ⁻¹	7.04	38.73	139.13	204.28
N ₃ : 90 kg N ha ⁻¹	7.45	47.70	165.19	225.14
SEm±	0.80	3.72	2.40	2.69
CD (P = 5%)	NS	NS	6.95	7.81

Table 4. Interaction of weed and nitrogen on dry weight of barley at 90 DAS

Treatment	N ₀ : Control (0 kg N ha ⁻¹)	N ₁ : 30 kg N ha ⁻¹	N ₂ : 60 kg N ha ⁻¹	N ₃ : 90 kg N ha ⁻¹	Mean
W ₁ : Weedy check (no weed control)	116.8	118.7	122.0	126.4	121.0
W ₂ : Weed-free	124.1	127.4	161.4	194.1	151.8
W ₃ : Application of Metsulfuron-methyl at 4 g ha ⁻¹ at 30 DAS	100.5	115.5	141.7	186.9	136.1
W ₄ : Application of 2,4-D at 0.5 kg ha ⁻¹ at 30 DAS	113.7	127.6	131.4	153.5	131.6
Mean	113.8	122.3	139.1	165.2	
SEm±	4.79				
CD (P = 5%)	13.9				

Table 5. Interaction of weed and nitrogen on dry weight of barley at harvest

Treatment	N ₀ : Control (0 kg N ha ⁻¹)	N ₁ : 30 kg N ha ⁻¹	N ₂ : 60 kg N ha ⁻¹	N ₃ : 90 kg N ha ⁻¹	Mean
W ₁ : Weedy check (no weed control)	130.0	163.4	176.8	192.0	165.6
W ₂ : Weed-free	161.3	179.6	230.8	267.1	209.7
W ₃ : Application of Metsulfuron-methyl at 4 g ha ⁻¹ at 30 DAS	164.6	180.3	208.8	221.1	193.7
W ₄ : Application of 2,4-D at 0.5 kg ha ⁻¹ at 30 DAS	122.7	173.0	200.7	220.4	179.2
Mean	144.6	174.1	204.3	225.1	
SEm±	5.38				
CD (P = 5%)	15.6				

Table 6. Effect of different treatments on yield attributes of barley

Treatment	Effective tillers (no. m ⁻¹)	Spike length (cm)	Grains spike ⁻¹	Test weight (g)
Weed management				
W ₁ : Weedy check (no weed control)	75.08	5.52	39.90	36.50
W ₂ : Weed-free	88.50	5.98	44.45	37.16
W ₃ : Application of Metsulfuron-methyl at 4 g ha ⁻¹ at 30 DAS	85.25	5.88	43.15	36.87
W ₄ : Application of 2,4-D at 0.5 kg ha ⁻¹ at 30 DAS	84.08	5.80	41.69	36.62
SE(m)±	2.54	0.08	1.34	0.54
C.D. at 5%	7.38	0.25	NS	NS
Nitrogen management				
N ₀ : Control (0 kg N ha ⁻¹)	71.33	5.02	35.83	35.87
N ₁ : 30 kg N ha ⁻¹	80.50	5.87	42.77	36.45
N ₂ : 60 kg N ha ⁻¹	84.58	5.99	45.30	37.08
N ₃ : 90 kg N ha ⁻¹	96.50	6.30	45.30	37.75
SEm±	2.54	0.08	1.34	0.54
CD (P = 5%)	7.38	0.25	3.90	NS

Table 7. Interaction of weed and nitrogen on spike length of barley

Treatment	N ₀ : Control (0 kg N ha ⁻¹)	N ₁ : 30 kg N ha ⁻¹	N ₂ : 60 kg N ha ⁻¹	N ₃ : 90 kg N ha ⁻¹	Mean
W ₁ : Weedy check (no weed control)	4.27	5.83	6.00	5.99	5.52
W ₂ : Weed-free	5.33	5.83	5.88	6.89	5.98
W ₃ : Application of Metsulfuron-methyl at 4 g ha ⁻¹ at 30 DAS	5.11	6.38	5.78	6.27	5.89
W ₄ : Application of 2,4-D at 0.5 kg ha ⁻¹ at 30 DAS	5.38	5.44	6.33	6.05	5.80
Mean	5.02	5.87	6.00	6.30	
SEm±	0.17				
CD (P = 5%)	0.50				

Table 8. Effect of different treatments on yield and harvest index of barley

Treatment	Grain yield (kg ha ⁻¹)	Straw yield (kg ha ⁻¹)	Biological yield (kg ha ⁻¹)	Harvest index (%)
Weed management				
W ₁ : Weedy check (no weed control)	1984	2574	4,972	36.09
W ₂ : Weed-free	2538	3,261	5,530	47.43
W ₃ : Application of Metsulfuron-methyl at 4 g ha ⁻¹ at 30 DAS	2494	3,019	5,468	44.77
W ₄ : Application of 2,4-D at 0.5 kg ha ⁻¹ at 30 DAS	2261	3,004	5,369	44.91
SE(m)±	38	52	82	0.97
C.D. at 5 %	111	151	238	2.81
Nitrogen management				
No: Control (0 kg N ha ⁻¹)	1294	1,751	3,145	41.06
N ₁ : 30 kg N ha ⁻¹	2068	2,752	4,850	43.12
N ₂ : 60 kg N ha ⁻¹	2817	3,510	6,510	43.61
N ₃ : 90 kg N ha ⁻¹	3098	3,846	6,834	45.40
SEm±	38	52	82	0.97
CD (P = 5%)	111	151	238	2.81

Table 9. Interaction of weed and nitrogen on straw yield of barley

Treatment	N ₀ : Control (0 kg N ha ⁻¹)	N ₁ : 30 kg N ha ⁻¹	N ₂ : 60 kg N ha ⁻¹	N ₃ : 90 kg N ha ⁻¹	Mean
W ₁ : Weedy check (no weed control)	1,353	2,696	2,830	3,418	2,574
W ₂ : Weed-free	1,852	3,002	3,634	4,558	3,261
W ₃ : Application of Metsulfuron-methyl at 4 g ha ⁻¹ at 30 DAS	1,925	2,727	3,747	3,678	3,019
W ₄ : Application of 2,4-D at 0.5 kg ha ⁻¹ at 30 DAS	1,874	2,583	3,828	3,731	3,004
Mean	1,751	2,752	3,510	3,846	
SEm±	104				
CD (P = 5%)	301				

Table 10. Interaction of weed and nitrogen on biological yield of barley

Treatment	N ₀ : Control (0 kg N ha ⁻¹)	N ₁ : 30 kg N ha ⁻¹	N ₂ : 60 kg N ha ⁻¹	N ₃ : 90 kg N ha ⁻¹	Mean
W ₁ : Weedy check (no weed control)	2,866	4,570	5,692	6,762	4,972
W ₂ : Weed-free	3,290	5,207	6,688	6,934	5,530
W ₃ : Application of Metsulfuron-methyl at 4 g ha ⁻¹ at 30 DAS	3,097	5,186	6,770	6,820	5,468
W ₄ : Application of 2,4-D at 0.5 kg ha ⁻¹ at 30 DAS	3,328	4,438	6,890	6,820	5,369
Mean	3,145	4,850	6,510	6,834	
SEm±	164				
CD (P = 5%)	476				

Table 11. Interaction of weed and nitrogen on barley harvest index

Treatment	N ₀ : Control (0 kg N ha ⁻¹)	N ₁ : 30 kg N ha ⁻¹	N ₂ : 60 kg N ha ⁻¹	N ₃ : 90 kg N ha ⁻¹	Mean
W ₁ : Weedy check (no weed control)	35.67	33.50	34.91	40.27	36.09
W ₂ : Weed-free	43.68	53.26	44.62	48.14	47.43
W ₃ : Application of Metsulfuron-methyl at 4 g ha ⁻¹ at 30 DAS	43.71	42.35	45.67	47.33	44.77
W ₄ : Application of 2,4-D at 0.5 kg ha ⁻¹ at 30 DAS	41.16	43.37	49.23	45.86	44.91
Mean	41.06	43.12	43.61	45.40	
SEm±	1.94				
CD (P = 5%)	5.62				

Table 12. Effect of different treatments on nitrogen content (%) and uptake (kg ha⁻¹) in grain and straw of barley

Treatment	N content in grain (%)	N uptake by grain (kg ha ⁻¹)	N content in straw (%)	N uptake by straw (kg ha ⁻¹)	Total N uptake (kg ha ⁻¹)
Weed management					
W ₁ : Weedy check (no weed control)	1.48	29.45	0.43	11.02	40.47
W ₂ : Weed-free	1.48	37.45	0.45	14.83	52.28
W ₃ : Application of Metsulfuron-methyl at 4 g ha ⁻¹ at 30 DAS	1.48	36.77	0.44	13.58	50.35
W ₄ : Application of 2,4-D at 0.5 kg ha ⁻¹ at 30 DAS	1.50	33.91	0.43	13.15	47.06
SE(m)±	0.03	0.92	0.02	0.74	1.10
C.D. at 5%	NS	2.66	NS	2.15	3.21
Nitrogen management					
N ₀ : Control (0 kg N ha ⁻¹)	1.52	19.64	0.43	7.53	27.17
N ₁ : 30 kg N ha ⁻¹	1.43	29.50	0.42	11.73	41.02
N ₂ : 60 kg N ha ⁻¹	1.50	42.18	0.46	16.12	58.31
N ₃ : 90 kg N ha ⁻¹	1.49	46.29	0.45	17.40	63.66
SEm±	0.03	0.92	0.02	0.74	1.10
CD (P = 5%)	NS	2.66	NS	2.15	3.21

Table 13. Economics of barley as influenced by weed and nitrogen management

Treatment	Cost of cultivation (₹ ha ⁻¹)	Gross return (₹ ha ⁻¹)	Net return (₹ ha ⁻¹)	B:C ratio
Weed management				
W ₁ : Weedy check (no weed control)	31351	76447	45096	2.42
W ₂ : Weed-free	43351	97524	54173	2.24
W ₃ : Application of Metsulfuron-methyl at 4 g ha ⁻¹ at 30 DAS	32526	94241	61715	2.88
W ₄ : Application of 2,4-D at 0.5 kg ha ⁻¹ at 30 DAS	32531	87713	55182	2.68
SEm±	-	-	-	-
CD (P = 5%)	-	-	-	-
Nitrogen management				
N ₀ : Control (0 kg N ha ⁻¹)	33709	50476	16767	1.51
N ₁ : 30 kg N ha ⁻¹	34830	80261	45431	2.32
N ₂ : 60 kg N ha ⁻¹	35351	107308	71958	3.07
N ₃ : 90 kg N ha ⁻¹	35871	117881	82010	3.32
SEm±	-	-	-	-
CD (P = 5%)	-	-	-	-

4. Conclusion

The study demonstrated that weed management and nitrogen levels significantly influenced growth, yield, nutrient uptake, and economic returns of barley. Effective weed control treatments, particularly W₂ and W₃, improved crop performance by minimizing competition and enhancing resource use efficiency. Nitrogen application showed a positive response, with N₃ and N₂ enhancing productivity and nutrient uptake. The interaction of weed management and nitrogen levels indicated that combined application of effective weed control with optimum nitrogen is essential. Economically, W₃ and N₂ proved to be more viable, suggesting an efficient and sustainable approach for barley cultivation under semi-arid conditions. (recommendations).

Disclaimer (Artificial Intelligence)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

Competing Interests

Authors have declared that no competing interests exist.

References

- Al-Gburi, B. K. H., Lahmod, N. R., Al-Thabhwai, S. H., and Al-Falloorji, S. A. K. (2024). Weed control in barley (*hordeum vulgare*) via herbicides that inhibit als and accase with increased seeding rate. *Sabrao Journal of Breeding and Genetics*, 56(5): 2136-2142. <https://doi.org/10.54910/sabrao2024.56.5.36>
- Al-Jayashi, M. T., Leiby, H. R., and Fadel, H. A. (2024, July). A novel of Study Effect of Herbicides and Nitrogen Rate on Yield and Components of the Barley Crop (*Hordeum vulgare* L.) in Iraq. In *IOP Conference Series: Earth and Environmental Science* (Vol. 1371, No. 5, p. 052081). IOP Publishing. <https://doi.org/10.1088/1755-1315/1371/5/052081>
- Arshad, M. A., Abbas, R. N., Policarpo Tonelli, F. M., Baloch, R., Ul Haq, M. Z., Ahmad, A., and Alshaharni, M. O. (2025). Enhancing wheat yield and quality through optimized weed control timing and seeding density. *Archives of Agronomy and Soil Science*, 71(1): 1-21. <https://doi.org/10.1080/03650340.2025.2570466>
- Bogale, A. A., Kende, Z., Tarnawa, A., Miko, P., Birkás, M., Kovács, G. P., & Percze, A. (2025). Precision nutrient and soil tillage management for sustainable winter barley production (*Hordeum vulgare* L.) and tillage impact on soil CO₂ emission. *Agronomy*, 15(1), 2. <https://doi.org/10.3390/agronomy15010002>

- Cammarano, D., Holland, J., Gianinetti, A., Baronchelli, M., and Ronga, D. (2024). Impact of Nitrogen and Water on Barley Grain Yield and Malting Quality. *Journal of Soil Science and Plant Nutrition*, 24(4): 6718-6730. <https://doi.org/10.1007/s42729-024-01999-0>
- Dolijanović, Ž., Roljević Nikolić, S., Šeremešić, S., Jug, D., Biljić, M., Pešić, S., and Kovačević, D. (2025). Effects of Conservation Tillage and Nitrogen Management on Yield, Grain Quality, and Weed Infestation in Winter Wheat. *Agronomy*, 15(7): 1742. <https://doi.org/10.3390/agronomy15071742>
- Dorado, J., Barroso, J., Peña, J. M., Luna, I. M., and Fernández-Quintanilla, C. (2025). Weed control efficacy and economic profitability of three contrasting cropping systems in semiarid rainfed zones. *Weed Science*, 73(1): e75. <https://doi.org/10.1017/wsc.2025.10052>
- Gomez, K. A., and Gomez, A. A. (1984). *Statistical Procedures for Agricultural Research* (2nd ed.). New York: John Wiley and Sons. <https://doi.org/10.1002/9781118986400>
- Hirich, E. H., Idrissi, O., Boughlala, M., Ouabbo, H., El Gharass, O., Dahan, R., Echchgadda, G., Bamouh, A., Amghar, I., Blanco Moreno, J. M., and Cicek, H. (2026). Novel integration of forage cereal–legume mixtures in conservation agriculture enhances weed control and barley productivity in Moroccan drylands. *Journal of the Science of Food and Agriculture*. Advance online publication. <https://doi.org/10.1002/jsfa.70370>
- Jackson, M. L. (1973). *Soil chemical analysis*. New Delhi: Prentice Hall of India Pvt. Ltd. <https://www.phindia.com/Books/BookDetail/9788120302181/soil-chemical-analysis-jackson>
- Kantwa, C. R., Saras, P. K., Vyas, K. G., Chaudhari, H. L., Choudhary, R. R., Patel, S. A., Singh, S. R. K., and Patel, B. J. (2025). Effect of wheat varieties and integrated nutrient management practices on nutrient content, uptake and soil nutrient status. *Indian Journal of Agricultural Research*, 59(2): 239-243. <https://doi.org/10.18805/IJARE.A-6050>
- Kaur, A., Purewal, S. S., Phimolsiripol, Y., and Punia Bangar, S. (2024). Unraveling the hidden potential of barley (*Hordeum vulgare*): an important review. *Plants*, 13(17): 2421. <https://doi.org/10.3390/plants13172421>
- Kumar, B., Hasanain, M., Raza, M. D. B., Yadav, R. B., Singh, R. K., Mishra, R., Ahmad, G., and Singh, D. (2022). Effect of weed management practices on nutrient uptake and productivity of wheat (*Triticum aestivum*). *Indian Journal of Agricultural Sciences*, 92(3): 405-407. <https://doi.org/10.56093/ijas.v92i3.122726>
- Kumar, N., Nath, C. P., and Hazra, K. K. (2022). Weed management in pulse crops: Challenges and opportunities. *Indian Journal of Weed Science*, 54(4): 397-410. <https://doi.org/10.5958/0974-8164.2022.00071.5>
- Kumar, R., and Singh, U. P. (2018). Performance of zero-till wheat (*Triticum aestivum* L.) with residue and weed management techniques under rice-wheat cropping system. *Agricultural Science Digest-A Research Journal*, 38(2): 113-117. <https://doi.org/10.18805/ag.D-4860>
- Kumar, V., and Jha, P. (2017). Influence of nitrogen rate, seeding rate, and weed removal timing on weed interference in barley and effect of nitrogen, weed response to herbicides. *Weed Science*, 65(1): 189-201. <https://doi.org/10.1614/WS-D-16-00047.1>
- Loukakis, L. O., Giannoulis, K. D., Garoufali, E., Karaviti, T., Sotirakoglou, K., Kotoulas, V., Papastilianou, P., & Economou, G. (2025). Integration of biostimulants alongside various advanced nitrogen fertilization practices improve the yield, quality, and sustainability of malting barley in Mediterranean conditions. *Agronomy*, 15(10), 2417. <https://doi.org/10.3390/agronomy15102417>
- Mendera, J., Jalata, Z., and Wagari, A. (2024). Effects of Weeding Frequency and NPS Fertilizer Rates on Barley (*Hordeum vulgare* L.) Yield Performance, Western Ethiopia. *International Journal of Agricultural Science and Food Technology*, 10(3): 114-123. <https://doi.org/10.17352/2455-815X.000215>
- Naeem, M., Minhas, W. A., Hussain, S., Ul-Allah, S., Farooq, M., Farooq, S., and Hussain, M. (2022). Barley-based cropping systems and weed control strategies influence weed infestation, soil properties and barley productivity. *Agriculture*, 12(4): 487. <https://doi.org/10.3390/agriculture12040487>
- Olsen, S. R., Cole, C. V., Watanabe, F. S., and Dean, L. A. (1954). Estimation of available phosphorus in soils by extraction with sodium bicarbonate (Circular No. 939). U.S. Department of Agriculture. <https://naldc.nal.usda.gov/catalog/CAT87201169>
- Pooniya, V., Biswakarma, N., Parihar, C. M., Swarnalakshmi, K., Lama, A., Zhiipao, R. R., and Singh, K. (2021). Six years of conservation agriculture and nutrient management in maize–mustard rotation: Impact on soil properties, system productivity and profitability. *Field crops research*, 260: 108002. <https://doi.org/10.1016/j.fcr.2020.108002>
- Puniya, R., Bazaya, B. R., Kumar, A., Sharma, B. C., Nesar, N. A., Bochalya, R. S., and Mehta, S. (2023). Effect of residue and weed management practices on weed flora, yield, energetics, carbon footprint, economics

- and soil quality of zero tillage wheat. *Scientific Reports*, 13(1): 19311. <https://doi.org/10.1038/s41598-023-45488-3>
- Safdar, M. E., Qamar, R., Maqbool, R., Ali, H. H., Nadeem, M., Nadeem, M. A., and Ashraf, M. A. (2022). Management strategies for optimizing barley yield through row spacing under variable weed competition duration. *Pakistan Journal of Agricultural Sciences*, 59(3): 467-476. <https://doi.org/10.21162/PAKJAS/22.9887>
- Saikia, J., Saikia, L., Phookan, D. B., and Nath, D. J. (2018). Effect of biofertilizer consortium on yield, quality and soil health of french bean (*Phaseolus vulgaris* L.). *Legume Research*, 41(5): 755-758. <https://doi.org/10.18805/LR.D-4460>
- Scapino, M., Meloni, R., and Blandino, M. (2025). A comparison of the agronomic management of a winter barley hybrid and a conventional genotype: effect of the seeding rate, soil tillage and nitrogen fertilization. *Frontiers in Agronomy*, 7: 1546989. <https://doi.org/10.3389/fagro.2025.1546989>
- Shekhawat, K., Singh, V. K., Rathore, S. S., Raj, R., and Das, T. K. (2021). Achieving sustainability in food systems: addressing changing climate through real time nitrogen and weed management in a conservation agriculture-based maize–wheat system. *Sustainability*, 13(9): 5010. <https://doi.org/10.3390/su13095010>
- Shpanev, A. M., and Smuk, V. V. (2025). Efficiency of herbicide treatment of spring barley crops under conditions of moisture deficiency in the North-West of Russia. *Russian Agricultural Sciences*, 51(5): 859-864. <https://doi.org/10.3103/S1068367425701502>
- Singh, G., Chopra, P., Manuja, S., Kaur, N., and Singh, G. (2025). Evaluation of herbicides for enhancing yield and economic benefits in barley (*Hordeum vulgare* L.). *International Journal of Bio-Resource and Stress Management*, 16(12): 1-7. <https://doi.org/10.23910/1.2025.6441>
- Sisay, K., Adissie, S., Teshome, H., and Degu, A. (2025). Enhancing yield and yield components of food barley (*Hordeum vulgare* L.) through optimum nitrogen and phosphorus levels in Eastern Amhara, Ethiopia. *International Journal on Food, Agriculture and Natural Resources*, 6(2): 62-68. <https://doi.org/10.46676/ij-fanres.v6i2.487>
- Subbiah, B. V., & Asija, G. L. (1956). A Rapid Procedure for the Determination of Available Nitrogen in Soils. *Current Science*, 25(8), 259-260. <https://www.currentscience.ac.in/Volumes/25/08/0259.pdf>
- Sunil, C. M., Hanagi, C., Manjunath, B., Gayathri, B., and Ananthakumar, M. A. (2025). Impact of large-scale demonstration on productivity and profitability of blackgram under rainfed conditions of Chamaraajanagara district, Karnataka (Zone-6). *Legume Research*, 48(7): 1196-120. <https://doi.org/10.18805/LR-5074>

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2026): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://pr.sdiarticle5.com/review-history/157607>