



## Metalaxyl Induced Changes in Protein Metabolism During Germination of Maize

N. Shobha<sup>1</sup>, S. Likhitha<sup>2</sup> and G. Savitha<sup>3\*</sup>

<sup>1</sup>Department of Biochemistry, PRIST University, Vallam, Thanjavur-613413, Tamilnadu, India.

<sup>2</sup>Department of Biochemistry, Maharani's Science College for Women, J.L.B. Road, Mysore, India.

<sup>3</sup>Department of Chemistry, Associate professor, Maharani's Science College for Women, J.L.B. Road, Mysore, Karnataka, India.

### Authors' contributions

This work was carried out in collaboration between all the authors. Authors NS and SL performed the study, statistical analysis, wrote the protocol and wrote first draft of the manuscript with literature. Author GS designed the study, corrected and prepared a final draft of manuscript. All authors read and approved the final manuscript.

### Article Information

DOI: 10.9734/IJPSS/2015/12374

#### Editor(s):

(1) Susana Rodriguez-Couto, Unit of Environmental Engineering, Paseo Manuel Lardizabal, Donostia-San Sebastián, Spain.

#### Reviewers:

(1) Anonymous, Research Institute of Subtropical Forestry, China.

(2) B. Vidya Vardhini, Dept. Of Botany, Telangana University, Dichpally, Nizamabad-503322, India.

(3) Anonymous, University of Maringá, Brazil.

Complete Peer review History: <http://www.sciencedomain.org/review-history.php?iid=651&id=24&aid=6155>

Original Research Article

Received 29<sup>th</sup> June 2014  
Accepted 25<sup>th</sup> August 2014  
Published 19<sup>th</sup> September 2014

### ABSTRACT

The present work was carried out to investigate the effect of a systemic fungicide metalaxyl as a seed treatment on the protein metabolism of maize during early germination. The study was carried out for 7 days after soaking the seeds in different concentrations of metalaxyl and a control was maintained. Biochemical analyses of whole seedlings were done up to 7 days. Changes in the amount of total soluble protein, total free amino acids, and the activity of protease were performed using standard methods. The results indicate that protein content was maximum on 4<sup>th</sup> and 6<sup>th</sup> day of germination in both control and treated seeds. Metalaxyl treatment resulted in decreased protein content (20-50% inhibition) over the untreated in a dose dependent manner till the 4<sup>th</sup> day of germination. But as the growth proceeded a proportionate increase in the protein content was observed on the 6<sup>th</sup> day of germination in fungicide treated seeds compared to the control. The specific activity of protease was decreased by 46%, 81%, 88%, 97% and 100% with 1.5mg, 3mg, 4.5mg, 6mg and 7mg concentration of metalaxyl on 3<sup>rd</sup> day of germination when compared to the control. An increase in the total free amino acids occurred during the germination and maximum

\*Corresponding author: E-mail: [savitha\\_93@yahoo.com](mailto:savitha_93@yahoo.com);

free amino acids content was observed on the 5<sup>th</sup> day. Metalaxyl treatment resulted in a dose dependent depletion of free amino acids. A significant increase in proline content was found to occur in treated seeds. From the overall findings, the present study gives an insight into a protective effect of the system with an increased production of proline and decreased protease activity and free amino acids and at the same time a higher protein content during later stages of germination for a particular concentration may be due to the synthesis of novel proteins as a defense mechanism indicating the dual role of metalaxyl.

*Keywords: Metalaxyl; germination; protease; proteins; free amino acids; proline.*

## 1. INTRODUCTION

The mobilization of seed storage proteins represents one of the most important post-germination events in the growth and development of seedling. Proteolytic enzymes play a central role in the biochemical mechanism of germination [1]. During germination period, the storage proteins are degraded by a variety of proteases which convert the insoluble storage proteins into soluble peptides and these peptides are further hydrolyzed to free amino acids. These free amino acids are mobilized to the embryonic axis to support its growth and also to provide energy [2]. Downey mildew of maize (*Zea mays*. L) caused by *Peronosclerospora* is one of the most destructive diseases of this crop in the tropical Asian countries [3]. Integrated approaches are used to manage the disease including crop rotation, planting resistant cultivars and the application of fungicides like mancozeb and metalaxyl. Metalaxyl (methyl *N*-(methoxyacetyl)-*N*-(2,6-xylyl)-DL-alaninate) compounds have been widely used for the control of Downey mildew in a number of crops. Metalaxyl is a systemic fungicide acts by suppressing sporangial formation, mycelia growth and establishment of new infection [4]. Much of our knowledge of reserve mobilization and its control processes during germination is very well understood. In contrast, available information on metalaxyl is limited to its effect on the pathogen and not much data is available on the effect of metalaxyl in protein metabolism of maize seeds. Hence, the present work is an attempt for furthering our knowledge towards better understanding of the effect of metalaxyl in germinating maize seeds focusing on the protein metabolic changes during germination.

## 2. MATERIALS AND METHODS

### 2.1 Collection of Seeds and Treatments

Maize seeds were procured from VC farm, University of Agriculture Science, Mandya, Karnataka. All the chemicals were purchased

from SLR and MERCK and the chemicals were of analytical grade. Seeds were surface sterilized with 0.1% mercuric chloride for 10 minutes and repeatedly washed with distilled water for 4-5 times. Seeds of uniform size were selected and soaked for 24 hours in distilled water (control) and with different concentrations (mg/g) of metalaxyl, 1.5, 3, 4.5, 6 and 7mg/g of the seeds (1:5 weight/volume) for 24 hours. Five seeds in triplicate were placed on petridish with 8-10 layer of soaked filter paper and incubated at 25<sup>0</sup>C both in light and dark condition. Uniform seedlings were selected and processed for further studies. Everyday filter paper was wetted with 10 ml of distilled water [5].

### 2.2 Preparation of Crude Extract

Around 1-2g of seedling treated with different concentrations of the metalaxyl and the untreated seeds (control) were taken each day, (up to 7 days) homogenized in ice cold saline (5ml) using pestle and mortar. The solution was centrifuged at 10000g for 10 minutes and the supernatant was used for further analysis.

### 2.3 Biochemical Studies

Protein was estimated as described by Lowry et al. [6] using BSA as standard. The activity of protease was determined following the procedure of Kunitz [7]. Total free amino acids were extracted and determined following the method of Sugano et al. [8]. Free proline content was estimated following the method of Bates et al. [9].

### 2.4 Statistical Analysis

The data are expressed as the mean  $\pm$  SEM analyzed by one-way analysis of variance (ANOVA) and Dunnett's *t*-test was used as the test of significance. P value <0.05 was considered as the minimum level of significance. All statistical tests were carried out using SPSS statistical software.

### 3. RESULTS AND DISCUSSION

#### 3.1 Total Protein


The protein content during different days of germination in control and fungicide treated seedlings is presented in Table 1. The protein content was maximum on 4<sup>th</sup> and 6<sup>th</sup> day of germination in both control and treated seeds. Metalaxyl treatment resulted in the decreased protein content (20-50% inhibition) over the untreated in a dose dependent manner till the 4<sup>th</sup> day of germination. The treated seeds showed similar pattern of protein content on different days of germination. However an increase in the protein content was seen on the 6<sup>th</sup> day of germination in treated seeds.

In the present study the protein content in germinated seeds were found to be increased due to the mobilization of storage nitrogen for the production of protein needed for the development of the young seedling and the observation was well in agreement with the studies conducted in germinated seeds [10]. A decline in the protein content was observed in treated seeds till the 4<sup>th</sup> day in a dose dependent fashion. The decrease in the protein content in fungicide treated maize may be due to osmotic shock effect by the fungicides which results in the release of protein and loss of membrane transport ability [11]. It has been suggested that the toxicant produced in the treated seeds may inhibit the protein synthesis by binding to the larger ribosomal subunit inducing change in the enzyme system ceasing ATP and NADP formation. The results of

our studies were in parallel, as the concentration of fungicide increases, the amount of protein gradually decreased. But as the growth proceeded a proportionate increase in the protein content was observed on the 6<sup>th</sup> day of germination in fungicide treated seeds compared to the control. The enhanced protein content may be attributed to the fact that as growth proceeds, metalaxyl may induce the expression of many defense related genes in plants which results in the synthesis of novel proteins. The results of our study correlates with the recent proteomic analysis which revealed the occurrence of novel proteins in several plant species under heavy metal stress [12].

#### 3.2 Activity of Protease

Protease is a hydrolytic enzyme which acts on proteinaceous substances to produce amino acids and amides. The specific activities of protease during germination of maize seeds treated with metalaxyl are shown in Fig 1a. Both control and treated seeds showed increasing specific activity up to 3<sup>rd</sup> day of germination and decreased then onwards. Specific activity of protease in maize seedlings at different concentration of metalaxyl on 3<sup>rd</sup> day were found to be 27, 9.1, 6, 1.6 and 0.3  $\times 10^{-7}$  mM whereas in control it is found to be 50  $\times 10^{-7}$  mM. The specific activity was decreased by 46%, 81%, 88%, 97%, and 100% with 1.5 mg, 3 mg, 4.5 mg, 6 mg and 7 mg concentration of Metalaxyl on 3<sup>rd</sup> day of germination when compared to the control.



**Fig. 1a. Effect of metalaxyl on the \*Specific activity of protease in the seedlings of maize**  
\*Specific activity of Protease is expressed as  $10^{-7}$  mM of amino acids released/mg of protein/min




Fig. 1b. Effect of metalaxyl on the activity of protease in the seedlings of maize

Table 1. Effect of metalaxyl on the total protein (mg/g) in the seedlings of maize

Concentration of metalaxyl (mg/g)	Seedling age in days								
	0	1	2	3	4	5	6	7	
Control	12.6±0.9	24.2±1	26.69±1.4	10.75±0.6	42.48±2.1	29.52±1.4	57±3.1	50.65±1	
1.5	10.7±0.7	22.82±1	24.45±1.5	9.21±0.3	34.09±1.4	21.54±1.2	76.4±2	60.49±1.9	
3.0	13.1±0.5	25.43±1.2	25.91±1.2	9.71±0.4	30.8±1.2	33.15±1.5	96.06±2.4	31.96±1.1	
4.5	34.4±1.2	27.61±1	25.69±0.4	2.71±0.1	26.78±0.9	33.69±1.6	67.51±2.3	52.4±2.1	
6.0	12.6±1	25.18±1.1	12.43±0.3	12.14±0.6	20.16±0.5	17.87±0.9	63.19±2.3	22.33±0.8	
7.0	10±0.5	21.12±0.9	23.15±1.1	7.65±0.3	25.45±0.7	15.91±0.7	47.52±1.2	31.21±0.9	

Increasing in proteolytic activity with concomitant reserve protein depletion agrees with the findings of earlier studies on other seeds; *Phaseolus vulgaris* [13], *Lupinus albus* [14] *Vicia sativa* [15] and *acrotyloma uniflorum* [16]. The result from our study is not parallel with the above data as there is a dose dependent inhibition of protease activity in metalaxyl treated maize seeds on 3<sup>rd</sup> day of germination. Impairment of proteasome functionality and decreased protease activities seems to be a common feature involved in metal toxicity in plants [17].

### 3.3 Free Amino Acid

The amino acid content in control and treated seeds during different days of germination is presented in the Fig. 2. A gradual increase in the amino acid content is seen and the maximum content of amino acid ( $41 \times 10^{-1}$  mg/g) was seen

on the 5<sup>th</sup> day in control and a gradual decrease was seen till the 7<sup>th</sup> day of germination. The free amino acid content was decreased effectively with the increasing concentration of fungicides and the maximum inhibition was seen with the highest concentration (7 mg/g). Only 30% of free amino acids were present in that concentration. An increase in the total free amino acids occurred during the germination and maximum free amino acids content was observed on the 5<sup>th</sup> day. Metalaxyl treatment resulted in a dose dependent depletion of free amino acids. There was no change in the pattern of amino acid profile on different days of germination in both treated and control seeds. These results suggests, the protective effect of amino acids against unfavorable condition, increasing the tendency of the system to maintain the homeostasis.

### 3.4 Proline content

Proline content in control and treated seeds are presented in the Fig. 3. The proline content increased in treated seeds and maximum proline content was seen with 3mg/g and in 6mg/g concentration of metalaxyl (0.797 mg/g, 0.705 mg/g) on 5<sup>th</sup> day of germination and decreased thereafter. The proline content was lowest in both control (0.043mg/g) on 5<sup>th</sup> day and in treated (7 mg/g) it was 0.06mg/g on 7<sup>th</sup> day. A significant increase in proline content was found to occur in treated seeds.

Proline is considered as an antioxidant free radical scavenging and biochemical indicator of

stress. It is considered as a metabolic measure of abiotic stress [18]. In the present work the stress caused by metalaxyl resulted in an increase in proline levels. In general water stress has been known to increase free proline in leaves [19]. Plant growing in stress condition need to produce specific proteins having higher proline content (hydroxyproline rich glycopeptide or proline rich glycopeptides) [20]. The high content of free proline is also observed in different cultivars of wheat under salt stress [21]. Effect of this change is subsequently contributed to higher production of glutamic acid which stressed plants need to synthesize phytochelatin [22].




Fig. 2. Effect of metalaxyl on the amino acid content (10<sup>-1</sup> mg/g) in the seedlings of maize




Fig. 3. Effect of metalaxyl on the proline content (mg/g) in the seedlings of maize

#### 4. CONCLUSION

The present study gives an insight into a protective effect of the system with an increased production of proline and decreased protease activity and free amino acids. At the same time, increased protein content was observed during the later stages of germination for a particular concentration of metalaxyl. This may be due to the synthesis of novel proteins involved in the defense mechanism and regulation process. This indicates the dual role of metalaxyl. Further work is in progress in understanding the structural and functional aspects of these proteins.

#### ACKNOWLEDGEMENT

We express our sincere thanks to Dr. N. G. Kademani, Head, Department of Studies in Biochemistry, Maharani's Science College for Women, Mysore, for providing the laboratory facility for this study.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

- Muntz K, Belozersky MA, Dunaevsky YE, Schlereth A, Tiedemann J. Stored proteinases and the initiation of storage protein mobilization in seeds during germination and seedling growth. *Journal of Experimental Botany*. 2001;52:1741–1752.
- Bewley JD, Black M. *Physiology and Biochemistry of Seeds in Relation to Germination*. Springer – Verlag, Berlin. 1983;1.
- Bonde MR. Epidemiology of downy mildew diseases of maize, sorghum and pearl millet. *Tropical Pest Management*. 1982;28(1):49-60.
- Fisher DJ, Hayes AL. Mode of action of the systemic fungicides furalaxyl, metalaxyl and ofurace. *Pestic. Sci*. 1982;13:330–339. DOI: 10.1002/ps.2780130316.
- ISTA. International rules for seed testing. In: Daper SR.(Ed), *Rules*; 2003. International seed testing association, Zurich, Switzerland, 1-520.
- Lowry OH, Rosenbrought NJ, Farr FL, Randall RJ. Protein measurement with Folin phenol reagent. *J. Biol. Chem*. 1951;193:265-275.
- Kunitz M. isolation of a crystalline protein compound of trypsin and of soybean trypsin-Inhibitor. *J. Gen. Physiol*. 1947;30(4):311-320.
- Sugano N, Tanaka T, Yamamoto E, Nishi A Phenylalanine ammonia lyase in carrot cells in suspension cultures. *Phytochemistry*. 1975;14:2435–2440.
- Bates LS, Waldran RP, Teare ID. Rapid determination of free proline for water-stress studies. *Plant Soil*. 1973;39:205-207.
- Gernah DI, Ariahe CC, Ingbian EK. Effects of malting and lactic acid fermentation on some chemicals and functional properties of maize. *Am. J. Food. Technol*. 2011;6:404-412.
- Amar L, Reinhold L. Loss of membrane transport ability in leaf cells and release of protein as a result of an osmotic shock. *Plant Physiol*. 1973;51:620-625.
- Elisabetta Gianazza, Robin Wait, Andrea Sozzi, Simona Regondi, Dolores Sac, Massimo Labra and Elisabetta Agradi. Growth and protein profile changes in *Lepidium sativum L.* plantlets exposed to cadmium. *Environmental and Experimental Botany*. 2007;59:179–187.
- Taneyama M, Okamoto T, Yamauchi D, Minamikawa T. Development of endopeptidase activity in cotyledons of *Vigna mungo* seedling: Effects of exogenously applied end products and plant hormones. *Plant Cell and Physiology*. 1996;37:19–26.
- Ferreira RB, Malo TS, Teixeira AN. Catabolism of the seed storage proteins from *Lupinus albus*: Fate of globulins during germination and seedling growth. *Australian Journal of Plant Physiology*. 1995;22:373–381.
- Schlereth A, Standhardt D, Mock HP, Muntz K. Stored cysteine proteinases start globulin breakdown in protein bodies of embryonic axis and cotyledons of germinating vetch (*Vicia sativa L.*) seeds. *Planta*. 2001;212:718–727.
- Rajeswari J, Ramakrishna Rao P. Storage protein degradation in germinating horse gram seeds. *Indian J Plant Physiol*. 2002;7:314-320.
- Wang C, Tian Y, Wang X, Geng J, Jiang J, Yu H, Wang C. Lead-contaminated soil induced oxidative stress, defense

- response and its indicative biomarkers in roots of *Vicia faba* seedlings. *Ecotoxicology*. 2010;19(6):1130-9.
18. Andurwulan N, Fardiaz D, Wattimena GA, Setty K. Antioxidant activity associated with lipid and phenolic mobilization during seed germination of *Pangium edule* Reinw. *J Agri Food Chem*. 1999;47:3158–63.
  19. Stewart CK, Lee JA. The role of proline accumulation in halophytes. *Planta*. 1974;120:1279-289.
  20. Ueda A, Yamamoto-Yamane Y, Takabe T. Salt stress enhances proline utilization in the apical region of barley roots. *Biochemical and Biophysical Research Communication*. 2007;355:61–66.
  21. Poustini K, Siosemardeh A, Ranjbar M. Proline accumulation as a response to salt stress in 30 wheat (*Triticum aestivum* L.) cultivars differing in salt tolerance, *Genet. Resour. Crop. Evol.* 2007;54(5): 925-934.
  22. Pavlíková D, Pavlík M, Stazková L, Motyka V, Száková J, Tlustoš P, Balík J. Glutamate kinase as a potential biomarker of heavy metal stress in plants. *Ecotoxicology and Environmental Safety*. 2008;70:223–230.

© 2015 Shobha et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

The peer review history for this paper can be accessed here:  
<http://www.sciencedomain.org/review-history.php?iid=651&id=24&aid=6155>